



ISOTOX[®] Lindane Spray No. 200

[INSECTICIDE]

NET CONTENTS
1 GALLON LIQUID

Active Ingredients	By Wt.
Lindane-Gamma Isomer of Benzene Hexachloride	20%
Aromatic Petroleum Derivative Solvent	59%
Inert Ingredients	21%
Lindane per gal. — 1.65 pounds	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

USE DESIGNATED AMOUNT IN 100 GALLONS WATER UNLESS OTHERWISE SPECIFIED. Repeat applications as necessary.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KALE, COLLARDS: Aphids, Thrips—1 pt. (Use up to 150 gals. diluted spray per acre). Do not apply to Cabbage, Broccoli, Brussels Sprouts, Cauliflower, and Kale after edible parts start to form. Apply to Collards only at seedling stage.

CELERY: Aphids, Thrips, Tarnished Plant Bugs and Certain Types of Lepidopterous Larvae—1 pt. (Use up to 150 gals. diluted spray per acre). Do not apply after the celery is half-grown or after it begins to bunch.

CUCUMBERS, MELONS, SQUASH, SUMMER SQUASH, PUMPKINS: Cucumber Beetles, Thrips— $\frac{1}{4}$ to $\frac{1}{2}$ pt. (Use up to 400 gals. diluted spray per acre). Squash Bugs on Squash—1 to 1½ pts. (Use up to 150 gals. diluted spray per acre). Aphids—1 pt. (Use up to 200 gals. diluted spray per acre). Do not apply to Cucumbers, Squash and Summer Squash within 1 day of harvest.




READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed. Avoid breathing spray mist. Avoid skin contact. May be absorbed through skin. In case of contact, wash immediately with soap and water. Do not store or transport near feed or food. Do not use on household pets or humans. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Do not apply to edible crops within 30 days of harvest. Do not apply to foliage of white or sweet potatoes. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption. When container is empty, immediately wash thoroughly and destroy. Never re-use.

Toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not contaminate any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not use where run-off is a frequent problem.



239-2363

GRAIN AND OTHER STORAGE BINS FOR GRAIN AND STORED PRODUCTS INSECTS (exposed stages): Saw-Toothed Grain Beetle, Granary Weevil, Confused Flour Beetle, Corn or Rice Weevil, Bean Weevil, Angoumois Grain Moth, Indian Meal Moth—2½ gals. per 100 gals. water (0.5% lindane). Apply at 2 gals. diluted spray per 1000 sq. ft. of bin area. Treat empty storage bins prior to filling.

GREENHOUSE NURSERY PLANTS: Aphids, Thrips, Loopers, Leaf-rollers, Plant Bug, Whitefly, Psyllid, Leaf Miners, Flea Beetles, Diabrotica Beetles—1 pt. **CAMILLIAS, AZALEAS, RHODODENDRONS**—Fuller Rose Weevil, Brachyrhinus, Cucumber Beetle Larvae on Roots—1 pt. Spray to saturate soil around plants in bed. **For Potted Plants:** Spray soil surface lightly, comparable to a light rain.

PINE BARK BEETLES: 1 quart in 10 gals. kerosene or No. 2 fuel oil and spray trunks thoroughly, or use 1 quart in 8 gals. water and wet trunks to run-off.

SOIL INSECTS: Wireworms, Cutworms, White Grubs, Diabrotica Larvae and Certain Other Soil Insects on Celery, Cucumbers, Kale, Lettuce, Melons, Pumpkins, Spinach, Tomatoes—**Pre-Plant Treatment**—2 to 2½ pts. per acre in water to cover. Spray directly on soil, then disc as soon as practicable to give thorough distribution in upper 4 to 6 inches of soil. Do not apply to root crops. Do not use in soil to be planted to root crops (except potatoes) within 1 year. Do not use in soil to be planted to potatoes within 2 years. **Garden Centipede** (Symphylid) on listed crops—Use 3 quarts per acre in water to cover (increase dosage to 5 quarts per acre in heavy infestations). Application same as above. Do not use this dosage on root crops, or in soils or on other crops grown in rotation with root crops, including peanuts.

SPINACH (Seedling stage only): Aphids, Leaf Miners—1 pt. (Use up to 200 gals. diluted spray per acre.)

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

CHEVRON CHEMICAL COMPANY

FORM 8145

EPA Reg. No. 239-2363

ORTHO DIVISION — San Francisco, California 94119

Product 1127

Made in U. S. A.

RICHMOND, CALIFORNIA

HONOLULU, HAWAII

SAN FRANCISCO, CALIFORNIA



DES MOINES, IOWA

ORLANDO, FLORIDA

PERTH AMBOY, NEW JERSEY