



ORTHO Diazinon Insect Spray

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING: Causes eye injury. Harmful if swallowed or absorbed through the skin. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin or clothing. Do not use on household pets or humans. Keep children and pets away from treated area until completely dry. Keep out of reach of domestic animals. **STATEMENT OF PRACTICAL TREATMENT:** In case of eye contact, wash eyes immediately with fresh water for 15 minutes and see a doctor. For skin contact, wash thoroughly with soap and water. Remove and launder contaminated clothing. If swallowed, DO NOT make person vomit. Call a doctor immediately. **Note to Physicians:** Emergency Information: call (415) 233-3737. This product contains a light petroleum solvent and may present an aspiration problem. Diazinon is an organophosphorus cholinesterase inhibitor. If signs and symptoms of cholinesterase inhibition develop, atropine is antidotal. 2-PAM may also be given in conjunction with atropine but should never be used alone.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate ornamental fishponds. Do not contaminate water, by cleaning of equipment or disposal of wastes. This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on treated areas. Do not apply when bees are actively visiting the crop. Application should be timed to provide the maximum possible interval between treatment and the next period of bee activity. **PHYSICAL OR CHEMICAL HAZARDS:** Do not use or store near heat or open flame.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

LAWNS: To Control: Ants, Armyworms, Bermuda Mites, Chiggers, Chinch Bugs, Clover Mites, Crickets, Cutworms, Dichondra Flea Beetle, Earwigs, European Crane Fly (larvae), Fleas, Lawn Billbugs, Lawn Moths, Leafhoppers, Sod Webworm, Brown Dog Ticks. Apply 4 fl oz (1/2 cup) in 15 gals. water to cover 500 sq. ft.

Notes: Thoroughly wet down grass a few hours before applying. For Lawn Billbugs, water lawn thoroughly immediately after treatment. To Control: Japanese Beetle Grubs, May Beetle Larvae (White Grubs), Mole Crickets. Apply 5 fl oz (1/2 tsp) in 15 gals. water to cover 500 sq. ft. Water lawn thoroughly after treatment.

Notes: Treat when damage to lawn is first noticed and grubs for European Crane Fly (larvae) are found in the soil or consult your local Agricultural Extension Service Specialist for proper time to treat in your area. For Mole Crickets, apply in May or June when pests are small.

LAWNS AND OTHER RECREATIONAL AREAS: To Aid in Control: Fire Ants. Apply 1 fl oz (2 Tbsp.) in a minimum of 1 gal. of water as a drench to mounds. **Notes:** Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Reapply as necessary. For best results, apply when temperatures are 65-80° F or in early morning or late evening hours. Treat new mounds as they appear.

FRUITS, VEGETABLES, ROSES, FLOWERS, SHRUBS & SHADE TREES

To Control: Aphids, Bagworms, Bean Beetles, Cabbageworms, Codling Moth, Cucumber Beetles, Cutworms, Flea Beetles, Holly Bud Moth, Leafhoppers, Leafminers, Lacebugs, Leafrollers, Mealybugs, Mites, Scale Crawlers, Thrips, Whiteflies. Apply 2 tsp (1/4 fl oz) per gal. water. Thorough coverage is important. Try to hit undersides of leaves and penetrate dense foliage to wet leaves and branches to dripping point. Repeat if needed.

FRUITS: Apples (14), Peaches (20), Oranges (21), Plums (10), Grapes (10), Cherries (10), Strawberries (5). **Note:** Start spraying after blooming and repeat at 10-14 day intervals.

VEGETABLES: Tomatoes (1), Beans (7), Cabbage (7), Carrots (10), Peppers (5), Potatoes (35), Onions (10), Cucumbers (7), Cantaloupes (3). **Note:** Apply at 7-10 day intervals as needed.

Note: Numbers in () indicate days required between last application and harvest.

ROSES & FLOWERS: Azaleas, Camellias, Chrysanthemums, Carnations, Gladioli, Pyracantha, Rhododendrons.

SHRUBS & SHADE TREES: Birch, Boxwood, Hawthorn, Holly, Lilac, Locust, Maple, Pine, Spruce, Sycamore, Willow.

Note: Do not use on certain ferns, poinsettia, hibiscus, papaya, pilea, gardenia, and orchids since injury may occur.

To Control: **Mediterranean Fruit Fly (larvae):** Apply 4 fl oz (1/2 cup) in 15 gals. water to cover 500 sq. ft. of soil beneath host plants listed above. Drench all soil within dripline. Make 3 applications at 14-16 day intervals as needed.

Combination Spray with Fungicide on Roses: ORTHO FUNGINEX® Rose Disease Control may be mixed with this product when listed insect(s) and disease(s) are found on roses. Follow directions and precautions on both labels.

NUISANCE PESTS: In outside areas

To Control: Ants, Box Elder Bugs, Chiggers, Clover Mites, Cockroaches, Earwigs, Fleas, Flies, Millipedes, Scorpions, Sowbugs (Pillbugs), Spiders (including Black Widow Spiders), Brown Dog Ticks. Apply 8 fl oz in 3 gals. water. Spray a 5 ft. band of soil around the house next to the foundation to a height of 2 to 3 feet. Thoroughly spray windows and door screens, sills, foundations, porch and patio walls and ceilings, entrance ways, walks, firewood piles, garbage cans, tree trunks, and into cracks and other places where insects can hide. **DO NOT USE IN HOME.** Do not treat plants or animals with this nuisance pest rate. Do not burn sprayed firewood within 2 weeks following treatment.

STORAGE AND DISPOSAL

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area. Do not store diluted spray. Keep tightly capped.

DISPOSAL: PRODUCT — Partially filled bottle may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash. **CONTAINER** — Do not reuse empty bottle. Rinse thoroughly before discarding in trash.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

FUNGINEX® Reg. TM of CILAMERCK GmbH Co. for Trifluine Fungicide.

Chevron Chemical Company © 1983

Ortho Consumer Products Division

San Francisco CA 94120 7144

Form 8130 M2 Product 1000 49 Made in U.S.A.

EPA Reg. No. 239-2364 AA

EPA Est. 239 MD 1

ORTHO


Kills up to 16 Common Garden Insects

Diazinon Insect Spray

CONTROLS LAWN INSECTS
Ants, Fleas, Sod Webworms,
Brown Dog Ticks, White Grubs
Kills Black Widow Spiders

CONTROLS GARDEN INSECTS
Aphids, Cucumber Beetles,
Cabbageworms, Codling Moth,
Whiteflies


This product must not be used on golf courses and sod farms.



Active Ingredients	By Wt
Diazinon (0.0 diethyl 0 (2 isopropyl 6 methyl 4 pyrimidinyl) phosphorothioate)	25%
Aromatic Petroleum	57%
Derivative Solvent	18%
Inert ingredients	

Keep out of reach of children
WARNING
See back panel for additional precautionary statements.

NET CONTENTS 1 QT.



10/2

ORTHO Diazinon Insect Spray

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING: Causes eye injury. Harmful if swallowed or absorbed through the skin. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin or clothing. Do not use on household pets or humans. Keep children and pets away from treated area until completely dry. Keep out of reach of domestic animals. **STATEMENT OF PRACTICAL TREATMENT:** In case of eye contact, wash eyes immediately with fresh water for 15 minutes and see a doctor for skin contact, wash thoroughly with soap and water. Remove and launder contaminated clothing. If swallowed, give water or milk to drink and telephone for medical advice. DO NOT make person vomit unless directed to do so by medical personnel. If medical advice cannot be obtained, then take person and product container to the nearest medical emergency treatment center or hospital. **Note to Physicians:** Emergency Information—call (475) 233-3737. This product contains a light petroleum solvent and may present an aspiration problem. Diazinon is an organophosphorus cholinesterase inhibitor. It is toxic and symptoms of cholinesterase inhibition develop; atropine is antidotal. 2 PAM may also be given in conjunction with atropine but should never be used alone.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water. Do not contaminate ornamental fishponds. Do not contaminate water by cleaning of equipment or disposal of wastes. This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on treated areas. Do not apply when bees are actively visiting the crop. Application should be timed to provide the maximum possible interval between treatment and the next period of bee activity. **PHYSICAL OR CHEMICAL HAZARDS:** Do not use or store near heat or open flame.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

LAWNS: To Control: Ants, Armyworms, Bermuda Mites, Chiggers, Chinch Bugs, Clover Mites, Crickets, Cutworms, Dichondra Flea Beetle, Earwigs, European Crane Fly (larvae), Fleas, Lawn Billbugs, Lawn Moths, Leafhoppers, Sod Webworm, Brown Dog Ticks. Apply 4 fl oz (1/2 cup) in 15 gals. water to cover 500 sq ft.

Notes: Thoroughly wet down grass a few hours before applying. For Lawn Billbugs, water lawn thoroughly immediately after treatment. To Control: Japanese Beetle Grubs, May Beetle Larvae (White Grubs), Mole Crickets. Apply 5 fl oz (10 Tbsp) in 15 gals. water to cover 500 sq ft. Water lawn thoroughly after treatment.

Notes: Treat when damage to lawn is first noticed and grubs (or European Crane Fly (larvae)) are found in the soil or consult your local Agricultural Extension Service Specialist for proper time to treat in your area. For Mole Crickets, apply in May or June when pests are small.

LAWNS AND OTHER RECREATIONAL AREAS: To Aid in Control: Fire Ants—Apply 1 fl oz (2 Tbsp) in a minimum of 1 gal. of water as a drench to mounds. **Notes:** Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Reapply as necessary for best results; apply when temperatures are 65-80°F or in early morning or late evening hours. Treat new mounds as they appear.

FRUITS, VEGETABLES, ROSES, FLOWERS, SHRUBS & SHADE TREES

To Control: Aphids, Bagworms, Bean Beetles, Cabbageworms, Codling Moth, Cucumber Beetles, Cutworms, Flea Beetles, Holly Bud Moth, Leafhoppers, Leafminers, Lacebugs, Leafrollers, Mealybugs, Mites, Scale Crawlers, Thrips, Whiteflies. Apply 2 tsp (1/4 fl oz) per gal. water. Thorough coverage is important. Try to hit undersides of leaves and penetrate dense foliage to wet leaves and branches to dripping point. Repeat if needed.

FRUITS: Apples (14), Apricots (10), Canberries (7), Cherries (10), Grapes (10), Nectarines (10), Oranges (7), Peaches (20), Pears (14), Plums (10), Strawberries (5). **Note:** Start spraying after blooming and repeat at 10-14 day intervals.

VEGETABLES: Beans (7), Broccoli (5), Cabbage (7), Cantaloupes (3), Carrots (10), Cauliflower (5), Cucumbers (7), Lettuce (10), Onions (10), Peas (1), Peppers (5), Potatoes (35), Radishes (10), Sweet Corn (1), Summer Squash (7), Tomatoes (1). **Note:** Apply at 7-10 day intervals as needed.

Notes: Numbers in parentheses indicate days required between last application and harvest.

ROSES & FLOWERS: Azaleas, Camellias, Chrysanthemums, Carnations, Gladioli, Pyracantha, Rhododendrons.

SHRUBS & SHADE TREES: Birch, Boxwood, Hawthorn, Holly, Lilac, Locust, Maple, Pine, Spruce, Sycamore, Willow.

Note: Do not use on certain ferns: poinsettia, hibiscus, papaya, plea, gardenia and orchids since injury may occur.

To Control: Mediterranean Fruit Fly (larvae). Apply 4 fl oz (1/2 cup) in 15 gals. water to cover 500 sq ft. or soil beneath host plants listed above. Drench all soil within drip line. Make 3 applications at 14-16 day intervals as needed.

Combination Spray with Fungicide on Roses: ORTHO FUNGIX® Rose Disease Control may be mixed with this product when listed insect(s) and disease(s) are found on roses. Follow directions and precautions on both labels.

NUISANCE PESTS: In outside areas.

To Control: Ants, Box Elder Bugs, Chiggers, Clover Mites, Cockroaches, Earwigs, Fleas, Flies, Millipedes, Scorpions, Sowbugs (Pillbugs), Spiders (including Black Widow Spiders), Brown Dog Ticks. Apply 8 fl oz in 3 gals. water. Spray a 5 ft. band of soil around the house next to the foundation to a height of 2 to 3 feet. Thoroughly spray windows and door screens, sills, foundations, porch and patio walls and ceilings, entrance ways, walks, firewood piles, garbage cans, tree trunks, and into cracks and other places where insects can hide. **DO NOT USE IN HOME.** Do not treat plants or animals with this nuisance pest rate. Do not burn sprayed firewood within 2 weeks following treatment.

STORAGE AND DISPOSAL

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool dry place, preferably in a locked storage area. Do not store diluted spray. Keep tightly capped.

DISPOSAL: PRODUCT—Partially filled bottle may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash. **CONTAINER:**—Do not reuse empty bottle. Rinse thoroughly before discarding in trash.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions. FUNGIX® Reg. TM of CEMERICK GmbH Co. for Triflorine Fungicide.

Chevron Chemical Company © 1983
Ortho Consumer Products Division
San Francisco CA 94120-7144
R Form B130 N7 Product 1000 49 Made in U.S.A.
EPA Reg. No. 239-2364-5A
EPA Est. 239-MO-1

ORTHO


Makes
up to
35 Gallons
Diluted
Spray

Diazinon Insect Spray

CONTROLS LAWN INSECTS
Ants, Fleas, Sod Webworms,
Brown Dog Ticks, White Grubs
Kills Black Widow Spiders

CONTROLS GARDEN INSECTS
Aphids, Cucumber Beetles,
Cabbageworms, Codling Moth,
Whiteflies

This product must not be used on golf courses and sod farms.



Contains Wetting Agent for More Effective Coverage.

← Copy A

Active Ingredients
Diazinon (0,0-diethyl-O (2-isopropyl-6-ethyl-4-pyrimidinyl) phosphorothioate)
Inert Ingredients
Contains petroleum distillates

By Vol.

25-75

Keep out of reach of children

WARNING

See back panel for additional
precautionary statements.


CONFIDENTIAL

PROPERTY OF

CHEVRON CHEMICAL COMPANY

FORM ACCEPTED BY EPA

NET CONTENTS 1 QT.



20/2