

PM 15
239-2364

ORTHO Diazinon Insect Spray
EPA Reg. No. 239-2364

COPY A

flush eyes with plenty of water and see a doctor.

COPY B


Contains ORTHEXTM, a special additive for better insect control. ORTHEXTM makes your spray spread more evenly and stick better to cover insects on smooth, rough or waxy plant surfaces.

COPY C


Do not contaminate water when disposing of equipment washwaters.

COPY D

ORTHEXTM



CONFIDENTIAL
PROPERTY OF
CHEVRON CHEMICAL COMPANY
UNTIL ACCEPTED BY EPA



10/2-

ORTHO Diazinon Insect S

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING: Causes eye injury. Harmful if swallowed or absorbed through the skin. Do not handle. Avoid contact with skin or clothing. Do not use on household pets or humans. Keep completely dry. Keep out of reach of domestic animals. **STATEMENT OF PRACTICAL TREATMENT:** If swallowed, give water or milk to drink and telephone for medical advice. DO NOT use medical personnel. If medical advice cannot be obtained, then take person and product to a center or hospital. **Note to Physicians:** Emergency Information — call (415) 233-3737. This product presents an aspiration problem. Diazinon is an organophosphorus cholinesterase inhibitor. Develop atropine is antidotal. 2 PAM may also be given in conjunction with atropine but is not highly toxic to bees exposed to direct treatment or to residues remaining on treated area. **ENVIRONMENTAL HAZARDS:** This product is toxic to fish and wildlife. Bees feeding on treated water. Do not contaminate ornamental fishponds. Do not use mammalian water by cleaning or water. Application should be timed to provide the maximum possible interval between treatment. **OR CHEMICAL HAZARDS:** Do not use or store near heat or open flame.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS.**

LAWNS: To Control: Ants, Armyworms, Bermuda Mites, Chiggers, Chinch Bugs, Clover Mites, Earwigs, European Crane Fly (larvae), Fleas, Lawn Billbugs, Lawn Moths, Leafhoppers, etc. (1/2 cup) in 15 gals. water to cover 500 sq. ft.

Notes: Thoroughly wet down grass a few hours before applying. For Lawn Billbugs, wet down grass. To Control: Japanese Beetle Grubs, May Beetle Larvae (White Grubs), Mole Crickets — 500 Sq. Ft. Water lawn thoroughly after treatment.

Notes: Treat when damage to lawn is first noticed and grubs (or European Crane Fly (larvae)) are small. Agricultural Extension Service Specialist for proper time to treat in your area. For Media small.

LAWNS AND OTHER RECREATIONAL AREAS: To Aid in Control: Fire Ants — Apply 1 fl. oz. per 100 sq. ft. Drench to mounds. **Notes:** Apply over and 6 inches around each mound. Apply gently for best results, apply when temperatures are 65-80°F or in early morning or late evening hours.

FRUITS, VEGETABLES, ROSES, FLOWERS, SHRUBS & SHADE TREES: To Control: Aphids, Bagworms, Bean Beetles, Cabbageworms, Codling Moth, Cucumber Beetles, Leafhoppers, Leafminers, Lacebugs, Leafrollers, Mealybugs, Mites, Scale Crawlers, Thrips. Thorough coverage is important. Try to hit undersides of leaves and penetrate dense foliage. Repeat if needed.

FRUITS: Apples (14), Apricots (10), Caneberries (7), Cherries (10), Grapes (10), Nectarines (10), Strawberries (15). **Note:** Start spraying after blooming and repeat at 10-14 day intervals.

VEGETABLES: Beans (7), Broccoli (5), Cabbage (1), Cantaloupes (3), Carrots (10), Cauliflower (1), Peppers (5), Potatoes (35), Radishes (10), Sweet Corn (1), Summer Squash (7), Tomatoes (1). **Note:** Numbers in () indicate days required between last application and harvest.

ROSES & FLOWERS: Azaleas, Camelias, Chrysanthemums, Carnations, Gladioli, Pyracantha, etc. **SHRUBS & SHADE TREES:** Birch, Boxwood, Hawthorn, Holly, Lilac, Locust, Maple, Pine, etc.

Note: Do not use on certain ferns, poinsettia, hibiscus, papaya, pines, gardenia, and orchids. To Control Mediterranean Fruit Fly (larvae): Apply 4 fl. oz. (1 1/2 cup) in 15 gals. water to above. Drench all soil within drip line. Make 3 applications at 14-16 day intervals as needed.

Combination Spray with Fungicide on Roses: ORTHO FUNGINEX® Rose Disease Control (sect(s) and disease(s) are found on roses. Follow directions and precautions on both labels.

NUISANCE PESTS: In outside areas.

To Control: Ants, Box Elder Bugs, Chiggers, Clover Mites, Cockroaches, Earwigs, Fleas, Flies, (Including Black Widow Spiders), Brown Dog Ticks — Apply 8 fl. oz. in 3 gals. water. Spray foundation to a height of 2-3 feet. Thoroughly spray windows and door screens, sills, foundations, walls, firewood piles, garbage cans, tree trunks, and into cracks and other places where not treat plants or animals with this nuisance pest rate. Do not burn sprayed firewood.

STORAGE AND DISPOSAL: Keep pesticide in original container. Do not put concentrate or dilute into food and foodstuffs. Store in a cool, dry place, preferably in a locked storage area. Do not store.

DISPOSAL: PRODUCT — Partially filled bottle may be disposed of by securely wrapping and discard in trash. **CONTAINER** — Do not reuse empty bottle. Rinse thoroughly before.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions. FUNGINEX® — Reg. TM of CELAMERCK GmbH Co. for Triflorine Fungicide.

Chevron Chemical Company © 1963

Ortho Consumer Products Division
San Francisco CA 94120-7144
R Form 8130 N2 Product 1000 49 Made in U.S.A.
EPA Reg. No. 239 7364-AA
EPA Est. 239 MO 1

2012