

JUN 22 1995

Mr. Frank Kamienski  
The Solaris Group  
Monsanto Company  
P.O. Box 5006  
San Ramon, CA 94583-0906

Dear Mr. Kamienski:

) Subject: Amendment - Add language for proprietary packaging  
ORTHO Diazinon Insect Spray  
EPA Reg. No. 239-2364  
Your submission dated March 22, 1995

The amendment referred to above, submitted in connection with registration under FIFRA section 3(c)(7)(a), is acceptable provided that you:

- 1) Submit and/or cite all data or other materials required for registration/reregistration of your product under FIFRA section 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.
- 2) Make the following revisions before your release the product for shipment under the amended labeling:
  - a) Revise the top front panel claim to "FOR OUTDOOR USE ONLY ON THE HOME LAWN AND HOME GARDEN".
  - b) As we discussed, the formula name will have to continue to appear on this and all diazinon products. Your request, however, will be presented for consideration to the diazinon reregistration team when the RED is being developed.
  - c) On page 3, the proposed revision to "Repeat as needed to maintain control." should be expanded to read "Repeat as needed and as directed to maintain control."
  - d) As we discussed, the limitation "USE CONTENTS WITHIN 6 MONTHS AFTER OPENING" must continue to appear on this product and all

29 11

other 25% diazinon EC products which are based on the current Ciba-Geigy formula and data. Future revised formulas may not require this limitation.

- e) In the proposed DISPOSAL directions, add "Tighten cap and" before "securely wrap...".

A stamped copy of the label is enclosed for your records.

- 3) Submit three (3) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Sincerely,

George T. LaRocca  
Product Manager (13)  
Insecticide Rodenticide Branch  
Registration Division (H7505C)

3 9 11

**BONUS 25% MORE 40 FL OZ AT REGULAR 32 FL OZ PRICE**

**(ICON) OUTDOOR USE ONLY**

ORTHO® Diazinon Insect Spray [ABN DIAZINON PLUS® Insect Spray]  
Makes up to 96 Gallons Diluted Spray[qt. size]

This product must not be used on golf courses and sod farms. *[This will appear in the upper 1/3 of the label]*

FOR OUTDOOR USE ONLY ON THE HOME LAWN AND IN THE HOME GARDEN  
DO NOT USE INSIDE THE HOME.  
FOR ROSES, FLOWERS, SHRUBS, FRUITS, VEGETABLES, HOME LAWNS & NUISANCE PESTS

**CONTROLS GARDEN INSECTS**

*Aphids, Cucumber Beetles, Cabbageworms, Codling Moth, Whiteflies*

**CONTROLS LAWN INSECTS**

*Ants, Fleas, Sod Webworms, Brown Dog Ticks, White Grubs*  
*Kills White Grubs*  
*Kills Black Widow Spiders*

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

Contains ORTHEX® Wetting Agent for More Effective Coverage

**NEW / NOW MEASURING CAP & POUR SPOUT**

**IMPROVED PACKAGE**

**NEW/NOW WITH MEASURING CAP**

**EASY POUR SPOUT**

**FAST & EASY MEASURING CAP**

JUN 22 1995

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 239-2364

**Active Ingredient**

Diazinon [~~0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate~~]  
.....25%

\*Inert Ingredients.....75%

\*Contains aromatic petroleum distillates

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See Back Panel Booklet for Additional Precautionary Statements

NET CONTENTS 32 fl oz (1 qt) 946 ml

NET CONTENTS 16 fl oz (1 pt) 473 mL

NET CONTENTS 64 fl oz (1/2 gal) 1.89 L

**NEW CONTENTS 40 FL OZ (1 QT & 8 FL OZ) 1.18 L**

3

4911

**Diazinon Plus Insect Spray controls a wide range of lawn and garden insect pests. Controls lawn and garden insects. For use on roses, flowers, trees, shrubs, lawns, nuisance pests, fruits, nuts, vegetables, and melons. May be applied with an ORTHO® DIAL 'n SPRAY®, hose-end, or tank type sprayer. Contains ORTHEX® which makes your spray spread more evenly on smooth, rough or waxy plant surfaces. Improved package with pour spout and measuring cap for fast, easy, accurate measuring. For the complete listing of all pests and plants, please read entire label.**

**[PICTURES OF BUGS] CONTROLS GARDEN INSECTS**

Aphids, Cucumber Beetles, Cabbageworms, Codling Moth, Whiteflies

**CONTROLS LAWN INSECTS**

Ants, Fleas, Sod Webworms, Brown Dog Ticks, White Grubs  
Kills White Grubs  
Kills Black Widow Spiders

**DO NOT USE INSIDE THE HOME. DO NOT USE ON HUMANS OR ANIMALS.**

*[THE FOLLOWING COPY IS FOR FIX-A-FORM BOOKLET TYPE LABELS WHICH CONSUMERS CAN OPEN, READ, AND RESEAL EASILY AT RETAIL AS WELL AS AT TIME OF USE]*

**Peel open for use directions and further information.**


**CAUTION:** Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye injury. See additional precautionary statements and use directions inside booklet. **PULL HERE TO OPEN LABEL. PRESS TO RESEAL.**

Medical Information call 1-800-454-2333  
Product Information call 1-800-225-2883

**Manufactured for**

The SOLARIS Group of  
Monsanto Company  
P.O. Box 5008  
San Ramon CA 94583-0808  
Form RS595-  
EPA Reg. No. 239-2364/ **Product 6019**  
EPA Est. 239-IA-3<sup>1</sup>, 58996-MO-1<sup>A</sup>  
Superscript is first letter of lot number

Made in USA  
UPC



4

5911

~~FOR GLUED LABELS WITH HANG TAGS AND PEEL/RESEAL LABELS~~

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL AND USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS.

**MEASURING CAP:** To open, squeeze cap sides while unscrewing. Fill cap to appropriate level for fast, easy, accurate measuring. Refer to label / table below for measuring instructions. Close cap tightly after use - no need to rinse.

- Level 1 = 1 teaspoon (tsp)
- Level 2 = 1 Tablespoon (Tbs) = 3 tsp = 1/2 fl oz
- Level 3 = 1 fl oz (2 Tbs)
- Level 4 = 2 fl oz (4 Tbs)

~~The following headings are for use if there is enough label space to have text in tables!~~

- ~~PESTS~~
- ~~PLANTS~~
- ~~DILUTION RATE~~
- ~~SPRAYER TYPE~~
- ~~PRODUCT TO MEASURE~~
- ~~AREA TO TREAT~~

tsp = teaspoon / Tbs = Tablespoon  
ROSES, FLOWERS, SHRUBS & SHADE TREES

**DILUTION RATE:** Fill cap to level 1 two times (2 tsp) per gallon water. To apply with an Ortho Dial 'n Spray Hose-End Sprayer: Pour product into sprayer jar. Do not add water. Set dial to 2 tsp and spray.

**HOW TO USE:** Use 2 tsp. (1/3 fl. oz.) per gal. of water. Spray when pests first appear. Spray plant to wet leaves and branches to dripping point. Repeat ~~as needed~~ applications only as directed to maintain control. **PESTS:** To Control: Aphids, Apple and Thorn Skeletonizers, Bagworms, Bean Beetles, Cabbageworms, Codling Moth, Cucumber Beetles, Cutworms, Dipterous Leafminers\* (such as Arborvitae, Azalea, Birch, Boxwood, Holly, and Oak), European Pine Shoot Moth, Flea Beetles, Hemlock Chermes, Hemlock Woolly Adelgid, Holly Bud Moth, Leafhoppers, Leafminers, Lacebugs, Leafrollers, Mealybugs, Mites (such as Carnation Bud, Carnation Shoot, Clover, Cyclamen, European Red, Privet, and Two-spotted), Oak Loopers, Obscure Root Weevil, Omnivorous Leaf-tiers, Pear Slugs, Scale Crawlers (such as Cottoncushion, Euonymus, Lecanium, Pine Needle, Pit, San Jose and Soft), Spidermites, Tent Caterpillars, Thrips, Webworms (such as Cotoneaster, Fall, Juniper and Mimosa), and Whiteflies. \*NOTE: Resistant

5

6911

leafminers will not be controlled by diazinon or similar type insecticides.

**PLANTS FOR:** Azaleas, Birch, Boxwood, Camellias, Chrysanthemums, Carnations, Gladioli, Hawthorne, Hemlock, Holly, Lilac, Locust, Maple, Pine, Pyracantha, Rhododendrons, Spruce, Sycamore, Willow. *Combination Spray with Fungicide on Roses: ORTHO FUNGINEX® Rose Disease Control may be mixed with this product when listed insect(s) and disease(s) are found on roses. Follow directions and precautions on both labels.*

Note: Do not use on Ferns, Poinsettia, Hibiscus, Papaya, Pilea, Gardenia, and Orchids since injury may occur.

To control Mediterranean Fruit Fly (larvae): Apply 3 fl. oz. per 12 gals. to cover 1000 sq. ft. of soil beneath host plants listed above. Drench all soil within dripline. Make 3 applications at 14 to 16 day intervals as needed.

**HOME LAWNS AND OTHER OUTDOOR HOME RECREATIONAL AREAS:** Measure the area to be sprayed. Calculate square feet by multiplying length by width. For example, 20' x 25' equals 500 sq ft. Spray the proper amount of product as directed. Thorough coverage is important for best results.

**DILUTION RATE/ PRODUCT TO MEASURE:** Fill cap to level 1 one and one half times (1 1/2 tsp) per gallon water. Spray evenly over 42 sq ft. To apply with an Ortho Dial 'n Spray Hose-End Sprayer: Fill cap to level 3 three times (3 oz) for every 500 sq ft of lawn area [ or fill cap to level 4 plus 1 cap to level 3 for every 500 sq ft of lawn area]. Spray evenly over desired area. (For other hose end sprayers)

**HOW TO USE:** Apply 3 fl oz (6 Tbs) per 12 gal of water to cover 500 sq. ft. (1-1/2 tsp. per gallon to cover 42 sq ft). Thoroughly wet grass a few hours before applying. **PESTS: Controls: To Control:** Ants, Armyworms, Bermudagrass Mites, Brown Dog Ticks, Chiggers, Chinch Bugs, Clover Mites, Crickets, Cutworms, Dichondra Flea Beetle, Digger Wasps, Earwigs, European Crane Fly (larvae), Fleas, Frit Flies, Grubs (Chafer, Japanese Beetle, May Beetle, Southern Chafer) Hyperodes Weevils, Millipedes, Lawn Billbugs, Lawn Moths, Leafhoppers, Rhodesgrass Scales, Sowbugs (Pillbugs), Sod Webworms, Springtails (Collembola). For Brown Dog Ticks, spray grass and shrubbery particularly near the house. To control: Digger Wasps, apply at dusk when wasps are not active. To control: Frit Flies, mow grass and water thoroughly before treatment, but delay application until grass is dry. Apply when crawlers first emerge. For Lawn Billbugs, Grubs, Southern Chafer (White Grubs), Hyperodes Weevils, and Mole Crickets water lawn thoroughly after treatment.

Notes: Consult your local Agricultural Extension Service for proper time to treat. For Mole Crickets, apply in May or June when pests are small.

**FIRE ANTS -- To Aid in Control: HOME LAWNS AND OTHER OUTDOOR HOME RECREATIONAL AREAS:** **DILUTION RATE:** Fill cap to level 3 (1 fl oz). Add to 1 gallon of water. [Or] Apply 1 fl oz (2 Tbs) in 1 gal of water as a drench to mounds. For best results, drench mounds apply when temperatures are 65 to 80°F or in morning or evening. Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Treat new mounds as they appear.

6

**NUISANCE PESTS OUTDOORS:** ~~In outside areas.~~ **DILUTION RATE:** ~~Fill cap to level 4 (2 fl oz) per gallon water.~~ ~~Apply 2 fl oz per gal of water.~~ Spray a 5 ft. band of soil around the house next to the foundation to a height of 2 to 3 ft. Do not apply to vinyl siding as staining may occur. Thoroughly spray walks, firewood piles, garbage cans, tree trunks, and into cracks and other places where insects can hide. Do not treat plants (other than tree trunks) at this rate. Do not burn sprayed firewood within 2 weeks following treatment.

**PESTS CONTROLS To Control:** Ants, Box Elder Bugs, Brown Dog Ticks, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Flies, Millipedes, Scorpions, Sowbugs (Pillbugs), Spiders (including Black Widow Spiders), Springtails (Collembola).

**VEGETABLES, FRUITS, MELONS AND NUTS**

**DILUTION RATE:** ~~Fill cap to level 1 two times (2 tsp) per gallon water.~~ ~~To apply with an Ortho Dial In Spray Hose-End Sprayer, Pour product into sprayer jar. Do not add water. Set dial to 2 tsp and spray.~~

**HOW TO USE:** Use 2 tsp (1/3 fl oz) per gal of water.

Spray plant thoroughly when pests first appear. Repeat only as directed to maintain control. Numbers in ( ) indicate days required between last application and harvest. Do not feed treated vines to ~~animals~~ livestock. Do not treat food crops grown in the greenhouse.

**VEGETABLES AND MELONS -- Foliar Spray:** Beans (lima, pole, and snap) +, fresh only (7), Broccoli (7) Cabbage (21), Cantaloupes (3), Carrots (14), Casabas (3), Cauliflower (7), Chinese Broccoli (10), Chinese Cabbage (Bok Choy and Napa) (10), Collards (10), Crenshaws (3), Cucumber (7), Endive (Escarole) (14), Honeydew (3), Kale (10), Lettuce (14) Muskmelons (3), Onions (14) (Bulb and Green), Parsnips (14), Peas +, fresh only (7), Peppers (5), Persian Melons (3), Potatoes (35), Radishes (14), Spinach (14), Summer Squash (7), Sweet Corn (7), Tomatoes (1), Turnips (14), Turnip Tops (14), Watermelons (3), and Winter Squash (3). Do not apply more than 3 gal. of water per 1000 sq. ft. of home vegetable garden. Do not apply more than 3 times per season for Beans and Peas, or more than 5 times per season for all other vegetable crops. Repeat sprays 7 to 10 days apart.

+NOTE: Do not use on dried bean or pea varieties. To protect bees, do not apply to bean or pea foliage if the crop or weeds in the treatment area are in bloom. Do not apply to corn during the pollen shed period. To Control: Aphids, Armyworms, Colorado Potato Beetle, Cucumber Beetles, Diamond Back Moth (larvae), Dipterous Leafminers \*\*, Flea Beetles, Grasshoppers, Imported Cabbage Worms, Leafhoppers, Mexican Bean Beetle, Mites, Spidermites, Onion Thrips, Thrips, Vinegar Fly, Whiteflies. \*\*NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.

**VEGETABLES - BEFORE PLANTING /**

~~Preplant: Do not use on dried bean or pea varieties.~~ To Control Cutworms: Cabbage, Carrots, Lettuce, Lima Beans, Peas, Pole Beans, Red (Table) Beets, Snap Beans, Sweet Corn, Tomatoes, Radishes, Turnips and Turnip Tops, Wireworms: Cabbage, Carrots, Lettuce, Peas, Snap Beans, Sweet Corn, Tomatoes. ~~Do not use on dried bean or pea varieties.~~

**DILUTION RATE:** ~~Fill cap to level 4 (2 fl oz). Add to 1 gallon of water. Spray over~~

350 sq. ft. To apply with an Ortho Dial 'n Spray Hose-End Sprayer: Fill cap to level 4 for every 350 sq. ft. of home garden area. Set dial to 2 fl. oz. and spray evenly over desired area. Before planting, apply 2 fl. oz. (4 Tbs.) in a minimum of 1 gal. of water to 350 sq. ft. Immediately mix into soil to the depth of 2 to 3 inches for Surface Cutworms, 3 to 6 inches for Subterranean Cutworms, and 4 to 8 inches for Wireworms. To Control Mole Crickets: Carrots, Lima Beans, Pole Beans, Red (Table) Beets, Snap Beans, Peas, Sweet Corn. 1 to 2 days before planting, apply 4 tsp. in a minimum of 1 gal. of water to 350 sq. ft. Fill cap to level 2 PLUS one cap measure to level 1 per gal. water (4 tsp.). Or fill cap to level 1 four times. Add to 1 gallon water. Spray evenly over 350 sq. ft. Immediately mix into the soil to a depth of 1 to 2 inches. To Control Root Maggots: Cabbage. Apply 2 fl. oz. (4 Tbs.) in a minimum of 1 gal. of water to 350 sq. ft. Fill cap to level 4 (2 fl. oz.). Add to 1 gallon water. Spray evenly over 350 sq. ft. Immediately mix into the soil to a depth of 2 to 3 inches. For Disease Control on Fruits and Vegetables apply ORTHO Multi-Purpose Fungicide DACTONIL 2787®. Follow label directions.

HOME FRUIT TREE ORCHARDS: Apricots, Cherries, Nectarines, Peaches, Plums, Prunes (21): Do not apply more than 4 gal. of water per 500 sq. ft. of home orchard. Do not apply more than 3 times per season. Repeat sprays for cherries, plums and prunes 7 to 10 days apart, for apricots, nectarines and peaches, 14 days apart. Note: To protect bees, do not apply to fruit trees when trees or substantial numbers of weeds around the trees are in bloom.

HOME NUT TREE ORCHARDS: ALMONDS (dormant season) CA only: Spray almonds only once to control San Jose and Parlatoria scale. Walnuts CA only (45): Do not apply to Walnuts more than 3 times per season or after husks crack open. Do not apply more than 6 gal. of water per 500 sq. ft. of home almond or walnut orchard. Repeat sprays for walnuts 14 days apart.

HOME VINEYARD OR BERRY PATCH: Grapes (28), Strawberries (5): Do not apply more than 3 gal. of water per 500 sq. ft. of grape vineyard or strawberry patch. Do not apply more than 5 times per season for grapes or 3 times per season for strawberries. Repeat sprays 7 to 10 days apart. Blackberries, Boysenberries, Dewberries, Loganberries, and Raspberries, CA, OR, WA only (7): Do not apply more than 5 gal. of water per 500 sq. ft. of caneberry patch. Do not apply more than 5 times per season. Repeat sprays 14 days apart. To Control: Aphids (such as Black Cherry, Leafcurl Plum, Mealy Plum, and Thistle), Cherry Fruit Flies, Dried Fruit Beetles (except California), Fruitworms (Maggots), Fruit Tree Leafrollers, Grape Berry Moth, Grape Leafrollers, Leafhoppers, Mealybugs, Mites (such as Brown, Two-Spotted Spider, and Cherry Rust), Raspberry Sawfly, Scale Crawlers (such as Olive, San Jose, Walnut, Italian Pear, and Frosted), Spider Mites, Strawberry Leafroller, Tentiform Leafminers, Thrips, Twig Borers, Walnut Caterpillars.

**STORAGE AND DISPOSAL**

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed and foodstuffs; Store in a cool, dry place, preferably in a locked storage area. Do not store near heat or open flame. Store in an area inaccessible to children and pets. USE CONTENTS WITHIN 6 MONTHS AFTER OPENING. Do not store diluted spray. Keep tightly



capped.

~~DISPOSAL: Securely wrap partially filled or empty container in several layers of newspaper and discard in trash. Do not reuse empty container. PRODUCT—Partially filled bottle may be disposed of by securely wrapping original container in several layers of newspaper tightly securing cap, and discard in trash. CONTAINER—Do not reuse empty bottle. Rinse thoroughly wrap in newspaper and discard in trash.~~

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS & DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye injury. Avoid breathing spray mist. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Wear a long sleeved shirt and long legged pants when spraying. Spray with the wind to your back. Do not spray on windy days. If clothing becomes wet from spray during use, remove clothing after spraying, wash affected body areas thoroughly with soap and water, and launder clothing before wearing again. **DO NOT USE INSIDE THE HOME.** Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. ~~(Child / Pet Icon)~~ Do not allow children or pets to go onto a treated area until the treated areas, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils such as spoons and measuring cups must not be used for food purposes after use in measuring pesticides. ~~(The preceding statement is optional if the package has the measuring cap — mandatory if it does not)~~ **FIRST AID STATEMENT OF PRACTICAL TREATMENT:** IF SWALLOWED: call a physician immediately. Do NOT induce vomiting unless no other treatment is available. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing. IF INHALED: Remove victim to fresh air and apply artificial respiration if indicated. Get medical attention. IF ON SKIN: wash promptly with soap and water. Rinse thoroughly. Get medical attention. If in eyes: Rinse eyes with plenty of water and call a physician immediately. Note to Physicians: Emergency Information call 1-800-454-2333. This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. This product contains aromatic petroleum distillates which may present an aspiration hazard. Gastric lavage may be indicated if product was taken internally.

**ENVIRONMENTAL HAZARDS:** This pesticide is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. For lawns, where irrigation (watering) is

10 9 11

recommended in the directions, or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product, however stop watering before puddling occurs. Keep out of water. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS: Do not use or store near heat or open flame.


NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Medical Information call 1-800-454-2333  
Product Information call 1-800-225-2883

ORTHO®, DIAL 'n SPRAY® Registered trademarks of Monsanto Company.  
DACONIL 2787® Registered trademark of ISK Biotech Corp.  
®Trademark of Monsanto Company  
©Monsanto Company 1994

**Manufactured for**

The SOLARIS Group of  
Monsanto Company  
P.O. Box 5008  
San Ramon CA 94583-0808  
Product 6019  
Form RS595-C  
EPA Reg. No. 239-2364  
EPA Est. 239-IA-3<sup>1</sup>, 58996-MO-1<sup>A</sup>  
Superscript is first letter of lot number  
Made in USA  
UPC



10

THE FOLLOWING EXAMPLES ARE TABLE HEADINGS AND LAYOUT WE HOPE TO BE ABLE TO USE. THIS WILL DEPEND ON THE AMOUNT OF LABEL SPACE AVAILABLE. Text in tables would be from registered copy.

**ROSES, FLOWERS, SHRUBS & SHADE TREES**

Spray thoroughly on leaves and branches to dilute point when pests first appear. Repeat spray only as needed to maintain control.

PESTS	DILUTION RATE / PRODUCT TO MEASURE	AREA TO TREAT
-------	------------------------------------	---------------

...text to come from registered label copy

**LAWNS**

Measure the area to be sprayed. Calculate square feet by multiplying length by width. For example, 20' x 25' equals 500 sq ft. Spray the proper amount of product as directed. Thorough coverage is important for best results.

PESTS	DILUTION RATE / PRODUCT TO MEASURE	AREA TO TREAT
...text to come from registered label copy		