



**ORTHO  
BUG-GETA®  
Snail & Slug Pellets**



**ORTHO**



**ORTHO**



**ORTHO**

**BUG-GETA®  
Snail & Slug  
Pellets**

**BUG-GETA®  
Snail & Slug  
Pellets**

**BUG-GETA®  
Snail & Slug  
Pellets**

ts and slugs. The highly  
ETA Snail & Slug Pellets

They are night feeders  
flower pots and around




LETT

**WARNING:** May be Fatal if Swallowed. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after using. Avoid contamination of food and foodstuffs. Keep out of reach of children and domestic animals. First Aid - If ingested, immediately induce vomiting and call a physician. Dispose of container promptly when empty.

This product may be toxic to wildlife. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

**IMPORTANT: KEEP PELLETS AWAY FROM DOGS, PETS AND CHILDREN. BAIT MAY BE ATTRACTIVE TO DOGS. USE AND STORE UNDER CONDITIONS WHERE**

**Kills Snails & Slugs**  
Remains effective after rain or sprinkling.



Active Ingredient	By Wt.
Metaldehyde	3.75%
Inert Ingredients	96.75%

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL WARNINGS AND DIRECTIONS. KEEP PESTICIDE IN ORIGINAL CONTAINER.**

**DIRECTIONS:** If ground is dry, water area to be treated thoroughly. ~~Use~~ the pellets around flowers, ornamental plants and lawn areas. Apply one pound of pellets to each 1000 sq. ft. Apply only to soil surface around plants. Do not apply to foliage or other plant parts. ~~Do not apply where food crops are grown.~~ Always wear protective gloves.

Bait areas likely to be infested at the beginning of the spring planting season before damage occurs. Treatment every two weeks should

A →

B →

# BUG-GETA<sup>®</sup> Snail & Slug Pellets

# BUG-GETA<sup>®</sup> Snail & Slug Pellets

# BUG-GETA<sup>®</sup> Snail & Slug Pellets

and slugs. The highly  
A Snail & Slug Pellets

They are night feeders  
over pots and around



LET

toned from wet ground,  
is attractive to children

but do not place in pots  
th plants. If soil is dry,

**WARNING:** May be Fatal if Swallowed. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after using. Avoid contamination of food and foodstuffs. Keep out of reach of children and domestic animals. First Aid - If ingested, immediately induce vomiting and call a physician. Dispose of container promptly when empty.

This product may be toxic to wildlife. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

**IMPORTANT:** KEEP PELLETS AWAY FROM DOGS, PETS AND CHILDREN. BAIT MAY BE ATTRACTIVE TO DOGS. USE AND STORE UNDER CONDITIONS WHERE THERE WILL NOT BE A HAZARD.

**NOTICE:** Buyer assumes all responsibility for safety and use not in accordance with directions.

**Chevron Chemical Company**

Ortho Division/San Francisco  
Calif. 94119

Richmond, Calif. Lafayette, Calif.  
Des Plaines, Ill. Perth Amboy, N.J.  
Atlanta, Ga.


Form 8431-A

EPA Reg. No. 239-2373-AA

Product 5096 Made in U.S.A.

## Kills Snails & Slugs

Remains effective after rain or sprinkling.



Active Ingredient	By Wt.
Metakdehyde	3.25%
Inert Ingredients	96.75%

**WARNING: KEEP OUT OF REACH OF CHILDREN.**  
**KEEP PELLETS AWAY FROM CHILDREN, DOGS & OTHER PETS.**  
**MAY BE FATAL IF SWALLOWED.**  
Read and follow all directions.  
Read warning statements — side panel.

**NET WEIGHT 1 POUND**

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL WARNINGS AND DIRECTIONS.**  
KEEP PESTICIDE IN ORIGINAL CONTAINER.

**DIRECTIONS:** If ground is dry, water area to be treated thoroughly. **A** → Toss the pellets around flowers, ornamental plants and lawn areas. Apply one pound of pellets to each 1000 sq. ft. Apply only to soil surface around plants. Do not apply to foliage or other plant parts. **B** → **Do not apply where food occurs or is grown.** Always wear protective gloves. Bait areas likely to be infested at the beginning of the spring planting season before damage occurs. Treatment every two weeks should give satisfactory control. **C** → Under conditions of frequent watering and high temperatures it may be necessary to treat more often.



# ORTHO BUG-GETA<sup>®</sup> Snail & Slug Pellets

