


239-2375


6-6-73

1044

ORTHO  
DIAZINON® Granules



TREATS UP TO 2500 SQUARE FEET



CONTROLS: Sod Webworms, Chinch Bugs, Cutworms,  
Japanese Beetle Grubs, Billbugs, Ants,  
Clover Mites, Sowbugs, Fleas, Crickets  
and Certain Other Lawn and Nuisance Pests

Active Ingredients	By Wt.
O,O-diethyl O-(2-isopropyl-4-methyl-6-pyrimidinyl)phosphorothioate . . . . .	2%
Inert Ingredients. . . . .	98%

U.S. Patent No. 2,754,243

CAUTION: Keep out of reach of children. See back panel for additional cautions.

NET WEIGHT 12 1/2 POUNDS


ORTHO DIAZINON® Granules Back Panel

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

DIRECTIONS

OVERALL LAWN PROTECTION



ORTHO DIAZINON® Granules control the following lawn insects on both grass and dichondra lawns: Chinch Bugs, Ticks, Ants, Sod Webworms (Lawn Moths), White Grubs (Japanese Beetle, European Chafer, Southern Chafer), Billbugs, Bermuda Mites, Crickets, Armyworms, Fleas, Leafhoppers, Sowbugs, Millipedes, Springtails.

WHEN TO APPLY

ORTHO DIAZINON® Granules should be applied when insects are present for control of the listed insects. For the control of Ants, Chinch Bugs, Sod Webworms (Lawn Moths), Armyworms, Leafhoppers, Cutworms and Bermuda Mites which have several generations per season, repeat application at 4-6 week intervals if insects are present.

204

To control White Grubs of Japanese Beetles, European Chafer and Southern Chafer, make applications anytime between late July and early October. For Billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

HOW TO APPLY

On most grass and dichondra lawns the contents of this bag should be spread evenly over an area of 2500 square feet (50' X 50'). For heavy lawns such as St. Augustine Grass, spread contents over 1600 square feet (40' X 40'). Use appropriate spreader settings listed in Table below. Note separate spreader settings listed for control of White Grubs and Billbugs.

For best results the lawn should be dry and mowed before application. Sprinkle lightly after application to wash granules off of foliage. For control of surface feeding insects, Armyworms and Sod Webworms (Lawn Moths), apply late in the evening and withhold heavy watering for 24 hours. To control Chinch Bugs, Cutworms, White Grubs, Billbugs and Ants, water the lawn thoroughly after application.

SPREADER SETTINGS  
(per 1000 sq. ft.)

Spreader Make and Model	Grass or Dichondra Lawn 5 lbs.	St. Augustine Lawn 7 1/2 lbs.	White Grubs and Billbugs 6 lbs.
ORTHO Lawn Spreader	4 3/4	5 3/4	5 1/4
Cyclone Lawn Spreader	4 3/4	5 3/4	5 1/4
Scott	6	7 1/2	6 3/4
Sears Roebuck Cross Country (19")	5 3/4	6 3/4	6 1/4
Montgomery Ward 89-108 (20")	7 3/4	8 1/2	8 1/4
Jackson 18-26 (18")	6	6 3/4	6 1/4
Lawn Beauty	6	6 3/4	6 1/4
Central Stamping	7 1/2	8	7 3/4

NOTE: Spreader settings are for covering lawns in one direction. Do not use criss-cross pattern. To insure complete coverage, overlap wheel tracks of all spreaders except the Cyclone Spreader and ORTHO Lawn Spreader. Variables inherent in equipment as well as accuracy of machine adjustment, may cause actual application rates to deviate slightly. Therefore, Chevron makes no warranty as to the uniformity of coverage obtained from the settings listed above.

SPOT TREATMENT

Ants: Apply one tablespoon of ORTHO Diazinon<sup>®</sup> Granules over and around each individual ant hill.

Digger Wasps: Apply one tablespoon of ORTHO Diazinon<sup>®</sup> Granules in and around each individual ground nest opening. Make the application about dusk when the wasps are not active.

BAND TREATMENT AROUND FOUNDATION

The following insects are commonly found around or near the foundation of houses as well as in the lawn. Certain of these insects may even enter the house.


- |                 |          |                          |
|-----------------|----------|--------------------------|
| Ants            | Crickets | Millipedes               |
| Brown Dog Ticks | Earwigs  | Sowbugs                  |
| Clover Mites    | Fleas    | Springtails (Collembola) |

For control of the above insects, apply a five-foot band of ORTHO Diazinon<sup>®</sup> Granules next to the foundation of the house. Distribute the granules uniformly using a rate of 8 ozs. for each 5' X 20' area to be treated.

CAUTION: Harmful if swallowed. Avoid breathing of dust. May be absorbed through skin. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Keep out of reach of children. Keep children and pets off treated areas until the following afternoon. Treat lawns in late afternoon. Do not reuse empty container.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes.

4074



NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.



\*DIAZINON® a registered trademark of CIBA-GEIGY Corporation

Chevron Chemical Company				
Ortho Division/San Francisco, Calif. 94119				
Richmond, Calif.	Lafayette, Calif.	Des Plaines, Ill.	Perth Amboy, N.J.	Atlanta, G
Form 8175-A	EPA Reg. No. 239-2375	Product 1000	Made in U.S.A.	