

# DIAZINON<sup>®</sup> Granules



# ORTHO

## EFFECTIVE ANT CONTROL

**KILLS HOME INVADING PESTS** — Ants, Brown Dog Ticks, Fleas, Clover Mites, Crickets, Earwigs, Millipedes, Sowbugs.

**KILLS LAWN INSECTS** — Chinch Bugs, Sod Webworms (Lawn Moths), White Grubs (Japanese Beetle, European Chafer, Southern Chafer), Ants, Billbugs, Bermuda Mites, Armyworms, Cutworms and Leafhoppers.

Active Ingredient	By Wt.
O,O-diethyl O-(2-isopropyl-4-methyl-6-pyrimidinyl) phosphorothioate	2%
Inert Ingredients	98%
U.S. Patent No. 2,754,243	

**CAUTION: KEEP OUT OF REACH OF CHILDREN.**

See back panel for additional cautions.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

## DIRECTIONS

**BAND TREATMENT AROUND FOUNDATION** for Home Invading Pests: The following insects and pests are commonly found around or near the foundation of houses as well as in the lawn. Certain of these insects may even enter the house.

Ants	Crickets	Millipedes
Brown Dog Ticks	Earwigs	Sowbugs
Clover Mites	Fleas	Springtails (Collembola)

For the control of the above insects, apply a five foot band of ORTHO DIAZINON Granules next to the foundation of the house. Distribute the granules uniformly using a rate of 8 oz. for each 100 sq. ft. area to be treated.

## NET WEIGHT 1 LB.

**Chevron Chemical Company**

Ortho Division/San Francisco CA 94119

Richmond CA 94804

Product 1000

Made in U.S.A.

Form 8175-B

EPA Reg. No. 239-2375

## SPOT TREATMENT

**ANTS:** Apply one level Tablespoonful of ORTHO DIAZINON Granules over and around each individual ant hill.

**DIGGER WASPS:** Apply one level Tablespoonful of ORTHO DIAZINON Granules in and around each individual ground nest opening. Make the application about dusk when the wasps are not active.

## OVERALL LAWN PROTECTION

ORTHO DIAZINON Granules control the following lawn insects on both grass and dichondra lawns: Chinch Bugs, Ticks, Ants, Sod Webworms (Lawn Moths), Cutworms, White Grubs (Japanese Beetle, European Chafer, Southern Chafer), Billbugs, Bermuda Mites, Crickets, Armyworms, Fleas, Leafhoppers, Sowbugs, Millipedes, Springtails.



## READ THE LABEL

**CAUTION:** Harmful if swallowed or absorbed through the skin. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling. Keep children and pets off treated area until completely dry.

**Note to Physicians:** Emergency Information - call (415) 233-3737. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Avoid contamination of feed and foodstuffs. Dispose of container promptly when empty by wrapping and placing in trash collection.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

ACCEPTED

APR 26 1976

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 239-2375

## When to Apply

ORTHO DIAZINON Granules should be applied when insects are present for control of listed insects. For control of Ants, Chinch Bugs, Sod Webworms (Lawn Moths), Armyworms, Leafhoppers, Cutworms and Bermuda Mites which have several generations per season, repeat applications at 4-6 week intervals if insects are present.

To control White Grubs of Japanese Beetles, European Chafer and Southern Chafer, make applications anytime between late July and early October. For Billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

## How to Apply

On most grass and dichondra lawns the contents of this canister (1 lb.) should be sprinkled evenly over area of 200 sq. ft. (10' x 20'). For heavy lawns such as St. Augustine Grass, spread contents over 128 sq. ft. (8' x 16'). For White Grub and Billbug control, spread contents evenly over 166 sq. ft. (10' x 16').

For best results the lawn should be dry and mowed before application. Sprinkle lightly after application to wash granules off foliage. For control of surface feeding insects, Armyworms and Sod Webworms (Lawn Moths), apply late in the evening and withhold heavy watering for 24 hours. To control Chinch Bugs, Cutworms, White Grubs, Billbugs and Ants, water the lawn thoroughly after application.

**NOTICE:** Buyer assumes all responsibility for safety and use not in accordance with directions.

DIAZINON<sup>®</sup> is a registered trademark of Ciba-Geigy Corporation.