



# ORTHO Diazinon Granules

- **KILLS HOME INVADING PESTS**—Ants, Brown Dog Ticks, Fleas, Clover Mites, Crickets, Earwigs, Millipedes, Sowbugs
- **KILLS LAWN INSECTS**— As Listed on Back Panel

This product must not be used on golf courses and sod farms.



Active Ingredient	By Wt.
Diazinon (0,0-diethyl-O-[2-isopropyl-6-methyl-4-pyrimidinyl] phosphorothioate)	2%
Inert Ingredients	98%

Keep out of reach of children

### CAUTION

See back panel for additional precautionary statements.

**NET WT. 1 LB.**



# ORTHO Diazinon Granules

**ACCEPTED**  
 MAR 24 1987  
 Under the Federal Insecticide,  
 Fungicide, and Rodenticide Act,  
 as amended, for the pesticide  
 registered under  
 EPA Reg. No. 239-2375

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS & DOMESTIC ANIMALS

#### CAUTION

Harmful if swallowed or absorbed through the skin. Causes eye irritation. Avoid contact with eyes, skin and clothing. Avoid inhalation of dust. Avoid contamination of food and feed. Wash thoroughly after handling. Keep children and pets off treated area until completely dry. Kitchen utensils such as measuring cups should not be used for food purposes after use in measuring pesticides.

#### STATEMENT OF PRACTICAL TREATMENT

If swallowed, DO NOT make person vomit unless directed to do so by a doctor. Immediately take person and container to a doctor or an emergency hospital. If this is not possible, telephone a physician, or emergency hospital, or Poison Control Center for medical advice. In case of eye contact, immediately flush eyes with fresh water for 15 minutes. If irritation persists, see a doctor. If on skin, wash thoroughly with soap and water. **Note to Physicians:** Emergency Information—call (415) 233-3737. Diazinon is an organophosphorus cholinesterase inhibitor. If signs and symptoms of cholinesterase inhibition develop, atropine sulfate is antidotal. 2-PAM may also be given in conjunction with atropine, but should not be used alone.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

**BAND TREATMENT AROUND FOUNDATION for Home Invading Pests:** The following insects and pests are commonly found around or near the foundation of houses as well as in the lawn. Certain of these insects may even enter the house.

Ants	Crickets	Millipedes
Brown Dog Ticks	Earwigs	Sowbugs
Clover Mites	Fleas	Springtails (Collembola)

For the control of the above insects, apply a five foot band of ORTHO Diazinon Granules next to the foundation of the house. Distribute the granules uniformly using a rate of 8 oz. for each 100 sq. ft. area to be treated. Water lightly after application.

#### SPOT TREATMENT

**ANTS:** Apply one level Tablespoonful of ORTHO Diazinon Granules over and around each individual ant hill. Water lightly after application.

**DIGGER WASPS:** Apply one level Tablespoonful of ORTHO Diazinon Granules in and around each individual ground nest opening. Make the application about dusk when the wasps are not active.

#### OVERALL LAWN PROTECTION

ORTHO Diazinon Granules control the following lawn insects on both grass and dichondra lawns: Chinch Bugs, Ticks, Ants, Sod Webworms (Lawn Moths), Cutworms, White Grubs (Japanese Beetle, European Chafer, Southern Chafer), Billbugs, Bermuda Mites, Crickets, Armyworms, Fleas, Leafhoppers, Sowbugs, Millipedes, Springtails.

#### When to Apply

ORTHO Diazinon Granules should be applied when insects are present for control of listed insects. For control of Ants, Chinch Bugs, Sod Webworms (Lawn Moths), Armyworms, Leafhoppers, Cutworms and Bermuda Mites which have several generations per season, repeat applications at 4-6 week intervals if insects are present.

To control White Grubs of Japanese Beetles, European Chafer and Southern Chafer, make applications anytime between late July and early October. For Billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

#### How to Apply

On most grass and dichondra lawns the contents of this canister (1 lb.) should be sprinkled evenly over area of 200 sq. ft. (10' x 20'). For heavy lawns such as St. Augustine Grass, spread contents over 128 sq. ft. (8' x 16'). For White Grub and Billbug control, spread contents evenly over 160 sq. ft. (10' x 16').

For best results the lawn should be dry and mowed before application. Sprinkle lightly after application to wash granules off foliage. For control of surface feeding insects, Armyworms and Sod Webworms (Lawn Moths), apply late in the evening and withhold heavy watering for 24 hours. To control Chinch Bugs, Cutworms, White Grubs, Billbugs and Ants, water the lawn thoroughly after application.

#### STORAGE AND DISPOSAL

**STORAGE:** Keep pesticide in original container. Do not put into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area.

**DISPOSAL: PRODUCT**—Partially filled canister may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash. **CONTAINER**—Do not reuse empty canister. Discard canister in trash.

**NOTICE:** Buyer assumes all responsibility for safety and use not in accordance with directions.

**Chevron Chemical Company** © 1984

Ortho Consumer Products Division  
 San Francisco CA 94120-7144  
 Product 5353 Made in U.S.A.  
 Form 8175-R  
 EPA Reg. No. 239-2375-AA  
 EPA Est. 239-MO-1

