

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust and contact with eyes, skin or clothing. Wash thoroughly after handling and before smoking or eating. May cause contact sensitization following repeated contact with skin of susceptible individuals. If sensitization reactions result, contact a physician. Do not store near food or feed products. Food utensils such as Tablespoons or measuring cups must not be used for food purposes after use in measuring pesticides. Do not contaminate ornamental fish ponds. Do not allow children or pets onto the treated area until granules have been watered in and the grass dried. **STATEMENT OF PRACTICAL TREATMENT:** If swallowed, call a physician immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing. If inhaled, remove victim to fresh air and apply artificial respiration if indicated. If on skin, wash promptly with soap and water. Rinse thoroughly. If in eyes, rinse eyes with plenty of water and call a physician immediately. **Note to Physicians:** Emergency Information—call 1-800-457-2022. This is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

ENVIRONMENTAL HAZARDS: This pesticide is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping granules. Where irrigation (watering) is recommended in the directions, or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), apply a minimum of 1/4 inch of water immediately after applying this product. Stop irrigation before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to waterfowl wetlands (swamps, bogs, marshes, and potholes). Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

BAND TREATMENT AROUND FOUNDATION for Home Inhabiting Pests: The following insects and pests are commonly found around or near the foundation of houses as well as in the lawn. Certain of these insects may even enter the house.

Ants	Clover Mites	Earwigs	Milkpedes	Springtails
Brown Dog Ticks	Crickets	Fleas	Sowbugs	(Collembola)

For the control of the above insects, apply a five foot band of ORTHO Diazinon Granules next to the foundation of the house. Distribute the granules uniformly using a rate of 8 oz. for each 100 sq. ft. area to be treated. Water lightly after application.

SPOT TREATMENT

Ants: Apply one level Tablespoonful of ORTHO Diazinon Granules over and around each individual ant hill. Water lightly after application.

Digger Wasps: Apply one level Tablespoonful of ORTHO Diazinon Granules in and around each individual ground nest opening. Make the application about dusk when the Wasps are not active.

OVERALL LAWN PROTECTION

ORTHO Diazinon Granules control the following lawn insects on both grass and dichondra lawns: Chinch Bugs, Brown Dog Ticks, Ants, and Webworms (Lawn Moths), Cutworms, White Grubs (Japanese Beetle, European Chafer, Southern Chafer), Billbugs, Bermudagrass Mites, Crickets, Armyworms, Fleas, Leafhoppers, Sowbugs, Milkpedes, Springtails.

When to Apply: ORTHO Diazinon Granules should be applied when insects are present and lawn is dry. For control of Ants, Chinch Bugs, Sod Webworms (Lawn Moths), Armyworms, Leafhoppers, Cutworms and Bermudagrass Mites which have several generations per season, repeat applications at 7 to 10 days, if necessary. Water lightly after application. To control White Grubs of Japanese Beetles, European Chafer and Southern Chafer, make applications anytime between late July and early October. For Billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

How to Apply: On most grass and dichondra lawns the contents of this canister (1 lb. 4 oz.) should be sprinkled evenly over area of 250 sq. ft. (10' x 25').

For best results the lawn should be dry and mowed before application. Sprinkle lightly after application to wash granules off foliage. For control of surface feeding insects, Armyworms and Sod Webworms (Lawn Moths), apply late in the evening and withhold heavy watering for 24 hours. To control Chinch Bugs, Cutworms, White Grubs, Billbugs and Ants, water the lawn thoroughly after application.

STORAGE AND DISPOSAL

STORAGE: Keep pesticide in original container. Do not put into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area.

DISPOSAL: PRODUCT—Partially used canister may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash. **CONTAINER**—Do not reuse empty canister. Discard canister in trash.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Chevron Chemical Company © 1992

Ortho Consumer Products Division

P.O. Box 5047 San Ramon CA 94583-0947

Form C407-B Product 5353

EPA Reg. No. 239-2375-AA EPA Est. 239-MO-1



PRECAUTIONARY STATEMENTS

DOMESTIC ANIMALS

swallowed, inhaled, or absorbed through skin. Avoid breathing dust and contact with thoroughly after handling and before smoking or eating. May cause contact sensitization with skin of susceptible individuals. If sensitization reactions result, contact a food or feed products. Food utensils such as Tablespoons or measuring cups must not be used in measuring pesticides. Do not contaminate ornamental fish ponds. Do not be treated area until granules have been watered in and the grass dried. **STATEMENT**: If swallowed, call a physician immediately. Drink one or two glasses of water and the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting person is unconscious or convulsing. If inhaled, remove victim to fresh air and apply. If on skin, wash promptly with soap and water. Rinse thoroughly. If in eyes, rinse and call a physician immediately. **Note to Physicians:** Emergency Information—call anticholinesterase inhibitor. If symptoms of cholinesterase inhibition are present, at antitoxin. 2-PAM is also antitoxin and may be administered, but only in conjunction with antitoxin.

This pesticide is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, are most susceptible. Because of the migratory habits of certain Atlantic Coast product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed label rates. Application rates above those recommended significantly increase waterfowl. Avoid overlapping granules. Where irrigation (watering) is recommended, i.e., ducks or geese, can be expected in the treated area after treatment (except for 1/4 inch of water immediately after applying this product. Stop irrigation before lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water, ditches, and potholes. Runoff may be hazardous to aquatic organisms in neighboring areas. Be killed at application rates recommended on this label. Do not apply where fish, wildlife are important resources. Do not contaminate water by cleaning of equipment near water.

to use this product in a manner inconsistent with its labeling. **STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND**

FOUNDATION for Home Inhabiting Pests: The following insects and pests are commonly found in houses as well as in the lawn. Certain of these insects may even enter the

Chinch Bugs	Earwigs	Millipedes	Springtails
Crickets	Fleas	Sowbugs	(Collembola)

insects, apply a five foot band of ORTHO Diazinon Granules next to the foundation of houses uniformly using a rate of 8 oz. for each 100 sq. ft. area to be treated. Water lightly

careful of ORTHO Diazinon Granules over and around each individual ant hill. Water 1/2 Tablespoonful of ORTHO Diazinon Granules in and around each individual ground nest about dusk when the Wasps are not active.

Control the following lawn insects on both grass and dichondra lawns: Chinch Bugs, Fleaworms (Lawn Moths), Cutworms, White Grubs (Japanese Beetle, European Chafer, Armyworms, Fleas, Leafhoppers, Sowbugs, Millipedes,

non Granules should be applied when insects are present and lawn is dry. For control Fleaworms (Lawn Moths), Armyworms, Leafhoppers, Cutworms and Bermudagrass weevils per season, repeat applications at 7 to 10 days, if necessary. Water lightly White Grubs of Japanese Beetles, European Chafer and Southern Chafer, make application July and early October. For Billbug control, apply when activity is first observed or indicates damage from these insects. Repeat as necessary.

and dichondra lawns the contents of this canister (1 lb. 4 oz.) should be sprinkled (18" x 25').

Soil should be dry and moved before application. Sprinkle lightly after application to wash soil of surface feeding insects, Armyworms and Sod Webworms (Lawn Moths), apply soil heavy watering for 24 hours. To control Chinch Bugs, Cutworms, White Grubs, lawn thoroughly after application.

DISPOSAL

original container. Do not put into food or drink containers. Avoid contamination of a cool, dry place, preferably in a locked storage area.

any filled canister may be disposed of by securely wrapping original container in plastic and discard in trash. **CONTAINER**—Do not reuse empty canister. Discard canister in trash.

responsibility for safety in directions.

© 1982
Peston
94583-0847

PA Est. 238-MO-1



ACCEPTED

MAR 11 1993

Under the Federal Insecticide, Fungicide, and Rodenticide Act, this pesticide is registered under EPA Reg. No.

- KILLS HOME INVA
Clover Mites, Cric
- KILLS LAWN INSE

NET

BEST