

ORTHO®  
Diazinon Granules

This product must not be used on golf courses and sod farms. *[This must appear in the upper 1/3 of the product label]*

KILLS HOME INVADING PESTS-Ants, Brown Dog Ticks, Fleas, Crickets, Earwigs, Millipedes, Sowbugs

KILL LAWN INSECTS-As Listed on Back Panel

EFFECTIVE OUTDOOR ANT CONTROL

*[FOR THE 1 LB. 4 OZ. SIZE ONLY  
25% MORE FREE  
20 OZ. FOR THE PRICE OF 16 OZ.]*

For outdoor use on home lawns and around the perimeter of the home only.

Active Ingredient

Diazinon [0,0-diethyl-0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate].....2%

Inert Ingredients .....98%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional precautionary statements.

1 lb/ 453g

For outdoor use on home lawns and around the perimeter of the home only. Kills home invading pests: Ants, Brown Dog Ticks, Clover Mites, Crickets, Earwigs, Fleas, Millipedes, Sowbugs, Springtails. Kills lawn insects as listed below.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.


Tbs = Tablespoon

HOME INVADING PESTS:

Ants	Clover Mites	Earwigs	Millipedes
Springtails	Brown Dog Ticks	Crickets	Fleas
(Collembola)	Sowbugs		

~~Apply a five ft band of ORTHO Diazinon Granules next to the foundation of the house. Distribute the granules uniformly using 8 oz for each 100 sq ft area. Water~~

12/22/93 11/11/1994



lightly after application.

Sprinkle a 2 foot band of granules evenly next to the house foundation. Use 8 oz. for each 100 sq. ft. area (2 ft by 50 ft). Repeat application at 3 week intervals if needed.

**SPOT TREATMENT**

Ants: Apply 1 level Tbs over and around each ant hill. Water lightly after application.

Digger Wasps: Apply 1 level Tbs in and around each ground nest opening. Apply about dusk when the Wasps are not active.

OVERALL LAWN PROTECTION - grass and dichondra lawns: Controls: Chinch Bugs, Brown Dog Ticks, Ants, Sod Webworms (Lawn Moths), Cutworms, White Grubs (Japanese Beetle, European Chafer, Southern Chafer), Billbugs, Bermudagrass Mites, Crickets, Armyworms, Fleas, Leafhoppers, Sowbugs, Millipedes, Springtails.

**When and How to Apply:**

Apply when insects are present. To control White Grubs of Japanese Beetles, European Chafer and Southern Chafer, make applications anytime between late July and early October. For Billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary. For best results the lawn should be dry and mowed before application. Sprinkle the contents of this canister (1 lb) evenly over area of 200 sq. ft. (10'x20'). Water lightly after application to wash granules off foliage. For control of surface feeding insects, Armyworms and Sod Webworms (Lawn Moths), apply late in the evening and withhold heavy watering for 24 hours. To control Chinch Bugs, Cutworms, White Grubs, Billbugs and Ants, water the lawn thoroughly after application.

**LAWN TREATMENT**

The lawn should be dry and mowed before application.

PEST	Amount to Use	Timing and Special Directions
------	---------------	-------------------------------

<p>Ants, Chinch Bugs, Brown Dog Ticks, Lawn Moths (Sod Webworms), Bermudagrass Mites, Crickets, Armyworms, Fleas, Leafhoppers, Sowbugs (Pillbugs), Millipedes, Springtails (Collembola)</p>	<p>Sprinkle 1 lb. per 200 sq ft (10 ft by 20 ft) evenly over lawn area</p>	<p>Apply when insects are present.</p> <p>Water lightly after treatment to wash granules off foliage.</p> <p>Repeat Application in 3 weeks if necessary. May be applied up to 4 times per year.</p>
<p>White Grubs (Japanese Beetle, European &amp; Southern Chafer)</p>	<p>☞</p>	<p>Treat anytime between late July and early October</p> <p>Water grass thoroughly with at least 1/4 inch of water after application; however, stop watering before puddling occurs. Consult the State Agricultural Extension Service for timing in your area.</p>
<p>Hyperodes Weevils</p>	<p>☞</p>	<p>Treat problem areas in mid-April and again in mid-May.</p> <p>Water grass thoroughly with at least 1/4 inch of water after application; however, stop watering before puddling occurs.</p>
<p>Billbugs</p>	<p>☞</p>	<p>Apply when chewed or brown grass indicates damage. Water grass thoroughly with at least 1/4 inch of water after application; however, stop watering before puddling occurs. If necessary, repeat application in 3 weeks.</p> <p>May be applied up to 4 times per year</p>

## STORAGE AND DISPOSAL

**STORAGE:** Keep pesticide in original container. Do not put into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferable in a locked storage area.

**DISPOSAL: PRODUCT-** Partially filled canister may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash.

**CONTAINER-** Do not reuse empty canister. Discard canister in trash.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS & DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust and contact with eyes, skin or clothing. Wash thoroughly after handling and before smoking or eating. May cause contact sensitization following repeated contact with skin of susceptible individuals. If sensitization reactions result, contact a physician. Do not store near food or feed products. Food utensils such as tablespoons or measuring cups must not be used for food purposes after use in measuring pesticides. Do not contaminate ornamental fish ponds. Do not allow children or pets onto the treated areas until granules have been watered in and the grass dried.

**FIRST AID:** If swallowed, call a physician immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing. If inhaled, remove victim to fresh air and apply artificial respiration if indicated. If on skin, wash promptly with soap and water. Rinse thoroughly. If in eyes, rinse eyes with plenty of water and call a physician immediately. Note to Physicians: Emergency Information-call 1-800-454-2333. This is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only on conjunction with atropine.

**ENVIRONMENTAL HAZARDS:** This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl (ducks or geese) feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds. Avoid overlapping granules. Follow all watering instructions in the directions for use, however, stop watering before puddling occurs. Do not apply this product to sites occupied by waterfowl, or within 75 ft. of any body of water that may attract waterfowl, such as ponds or lakes. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water. Runoff may be hazardous to fish and other aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

~~This pesticide is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping granules. Where irrigation (watering) is recommended in the directions, or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), apply a minimum of 1/4 inch of water immediately after applying this product. Stop irrigation before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.~~

) NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

▣ Medical Information call 1-800-454-2333  
Product Information call 1-800-225-2883

©Trademark of Monsanto Company

©Monsanto Company 1994

) The SOLARIS Group of  
Monsanto Company  
P.O. Box 5008  
San Ramon CA 94583-0808  
Form \_\_\_\_\_ Product 5353  
EPA Reg. No. 239-2375 EPA EST. 58996-MO-1  
Made in USA