

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER

Causes permanent eye damage. May cause allergic skin reactions. Harmful if swallowed or inhaled. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Avoid breathing dust or vapor. Wash hands after use. Remove and launder clothing before reuse.

STATEMENT OF PRACTICAL TREATMENT

In case of eye contact, immediately flush eyes with fresh water for 15 minutes and see a doctor. If swallowed, give a large amount of water to drink, induce vomiting, and call a doctor. For skin contact, wash with plenty of soap and water. See a doctor if irritation occurs or rash develops.

Note to Physicians: Emergency Information - call (415) 233-3737

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ADHERE TO ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal or cleaning equipment.

STORAGE

Keep in original container. Store in cool, dry place. Protect from excessive heat. Do not store near food or feed.

ESTICIDE DISPOSAL

Do not pour, flush, or dump into water that cannot be used, or drain into sewer. Dispose of according to applicable federal, state and local regulations.

CONTAINER DISPOSAL

Triple empty bag by shaking and tapping sides and bottom to dislodge particles. Empty residue into application equipment. Then place bag in a sanitary landfill or by incineration if allowed by state and local authorities.



ORTHO

ORTHOOCIDE®

Soil Treater "X"

(FUNGICIDE)

Active Ingredients	By Wt.
*Captan	9.5%
Related Derivatives	0.5%
Pentachloronitrobenzene	10.0%
Inert Ingredients	80.0%
*N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide	

KEEP OUT OF REACH OF CHILDREN
DANGER

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS.

ACCEPTED

APR 15 1982

NET WEIGHT

Chevron Chemical Company

Ortho Agricultural Chemicals Division
San Francisco CA 94119 Richmond CA 94804

Product 2324 Made in U.S.A.

Form 7029 I

EPA Est. 239-CA-5

EPA Reg. No. 239-2382-AA

ORTHOOCIDE® Soil Treater "X" is a fungicide for use on fungicide seed and soil. It is a systemic fungicide that kills fungi in the soil and on the seed. It is used to control seed and soil-borne fungi. It is used on a wide range of crops. It is used on a wide range of crops. It is used on a wide range of crops.

COTTON SEED: Planter Box Application—Damp Off, Rhizoctonia, Seed and Seedling Rot, Acid, Reginald and Fuzzy Seed. Use 2 1/2 to 3 lbs. per acre. Apply to seed through planter box at time of planting. This is a pre-planting treatment which must be followed by additional fungicide treatments. Planting rate must be adjusted after adding the required dosage of ORTHOOCIDE® Soil Treater "X" to planter box. Do not use treated seed for food, feed or oil.

COTTON: Furrow Planting Treatment—Damp Off, Rhizoctonia, Seed and Seedling Rot. Use 12 lbs. per acre based on 40-inch row spacing. Apply to seed through planter box at time of planting. This is a pre-planting treatment which must be followed by additional fungicide treatments. Planting rate must be adjusted after adding the required dosage of ORTHOOCIDE® Soil Treater "X" to planter box. Do not use treated seed for food, feed or oil.

BEANS: Root and Stem Rot. 15 to 20 lbs. per acre, based on 36-inch row spacing. Apply in the furrow and over the seed and turn into soil and broadcast. Calibrate dusting equipment to apply 2 1/2 to 3 lbs. per 145 linear feet of row. Do not feed treated beans to livestock.

PEPPERS: Damping Off in Seed Bed. 25 lbs. per acre (10 lbs. per 1,000 sq. ft.). Apply as a pre-planting soil treatment by dusting over the seed bed. Do not use treated seed for food, feed or oil. For Field Seeded Peppers. Dust in seed furrow at planting time using 20 to 30 lbs. per acre (equivalent to 10 to 12 lbs. per 1,000 sq. ft.) of single row with 40" row spacing.

LAWN: Brown Patch, Dollar Spot. One day before treatment, mow lawn and remove clippings. Do not water lawn after treatment. Apply ORTHOOCIDE® Soil Treater "X" at 1 lb. per 1,000 sq. ft. of lawn. Do not use treated lawn for food, feed or oil. Following treatment, water lawn and mow lawn every 3 to 4 days. Do not mow lawn with rotary mower. Water lawn regularly every 3 to 4 days. On New Lawns, first treatment should be applied when first seeds germinate and repeat every 10 days until lawn is well established.

BULBIOUS IRIS: Crown Rot, Black Rot. 5 lbs. per 100 sq. ft. Spread on the soil surface and water in to a depth of 4 to 6 inches. Best results are obtained if applied in the fall of the year.

LARKSPUR, CALENDULA, SWEET PEA, SNAPDRAGON: Rhizoctonia, Stem Rot, Damping Off, Root Rot. 5 lbs. per 100 sq. ft. Spread on the soil surface and water in to a depth of 4 to 6 inches before planting.

CAMELLIA: Flower Blight. 5 lbs. per 100 sq. ft. Apply and water during early winter. Do not use treated plants for food, feed or oil. During the winter, water plants regularly and mow lawn every 3 to 4 days. Do not mow lawn with rotary mower. Water lawn regularly every 3 to 4 days.

CARNATIONS, AFRICAN VIOLETS, FLOINSETTIA, CHRYSANTHEMUMS, BEGONIA: Stem Rot, Damping Off, Root Rot. 5 lbs. per 100 sq. ft. Apply as a pre-planting soil treatment by dusting over the seed bed. Do not use treated seed for food, feed or oil.

GREENHOUSE BENCH TREATMENT: On all crops for listed diseases. Mix into top 3 to 4 inches of soil before planting at rates given above.

Do not feed treated seed to animals. Do not use treated seed for food, feed or oil. Do not use treated seed for food, feed or oil. Do not use treated seed for food, feed or oil. Do not use treated seed for food, feed or oil.

Do not apply under conditions involving possible drift to or contamination of food, forage, feed or animals.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the label. It does not warrant that the material will be effective against all diseases or pests. 2. This material and all other factors beyond Chevron's control present a risk of injury to crops and property. Such risks include, but are not limited to, the use of this material on crops and property. Such risks include, but are not limited to, the use of this material on crops and property. Such risks include, but are not limited to, the use of this material on crops and property.

Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1. above) resulting from handling, storage, and use of this material.