


ORTHO DIAZINON Insect Spray

ORTHO

DIAZINON

Insect Spray



Active Ingredients
10.0 diethyl 0-12 isopropyl 6-methyl-4
pyrimidinyl phosphorothioate 25%
Aromatic Petroleum Derivative Solvent 57%
Inert Ingredients 18%
U.S. Pat. No. 2,754,145

WARNING: Keep out of reach of children.
Read warning statement on back panel.

NET CONTENTS 1 QUART LIQUID

THIODAN®-Methyl Parathion-Toxaphene 2.5-2.5-10 Dust



ORTHO

(INSECTICIDE)

Active Ingredients	By Wt.
*Endosulfan	2.5%
O,O-dimethyl O-p-nitrophenyl thiophosphate	2.5%
**Toxaphene	10.0%
Inert Ingredients	85.0%
*Hexachlorohexahydromethano-2,4,3-benzodioxathiepin oxide	
**Technical Chlorinated Camphene	
(Chlorine Content: 67% to 69%)	
U. S. Pat. No. 2,799,685	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH DANGER STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT DILUTE INTO FOOD OR DRINK CONTAINERS. BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

DIRECTIONS

This product is formulated for application by ground or air equipment.

BROCCOLI, CABBAGE, CAULIFLOWER, LETTUCE: Aphids, Armyworms, Cabbage Loopers, Cabbage Worms, Imported Cabbage Worms, Climbing Cutworms, Surface-feeding Cutworms, Flea Beetle—25 to 40 lbs. per acre. Do not apply

NET WEIGHT 50 LBS.


Chevron Chemical Company

Ortho Division/San Francisco, Calif. 94119

Richmond, California Fresno, California Honolulu, Hawaii
Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 8262 EPA Reg. No. 239.2384 Product 5159 Made in U.S.A.

DANGER:
KEEP OUT OF REACH OF CHILDREN
DO NOT TAKE INTERNALLY
DO NOT INHALE
DO NOT GET ON SKIN



POISON



ANTIDOTE: If Swallowed. Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. If On Skin. In case of contact, remove contaminated clothing and immediately wash skin with soap and water. **CALL A PHYSICIAN IMMEDIATELY!** Note to Physicians: This product is a combination of an organic phosphate and chlorinated hydrocarbons. Atropine is antidotal, but prolonged (at least 12 hours) observation should be carried out for signs of central nervous system excitation which should be treated symptomatically if they appear.

DANGER: Poisonous if swallowed, inhaled or absorbed through skin. Wear a mask or respirator of a type passed by the U. S. Bureau of Mines for methyl parathion protection. Rapidly absorbed through skin! Do not get in eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse. Keep all unprotected persons out of operating area in vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticides and volatile residues have dissipated. Do not store or transport near feed or food. Do not use on young plants. Do not use on sorghum, milo. **Do not apply under conditions involving possible drift to food, forage, or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not contaminate any body of water, woodlands, or other non-crop areas. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

ACCEPTED

MAR 10 1972

after edible parts start to form or within 30 days of harvest. Do not apply to Cabbage after heads start to form. Do not apply to Leaf Lettuce after seedling stage or to head lettuce after heads begin to form. Remove plants from field at harvest. Do not feed crop waste to livestock.

TOMATOES, PEPPERS: Aphids, Armyworms, Hornworm, Pepper Maggot—25 to 30 lbs. per acre. Do not apply within 15 days of harvest.

SPINACH: Aphids, Crown Mite, Armyworm—20 lbs. per acre. Do not apply more than once. Do not apply within 15 days of harvest.

STRAWBERRIES: Aphids, Spittlebug—30 lbs. per acre. Do not apply after fruit starts to form.

CELERY: Aphids, Flea Beetles, Leafhoppers, Spider Mites—20 lbs. per acre. Do not apply after plants start to bunch or are half mature.

CONDITIONS OF SALE: 1. Chevron Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the use stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risk and liability, except that claimed by Chevron under the above, resulting from handling, storage, and use of the material.

THIODAN is a registered trademark of Chevron Chemical Company.