

Dormant Disease Control



ORTHO

(LIME SULFUR FUNGICIDE)

PREVENTS PEACH LEAF CURL
Also Aids in Control of Black Spot
and Powdery Mildew on Roses

Active Ingredient	By Wt.
Calcium Polysulfide	26%
Inert Ingredients	74%
alpha-[p-(1,1,3,3-Tetramethylbutyl)phenyl]-omega-hydroxypoly(oxyethylene)	10%
Density Baume at 60 F	29

DANGER: KEEP OUT OF REACH OF CHILDREN.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL DANGER STATEMENTS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

ORTHO Dormant Disease Control helps to prevent foliage damage from occurring during the growing season by killing disease-causing spores that over-winter on buds, twigs and bark and by providing a protective covering of fruit surfaces before they open in the spring. Since many of the disease organisms covered on this label also over-winter on old leaves they should be raked up in the fall and disposed of.

NET CONTENTS

LIQUID

Chevron Chemical Company
Ortho Division/San Francisco CA 94119
Product 5154 Made in U.S.A.
Form 8184 A

Richmond CA 94804

EPA Reg. No. 239-2391-AA



READ THE LABEL

DANGER: Harmful if swallowed or absorbed through the skin. Concentrate causes eye and skin damage. Do not get in eyes, on skin or on clothing. Avoid breathing spray mist. Wear water-proof gloves and face shield or goggles when handling concentrate. In case of contact, immediately flush eyes or skin with plenty of water. For eye or skin contact, call physician. Wash hands after use.

Note to Physicians: Emergency Information - call 415-333-3737

Avoid spraying on painted surface as staining may occur. Keep bottle tightly closed when not in use. If a crust should form on the surface, break through the crust and shake thoroughly. Product quality is not impaired by crystallization. When container is empty, immediately wash thoroughly and destroy. Never reuse.

DIRECTIONS


Apply ORTHO Dormant Disease Control to thoroughly cover all exposed plant parts. Use at the rate of 10 tablespoons (5 fl. oz.) per gallon of water in an ORTHO SPRAYETTE, tank type or power sprayer. Do not spray when the temperature is over 80° or under 45°. Do not store diluted spray. Make new dilution for each use.

PEACHES: Peach Leaf Curl, Peach Blight—Timing is very important. **Late Fall**—Spray in October or November after leaf drop but before heavy rains occur. **Mid-Winter**—Apply a full coverage spray in January or February before buds begin to swell. Note: If buds have begun to swell or open, it is too late to obtain satisfactory control of Peach Leaf Curl Disease as infection has already occurred. For best results we recommend a combination of Late Fall and Mid-Winter applications.

ROSES: Black Spot, Powdery Mildew, Rust—Prune and destroy all old leaves. Spray in the spring before buds begin to swell.

PEARS: Blister Mite—Apply in the fall before the leaves start to turn color and again in the spring just as the buds begin to swell.

BLACKBERRIES, RASPBERRIES: Leaf Spot, Cane Blight, Powdery Mildew—Prune out old canes and destroy before fall rains. Apply in late fall and again in the spring as the buds begin to swell.



NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.