



ORTHO Dormant Disease Control

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER: Corrosive to eyes. Concentrate causes eye damage and skin irritation. Harmful if swallowed or absorbed through the skin. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Avoid breathing spray mist. **WASH OF PRACTICAL TREATMENT:** In case of eye contact, immediately flush eyes with fresh water for 15 minutes and see a doctor. Wash skin thoroughly with soap and water. If swallowed, give large quantities of water, induce vomiting and see a doctor immediately. **Note to Physicians:** Emergency Information—call (815) 233-3737.

ENVIRONMENTAL HAZARDS: Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. **PHYSICAL OR CHEMICAL HAZARDS:** MAY GIVE OFF HIGHLY TOXIC AND EXTREMELY FLAMMABLE HYDROGEN SULFIDE GAS IF MIXED WITH ACIDS.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS

ORTHO Dormant Disease Control helps to prevent foliage diseases from occurring during the growing season by killing disease causing spores that over winter on buds, twigs and bark and by providing a protective covering of the buds before they open in the spring. Since many of the disease organisms covered on this label also over winter on old leaves they should be raked up in the fall and disposed of.

DIRECTIONS
Apply ORTHO Dormant Disease Control to thoroughly cover all exposed plant parts. Use at the rate of 10 Tablespoons (5 fl. oz.) per gallon of water in an ORTHO SPRAYETTE tank type or power sprayer. Do not spray when the temperature is over 80°F or under 45°F. Make new dilution for each use.

PEACHES: Peach Leaf Curl, Peach Blight, Peach Shot, Peach Yellows, Peach Rust, Peach Scale, Peach Mite, Peach Nematode, Peach Root Rot, Peach Tree Canker, Peach Tree Borer, Peach Tree Weevil, Peach Tree Aphid, Peach Tree Spider Mite, Peach Tree Scale, Peach Tree Nematode, Peach Tree Root Rot, Peach Tree Canker, Peach Tree Borer, Peach Tree Weevil, Peach Tree Aphid, Peach Tree Spider Mite, Peach Tree Scale, Peach Tree Nematode.

Curt Disease as infection has already occurred. For best results we recommend a combination of Late Fall and Mid Winter applications.

ROSES: Black Spot, Powdery Mildew, Rust. — Prune and destroy all old leaves. Spray in the spring before buds begin to swell.

PEARS: Blister Mite. — Apply in the fall before the leaves start to turn color and again in the spring just as the buds begin to swell.

BLACKBERRIES, RASPBERRIES: Leaf Spot, Cane Blight, Powdery Mildew. — Prune out old canes and destroy before fall rains. Apply in late fall and again in the spring as the buds begin to swell.

Apply spraying on painted surface as staining may occur. Keep bottle tightly closed when not in use. If a crust should form on the surface break through the crust and shake thoroughly. Product quality is not impaired by crystallization.

STORAGE AND DISPOSAL

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area. Do not store diluted spray.

DISPOSAL: **PRODUCT:** Partially filled bottle may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash. **CONTAINER:** Do not reuse empty bottle. Rinse thoroughly before discarding in trash.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Chevron Chemical Company
Ortho Consumer Products Division
P.O. Box 400, CA 94502-0400
P.O. Box 400, CA 94502-0400



ORTHO

Dormant Disease Control

(LIME SULFUR FUNGICIDE) PREVENTS PEACH LEAF CURL
Also Aids in Control of Black Spot and Powdery Mildew on Roses

