

1070
P1021
239-2391

ORTHO Dormant Disease Control
EPA Reg. No. 239-2391
Submitted: March 21 1986

COPY A

COMBINATION WINTER SPRAY - Fruit Trees, Roses and Deciduous (no leaves in winter) Hedge Plants, Shade Trees and Shrubs (Dormant, deciduous only) - San Jose Scale, Rose Scale, Oyster Shell Scale, Brown Apricot Scale, Black Scale, Overwintering Insect Eggs, Many Overwintering Fungus Spores or Plant Diseases: FOR BEST RESULTS use 1½ pts. ORTHO Dormant Disease Control and ½ pt. VOLCK Oil Spray in 6 gals. water (7½ Tablespoonfuls ORTHO Dormant Disease Control and 2½ Tablespoonfuls VOLCK Oil Spray per gallon). Keep agitated during spraying.

COPY B

If swallowed, give water or milk to drink and telephone for medical advice. DO NOT make person vomit unless directed to do so by medical personnel. If medical advice cannot be obtained, then take person and product container to the nearest medical emergency treatment center or hospital. If inhaled, remove person from exposure area.

CONFIDENTIAL
PROPERTY OF
CHEVRON CHEMICAL COMPANY
UNTIL ACCEPTED BY EPA

ACCEPTED
JUL 15 1986
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 239-2391



ORTHO Dormant Disease Control

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER: Corrosive to eyes. Concentrate caused eye damage and skin irritation. Harmful if swallowed or absorbed through the skin. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Avoid breathing spray mist. **STATEMENT OF PRACTICAL TREATMENT:** In case of eye contact, immediately flush eyes with fresh water for 15 minutes, and see a doctor. Wash skin thoroughly with soap and water. If swallowed, give large quantities of water, induce vomiting and see a doctor immediately. Note to Physicians: Emergency Information—call (415) 233-3737.

ENVIRONMENTAL HAZARDS: Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. **PHYSICAL OR CHEMICAL HAZARDS: MAY GIVE OFF HIGHLY TOXIC AND EXTREMELY FLAMMABLE HYDROGEN SULFIDE GAS IF MIXED WITH ACIDS.**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

ORTHO Dormant Disease Control helps to prevent foliage diseases from occurring during the growing season by killing disease causing spores that over-winter on buds, twigs and bark and by providing a protective covering of the buds before they open in the spring. Since many of the disease organisms covered on this label also over-winter on old leaves, they should be raked up in the fall and disposed of.

DIRECTIONS

Apply ORTHO Dormant Disease Control to thoroughly cover all exposed plant parts. Use at the rate of 3/4 Tablespoons (5 fl. oz.) per gallon of water in an ORTHO SPRAYETTE, tank type or power sprayer. Do not spray when the temperature is over 80°F or under 45°F. Make new dilution for each use.

PEACHES: Peach Leaf Curl, Peach Blight—Timing is very important. Late Fall—Spray in October or November after leaf drop but before heavy rains occur. Mid-Winter—Apply a full coverage spray in January or February before buds begin to swell. Note: If buds have begun to swell or open, it is too late to obtain satisfactory control of Peach Leaf Curl Disease as infection has already occurred. For best results we recommend a combination of Late Fall and Mid-Winter applications.

ROSES: Black Spot, Powdery Mildew, Rust—Prune and destroy all old leaves. Spray in the spring before buds begin to swell.

PEARS: Blister Mite—Apply in the fall before the leaves start to turn color and again in the spring just as the buds begin to swell.

BLACKBERRIES, RASPBERRIES: Leaf Spot, Cone Blight, Powdery Mildew—Prune out old canes and destroy before fall rains. Apply in late fall and again in the spring as the buds begin to swell.

Avoid spraying on painted surface as staining may occur. Keep bottle tightly closed when not in use. If a crust should form on the surface, break through the crust and shake thoroughly. Product quality is not impaired by crystallization.

STORAGE AND DISPOSAL

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area. Do not store diluted spray.

DISPOSAL: PRODUCT—Partially filled bottle may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash.

CONTAINER—Do not reuse empty bottle. Rinse thoroughly before discarding in trash.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Chevron Chemical Company © 1984

Ortho Consumer Products Division
San Francisco CA 94120-7144
Richmond CA 94804-0036
Product 5154 Made in U.S.A.
Form 8484-H

EPA Est. 239-CA-1, 239-MO-1
EPA Reg. No. 239-2391-AA



(LIMITED)
Also

