



ORTHO Dormant Disease Control

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER: Corrosive to eyes. Concentrate causes eye damage and skin irritation. Avoid if swallowed or absorbed through the skin. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Avoid breathing spray mist. **STATEMENT OF PRACTICAL TREATMENT:** In case of eye contact, immediately flush eyes with fresh water for 15 minutes and see a doctor. Wash skin thoroughly with soap and water. If swallowed, give water or milk to drink and telephone for medical advice. **DO NOT** make person vomit unless directed to do so by medical personnel. If medical advice cannot be obtained, then take person and product container to nearest medical emergency treatment center or hospital. If inhaled, remove person from exposure area. **Note to Physicians:** Emergency Information—call (415) 233-3737. **ENVIRONMENTAL HAZARDS:** Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. **PHYSICAL OR CHEMICAL HAZARDS: MAY GIVE OFF HIGHLY TOXIC AND EXTREMELY FLAMMABLE HYDROGEN SULFIDE GAS IF MIXED WITH ACIDS.**

DIRECTIONS FOR USE

is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THE LABEL USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS

ORTHO Dormant Disease Control helps to prevent foliage diseases from occurring during the growing season by killing disease causing spores that overwinter on buds, twigs and bark and by providing a protective covering of the buds before they open in the spring. Since many of the disease organisms covered on this label also overwinter on old leaves, they should be raked up in the fall and disposed of.

DIRECTIONS

Apply ORTHO Dormant Disease Control to thoroughly cover all exposed plant parts at the rate of 10 Tablespoons (5 fl. oz.) per gallon of water in an ORTHO SPRAY-ITE tank type or power sprayer. Do not spray when the temperature is over 60°F or when the foliage is new growth for each use.

PEACH LEAF CURL: Peach Blight. Timing is very important. Late fall—begin in October or November after leaf drop but before heavy rains occur. Mid-fall—Apply a full coverage spray in January or February before buds begin to swell. **Note:** If buds have begun to swell or open, it is too late to obtain satisfactory control of Peach Leaf Curl Disease as infection has already occurred.

For best results we recommend a combination of Late Fall and Mid Winter applications.

ROSES: Powdery Mildew, Rust— Prune and destroy old leaves. Spray in the spring before buds begin to swell.

LEAVES: Blister Mite— Apply in the fall before the leaves start to turn color and again in the spring just as the buds begin to swell.

BLACKBERRIES, RASPBERRIES: Leaf Spot, Cane Blight, Powdery Mildew— Prune out old canes and destroy before late rains. Apply in late fall and again in the spring as the buds begin to swell.

Avoid spraying on painted surfaces as staining may occur. Keep bottle tightly closed when not in use. If a crust should form on the surface, break through the crust and shake thoroughly. Product quality is not impaired by crystallization.

STORAGE AND DISPOSAL


STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area. Do not store diluted spray.

DISPOSAL: PRODUCT— Partially filled bottle may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash. **CONTAINER—** Do not reuse empty bottle. Rinse thoroughly before discarding in trash.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Chevron Chemical
Company

Ortho Consumer Products Division
San Francisco CA 94120 7144
Product 5514 Made in U.S.A.
Form E184-1
EPA Est. 233 MO 1
EPA Reg. No. 233-2331-AA



BEST AVAILABLE COPY



ORTHO

Dormant Disease Control

(LIME SULFUR FUNGICIDE) PREVENTS PEACH LEAF CURL
Also Aids in Control of Black Spot and Powdery Mildew on Roses




Active Ingredient
Calcium Polysulfide
Inert Ingredients
alpha to 1133 Tetramethylphenyl
alpha to 1133 Hydroxyethylphenyl
Density Baumé at 60°F

By Wt.
26%
74%

Keep out of reach of children
DANGER
See side panel for additional
precautionary statements.


NET CONTENTS 1 QT.



COMBINATION WINTER SPRAY - Fruit Trees, Roses and Deciduous (no leaves in winter) Hedge Plants, Shade Trees and Shrubs (Dormant, deciduous only) - San Jose Scale, Rose Scale, Oyster Shell Scale, Brown Apricot Scale, Black Scale, Overwintering Insect Eggs, Many Overwintering Fungus Spores or Plant Diseases: FOR BEST RESULTS use 5 Tbs. ORTHO Dormant Disease Control and 5 Tbs. VOLCK Oil Spray in 1 gal. water. Keep agitated during spraying.

COPY B

If swallowed, give water or milk to drink and telephone for medical advice. DO NOT make person vomit unless directed to do so by medical personnel. If medical advice cannot be obtained, then take person and product container to the nearest medical emergency treatment center or hospital. If inhaled, remove person from exposure area.





ORTHO Dormant Disease Control

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER: Contains a highly concentrated caustic base. Avoid contact with skin, eyes, or clothing. Wash exposed areas thoroughly with soap and water. If contact occurs, flush with water for 15 minutes and see a doctor. Wash the exposed area with water for 15 minutes and see a doctor. Wash the exposed area with water for 15 minutes and see a doctor. Wash the exposed area with water for 15 minutes and see a doctor.

ENVIRONMENTAL HAZARDS: Do not apply directly to water. Do not apply where run-off may occur. Do not apply when weather conditions favor drift from areas treated. **PHYSICAL OR CHEMICAL HAZARDS:** MAY GIVE OFF HIGHLY TOXIC AND EXTREMELY FLAMMABLE HYDROGEN SULFIDE GAS IF MIXED WITH ACIDS.

DIRECTIONS FOR USE

Use in accordance with label directions and use that product in a manner inconsistent with its labeling.

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS

ORTHO Dormant Disease Control helps to prevent foliage diseases from occurring during the growing season by killing dormant spores that are on leaves, buds, and bark of the growing plant before they begin to grow. Since many of the diseases are carried on the foliage of the plant, they should be killed up in the fall and destroyed.

DIRECTIONS

Apply ORTHO Dormant Disease Control to thoroughly cover all exposed plant parts. Use the rate of 1/2 to 1/4 ounce per gallon of water in a spray or in a dust. Use the rate of 1/2 to 1/4 ounce per gallon of water in a spray or in a dust. Use the rate of 1/2 to 1/4 ounce per gallon of water in a spray or in a dust.

PEACHES: Peach Leaf Curl, Peach Blight - Young in very important. Late Fall - Spray in October or November after last drop but before heavy rains occur. Mid-Winter - Apply a full coverage spray in January or February before buds begin to swell. Note: If buds have begun to swell or open, use the rate to obtain satisfactory control of Peach Leaf Curl disease if infection has already occurred.

ROSES: Black Spot, Powdery Mildew, Rust - Fall and destroy of old leaves. Spray in the spring before buds begin to swell.

PEARS: Blight, Rust - Apply in the fall before the leaves start to turn color and again in the spring just as the buds begin to swell.

BLACKBERRIES, RASPBERRIES: Leaf Spot, Canine Blight, Powdery Mildew - Plant in the fall and destroy of old leaves. Apply in late fall and again in the spring as buds begin to swell.

Leaf spraying on painted surfaces as staining may occur. Keep these surfaces covered with paint. If a crust should form on the surface, break it off, wash the surface and shake thoroughly. Product quality is not impaired by crystallization.

STORAGE AND DISPOSAL

STORAGE: Keep product in original container. Do not put contents of container in food or drink containers. Avoid contact with skin and clothing. Store in a cool, dry place. Do not store in a hot place. Do not store in a hot place.

DISPOSAL: **PRODUCT:** - Pour into a container with a cap and dispose of it in a safe place. **CONTAINER:** - Do not reuse empty container. Rinse thoroughly before discarding in trash.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Chevron Chemical
Company © 1981

Ortho Chemical Products Division
San Francisco, CA 94120
Rt. 1, Box 1000
P.O. Box 1000
P.O. Box 1000

EPA Reg. No. 720-2701 AA



BEST AVAILABLE COPY

Dise

(LIME SULFUR FU
Also Aids in Cont



Peach Leaf Curl