




ORTHO

Dormant Disease Control

(LIME SULFUR FUNGICIDE) PREVENTS PEACH LEAF CURL
Also Aids in Control of Black Spot and Powdery Mildew on Roses



Active Ingredient
Calcium Polysulfide 25%
Inert Ingredients 75%
alpha (p-t) 1,3,3-tetramethylbutylphenyl,
omega-hydroxypolyoxyethylene 10%
Density Approx at 60 29

Keep out of reach of children
DANGER
See side panel for additional
precautionary statements.

NET CONTENTS 1 QT.

11 11

11 11

11 11

2.

3/23 3/24/87



ORTHO Dormant Disease Control

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER:

Irritative to eyes. Concentrate causes eye damage and skin irritation. Harmful if swallowed or absorbed through the skin. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Avoid breathing spray mist. **STATEMENT OF PRACTICAL TREATMENT:** In case of eye contact immediately flush eyes with fresh water for 15 minutes and see a doctor. Wash skin thoroughly with soap and water. If swallowed, give water or milk to drink and telephone for medical advice. DO NOT make person vomit unless directed to do so by medical personnel. If inhaled, get fresh air. If inhaled, then take person and product container to the nearest hospital emergency treatment center or hospital. If inhaled, remove person from exposure area. **Note to Physicians:** Emergency Information: Call 415-239-3737.

ENVIRONMENTAL HAZARDS: Do not apply directly to water. Do not apply where run-off is expected. Do not apply when weather conditions favor drift from areas treated. **PHYSICAL OR CHEMICAL HAZARDS:** MAY GIVE OFF HIGHLY TOXIC AND EXTREMELY FLAMMABLE HYDROGEN SULFIDE GAS IF MIXED WITH ACIDS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS

ORTHO Dormant Disease Control helps to prevent foliage diseases from occurring during the growing season by killing disease causing spores that overwinter on buds, twigs and bark and by providing protective covering of the buds before they open in the spring. Some of the plant diseases covered on this label also overwinter on old leaves. These old leaves should be removed from the tree and disposed of.

Apply ORTHO Dormant Disease Control to the foliage of the plant, exposing the entire plant. Use at the rate of 10 Tbs. (5 fl. oz.) per gal. of water. ORTHO Dormant Disease Control is a water soluble concentrate. It should be applied before the temperature is over 80° F. or under 45° F. Make new solution if needed.

PEACHES: Peach Leaf Curl, Peach Blight, Shot Hole Fungus. Timing is very important. **Late Fall:** Apply ORTHO Dormant Disease Control to the foliage of the tree before heavy frost. **Mid Winter:** Apply ORTHO Dormant Disease Control to the foliage of the tree before buds begin to swell. For best results, apply ORTHO Dormant Disease Control to the foliage of the tree before buds begin to swell. For best results, apply ORTHO Dormant Disease Control to the foliage of the tree before buds begin to swell. For best results, apply ORTHO Dormant Disease Control to the foliage of the tree before buds begin to swell.

ROSES: Black Spot, Powdery Mildew, Rust. Remove and destroy old leaves. Spray in the spring before the buds begin to swell.

PEARS: Blister Mite. Apply in the fall before the leaves start to turn color and again in the spring before the buds begin to swell.

BLACKBERRIES, RASPBERRIES: Leaf Spot, Cane Blight, Powdery Mildew. Remove old canes and destroy before fall rains. Apply in late fall before the leaves begin to swell.

COMBINATION WINTER SPRAY: Fruit Trees, Roses and Deciduous tree leaves in winter. Hedge Plants, Shade Trees and Shrubs (Dormant, deciduous only). San Jose Scale, Rose Scale, Doryphora Scale, Brown Apricot Scale, Black Scale, Overwintering Eggs, Many Overwintering Fungus Spores or Plant Diseases. For best results, use 5 Tbs. ORTHO Dormant Disease Control and 5 Tbs. VOOCH (Vine Oil) per gallon of water. Isolate during spraying.

Avoid spraying on painted surface as staining may occur. Do not spray on plants that are not in use. If a crust should form on the surface, break it off and spray again. Spray thoroughly. Product quality is not impaired by crystallization.

STORAGE AND DISPOSAL

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute food or drink containers. Avoid contamination of feed and animal stuffs. Store in a dry place, preferably in a locked storage area. Do not store in a heated space.

DISPOSAL: Partially filled bottle may be disposed of by burying in a deep hole or by pouring in several layers of sawdust or other absorbent material. **CONTAINER:** Do not reuse empty bottle. Rinse thoroughly before disposal.

NOTICE: Buyer assumes all responsibility for safety of use. Not for sale where prohibited.

Chevron Chemical Company © 1984

Orth