



ORTHO



ORTHO

Covers Up To
12,800 Sq. Ft.

WEED-B-GON[®]
For Southern
Grasses

WEED-B-GON[®]
For Southern
Grasses

CONTROLS

Black Medic
Chickweed, Clover
Cudweed, Dichondra
Evening Primrose
Pigweed
Bracted Plantain
Purslane
Shepherdspurse
Sow Thistle
Toadflax
Florida Betony
Florida Pusley
Oxalis
Sensitive Mimosa
Spurge
Wild Geranium

FOR CONTROL OF WEEDS

in St. Augustine, Bermuda, Zoysia,^{*}
Centipede and Bahiagrass Lawns

Active Ingredients

Isobutyl ester of 2-Methyl-4-chlorophenoxypropionic acid

Methyl 2-(3-hydroxyfluorene-9-carboxylate)

Methyl 9-hydroxyfluorene-9-carboxylate

Methyl 2-(3,4-dichloro-5-hydroxyfluorene-9-carboxylate)

Inert Ingredients

*Equivalent to 11.65% MCPP, 2-(2-Methyl-4-chlorophenoxy)propionic acid

Patent Pending

By Wt.

17.13%

26%

1.65%

0.4%

78.4%

CAUTION: Keep out of reach of children.
See back panel for additional cautions.

NET CONTENTS 1 QUART



ORTHO



ORTHO

WEED-B-GON[®]
For Southern Grasses

WEED-B-GON[®]
For Southern
Grasses

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

WEED-B-GON For Southern Grasses containing MCPP and Maintain is designed to control broadleaf weeds in southern or warm season grasses without injuring the lawngrass when used as directed. Do not store diluted spray.

DIRECTIONS FOR USE ON LAWNS: For best control, WEED-B-GON For Southern Grasses should be applied in the spring or fall when weeds are young and actively growing. If summer applications are necessary, temporary discoloration or injury may occur to St. Augustine Grass; however, St. Augustine and other southern grasses will not be permanently injured if applied as directed.

Make one application to control Black Medic, Chickweed, Clover, Cudweed, Dichondra, Evening Primrose, Pigweed, Bracted Plantain, Purslane, Shepherds-purse, Sow Thistle and Toadflax.

Make 2 applications 2-3 weeks apart to control the following hardy weeds: Florida Betony, Florida Pusley, Oxalis, Sensitive Mimosa, Spurge and Wild Geranium.

HOW TO USE: For best results, use an ORTHO SPRAYETTE or tank type sprayer. Refer to table for recommended dilution. If lawn needs watering, water thoroughly before application. Do not water for 24 hours after application. Spray entire lawn, wetting weeds and lawn grasses; do not over-spray. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants.

Note: The activity of WEED-B-GON For Southern Grasses may not be evident for at least 2 to 3 weeks after application and complete control of listed weeds may take 4-6 weeks.

NEWLY SEEDED LAWNS may be treated after they have been mowed at least 3 times. Do not apply to Dichondra, Lippia, Clover, Bluegrass or Fescue Lawns.

DILUTION TABLE

Area to Treat	Water To Use	Amount of WEED-B-GON For Southern Grasses To Use
200 sq. ft.	1 gallon	1 Tablespoonful
4200 sq. ft.	6 gallons	6 Tablespoonfuls (3 ounces)

IMPORTANT: Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Vapors may injure susceptible plants in immediate vicinity. Do not reseed for 3 to 4 weeks after use.

CAUTION: Harmful if Swallowed. Do not breathe vapor or spray mist. Avoid contact with skin, eyes and clothing. In case of contact, flush with plenty of water. For eyes, get medical attention. Wash hands after use. Keep children and animals away from treated areas until these areas are dry.

Note to Physicians: Emergency information - call (415) 233-3737. **DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.** Avoid spray contact or drift to susceptible plants as this product may injure flowers, shrubs, garden vegetables, fruit trees and other crops. Coarse sprays are less likely to drift. When container is empty, immediately wash thoroughly and destroy. Never re-use. Do not store near feed, foodstuffs, fertilizers, seeds, insecticides or fungicides. When ORTHO Sprayer is used and rinsed thoroughly as directed on sprayer jar immediately after applying WEED-B-GON For Southern Grasses, it can be used to apply insecticides, fungicides and liquid fertilizers to ornamental plants.

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions. Contains Chlorfuretol, a Product of E. Merck, Darmstadt, West Germany.

Chevron Chemical Company

Ortho Division, San Francisco, CA 94119
Rm. 400, CA 94804
Phone: 936-3141, Prod. 5277, Major - S.A.

EPA Est. 1169-NJ-1
EPA Reg. No. 239-2403