

239-2423

12/9/1974

1/2

Lawn Insect Spray

Treats up to 4,000 Sq. Ft.

Contains Dursban*

For Grass and Dichondra Lawns

Controls: Armyworms, Sod Webworms (Lawn Moth Larvae), Chinchbugs, Dichondra Flea Beetle, Fiery Skipper Larvae, Cutworms, Crickets, Earwigs, Grasshoppers, Brown Dog Ticks and certain other insects.

Active Ingredients:	By Wt.
Chlorpyrifos [O,O-diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate]	5.3%
Aromatic petroleum derivative solvent	3.0%
Petroleum distillate	87.0%
Inert Ingredients	4.7%
U.S. Patent No. 3,244,586	

*Dursban® is a registered trademark of The Dow Chemical Company

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

DIRECTIONS

Place required amount of ORTHO Lawn Insect Spray in ORTHO Lawn Sprayer jar (see chart). Fill jar with water, mix, attach to sprayer and apply as recommended. Use at once. For best results the lawn should be moist at time of application and care should be taken to obtain complete and uniform coverage of pest infested areas. Spray when pests first appear. Re-treat when needed.

NET CONTENTS 1 QUART LIQUID

Chevron Chemical Company

Ortho Division/San Francisco CA 94119 Richmond CA 94804

Product 5284 Made in U.S.A.

Form 8863-A

EPA Reg. No. 239-2423-AA

NOTE: Mosquitoes coming to rest on areas treated for the control of Lawn and Home Invading Pests will be killed for varying periods of time after treatment, depending on exposure of treated areas to weathering conditions.

NOTE: Do not use on azaleas, camellias, poinsettias, rose bushes or variegated ivy because of possible injury to these plants.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: May be harmful if swallowed or absorbed through the skin. Do not breathe vapors or spray mist. Avoid eye and skin contact. Wash thoroughly after handling. Keep children and pets off treated areas until spray has dried.

Note to Physicians: Emergency Information - call (415) 233-3737. Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is antidotal. Do not use on vegetables or food crops. Keep away from food, feedstuffs, and domestic water supplies. **DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.** Handle concentrate in a ventilated area. When container is empty, immediately wash thoroughly and destroy. Never re-use.

This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on plants. Protective information may be obtained from your Cooperative Agricultural Extension Service.

ORTHO

RECEIVED
12 NOV 15
REGISTRATION OFFICE
PESTICIDES, EPA

2/2

Pests	Amount Lawn Insect Spray to Use	Comments
Lawn Pests		
Armyworm* (Western U.S. only)	1/2 cup (4 fl. oz.) in 15 gals. water to treat 500 sq. ft. of lawn. One	Mow lawn and water well 1-2 days before treatment. Heavy watering or irrigation should be withheld for 2-3 days following treatment. Spray when pests first appear. Repeat treatments may be necessary at 4-6 week intervals. Sprays are most effective if applied during 2-hour period prior to darkness when lawn insects become active.
Lawn Moth Larvae (Sod Webworms)	tablespoon in 1 gal. covers 62 sq. ft. 1/2	
Dichondra Flea Beetle*	pt. in 30 gals. covers	
Chinchbugs	1,000 sq. ft.	
Cutworms*		
Fiery Skipper Larvae (California and Hawaii only)		
Tropical Grass Webworms (Hawaii only)		
Grasshoppers		
Crickets		
*including Dichondra Lawns		
White Grubs in Lawns		
Certain White Grubs: (Larvae of European chafer and Japanese Beetle)	2 cups (16 fl. oz.) in 15 gals. water to treat 500 sq. ft. Four tablespoons in 1 gal. covers 62 sq. ft. One quart in 30 gals. covers 1,000 sq. ft.	Water lawn thoroughly following application to wash the insecticide into the soil. For best results treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local Agricultural Extension Service Specialist for proper time to treat in your area.
Home Invading Pests (Outside Dwellings)		
Ants	1/2 cup (4 fl. oz.) in 15 gals. water to treat	To help prevent infestation of buildings, treat a band of soil 6-10 feet wide around and adjacent to the building, also the building foundation to a height of 2-3 feet when pests are active and may find entrance. Do not treat pets with this product.
Brown Dog Ticks	500 sq. ft. One table-	
Earwigs	spoon in 1 gal. covers	
Crickets	62 sq. ft. 1/2 pint in	
Sowbugs (Pillbugs)	30 gals. covers 1,000 sq. ft.	

ACCEPTED
239-2423
DEC 9 1974

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.