

**Chevron**  
**ORTHO Liquid Lawn Disease Control**

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS & DOMESTIC ANIMALS**  
**WARNING:** Causes eye and skin irritation. Skin contact may cause allergic reaction. Do not get in eyes, on skin or on clothing. Wear rubber gloves when handling. Wash skin and hands thoroughly after using. Remove and wash contaminated clothing before reuse. Keep children and pets off treated area until dry. **STATEMENT OF PRACTICAL TREATMENT:** In case of eye contact, immediately flush eyes with fresh water for 15 minutes and see a doctor. In case of skin contact, wash skin with soap and water. Get medical attention if irritation persists. **Note to Physicians:** Emergency information call (415) 233-3737. **ENVIRONMENTAL HAZARDS:** This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

**DIRECTIONS FOR USE**  
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

ORTHO Liquid Lawn Disease Control gives effective control of common disease of grass lawns including Brown Patch (*Rhizoctonia*), Helminthosporium Leaf Spot and Melting Out (*Helminthosporium*), Dollar Spot (*Sclerotinia*), Copper Spot (*Gloeosporangium*), Red Thread (*Corticium*), Curvularia Leaf Spot (*Curvularia*), Gray Leaf Spot (*Piricularia*), and Stem Rust (*Puccinia*) of Bluegrass, also Leaf Spot (*Alternaria*) and (*Anthraco*) in Dichondra lawns (California only).

**DIRECTIONS**  
 One day before applying ORTHO Liquid Lawn Disease Control, mow lawn, collect clippings and water if necessary. May be applied with an ORTHO SPRAYETTE or ORTHO Lawn and Garden Sprayer (Hollow setting) tank type or power sprayer. Shake bottle well before using. For best results when mowing, place a small amount of water in sprayer, add the appropriate amount of ORTHO Liquid Lawn Disease Control, then add water to bring to desired level and agitate gently to uniformly mix the solution. Apply spray evenly. Six gallons diluted spray treats 1,000 sq. ft. Agitate occasionally while spraying. Do not water or mow lawn for 48 hours after application. Make new dilution for each use. Do not apply when temperature is above 90 F to prevent slight burning or temporary discoloration of grass lawns. Do not graze treated areas or feed treated plant parts to livestock.

Copy A

MAY 27 1983  
 233-2432 Copy D

GRASS LAWN DISEASES	WINTER SEASON
Copper Spot ( <i>Gloeosporangium</i> )	70-90 F, low fertility
Curvularia Leaf Spot	Summer, 80-90 F, heavy thatch
Dollar Spot ( <i>Sclerotinia</i> )	70-80 F, low fertility
Gray Leaf Spot ( <i>Piricularia</i> )	Summer, high N, high humidity
Helminthosporium Leaf Spot & Melting Out	Summer, 80-90 F
Large Brown Patch ( <i>Rhizoctonia</i> )	Most, overcast
Red Thread ( <i>Corticium</i> )	Summer, 75-85 F, warming trend
Stem Rust of Bluegrass ( <i>Puccinia</i> )	Cool wet, 55-72 F
Dichondra Diseases - (CA only)	70-75 F, high light intensity, humidity
Alternaria Leaf Spot	Winter & Spring, rainy, cool, 55-72 F
Anthraco	Summer, 70-85 F

Mix in 1 qt. of Water to Cover 1,000 Sq. Ft.  
 (1) (Preventive\*), 4-8 fl. oz. (Curative\*\*) 8-16 fl. oz.  
 (2) (Preventive\*), 8-12 fl. oz. (Curative\*\*) 12-16 fl. oz.  
 Note: For 3 gal. rate, cut above rates in half and apply to cover 500 sq. ft.  
 \*Preventive Program: Begin applications when disease threatens or when conditions favor disease development. During periods of very wet or damp weather apply every 7 days. Use lower rates when disease conditions are light to moderate, higher indicated rates when conditions are severe.  
 \*\*Curative Program: If disease is already established, use a curative rate. Higher indicated rates should be applied under severe conditions. Make at least 3 successive applications at 7 day intervals. Continue treatments, using PREVENTIVE PROGRAM until lawn has recovered.

**STORAGE AND DISPOSAL:** Do not contaminate feed or foodstuffs. Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Do not reuse empty container. Wrap container and put in trash collection.

**NOTICE:** Buyer assumes all responsibility for safety and use not in accordance with directions.

**Chevron Chemical Company**  
 Ortho Consumer Products Division  
 San Francisco CA 94119  
 Richmond CA 94804  
 Product 1000 Made in U.S.A.  
 Form 8967-G  
 EPA Reg No. 233-2432-AA  
 EPA Est. 239 MD 1




BEST DOCUMENT AVAILABLE

**Chevron**  
**ORTHO**  
**Liquid Lawn Disease Control**

Copy C  
 Treats up to 6000 sq. ft.

Contains DACONIL 2787\* — A Broad-Spectrum Liquid Fungicide for Use on Grass and Dichondra Lawns. Controls Lawn Diseases: Brown Patch, Leaf Spots, Melting Out, Red Thread, Copper Spot, Dollar Spot and Rust in Grass Lawns; and Alternaria Leaf Spot and Anthracnose in Dichondra Lawns



Active Ingredient	By Wt
**Chlorothalonil (tetrachloro-5-phthalonitrile)	29.6%
Inert Ingredients	70.4%

\*DACONIL 2787 is a registered trademark of Diamond Shamrock Corporation  
 \*\*Covered under U.S. Patent No. 3,290,353 and No. 3,331,735

Keep out of reach of children  
**WARNING**  
 See side panel for additional precautionary statements.  
**NET CONTENTS 1 QT.**

ACCEPTED

MAY 27 1983

239-2432

ORTHO Liquid Lawn Disease Control  
EPA Reg. No. 239-2432  
APR 19 1983

COPY A

"IMPORTANT: ORTHO Liquid Lawn Disease Control is effective on a wide range of diseases, however proper identification of specific lawn disease is essential for control. Take a specimen of the disease to a professional garden center or contact the local County Extension Service for positive identification before treatment."

COPY B

"FOR ESTABLISHED LISTED DISEASES AND SEVERE CONDITIONS - Apply 12 fl. oz. (1½ cups) in 6 gallons of water to cover 1,000 sq. ft. During periods of very wet or damp weather, make 3 successive applications at 7 day intervals. Continue treatments following rate and intervals listed below."

FOR LIGHT DISEASE INCIDENCE AND MILD CONDITIONS - Apply 8 fl. oz. (1 cup) in 6 gallons of water to cover 1,000 sq. ft. Apply at a 10-14 day interval depending on severity of disease and conditions favoring disease development until lawn has recovered. NOTE: For 500 sq. ft. of lawn, use half the listed amount of product in 3 gallons of water.

COPY C

Treats up to 4,000 sq. ft.

COPY D

CONDITIONS

**CONFIDENTIAL**  
**PROPERTY OF**  
**CHEVRON CHEMICAL COMPANY**  
**UNTIL ACCEPTED BY EPA**