

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

WARNING: Causes eye and skin irritation. Skin contact may cause allergic reaction. Do not get in eyes, on skin or on clothing. Wear rubber gloves when handling. Wash skin and hands thoroughly after using. Remove and wash contaminated clothing before reuse. Keep children and pets off treated area until dry.

STATEMENT OF PRACTICAL TREATMENT: In case of eye contact, immediately flush eyes with fresh water for 15 minutes and see a doctor. In case of skin contact, wash skin with soap and water. Get medical attention if irritation persists. **Note to Physicians:** Emergency Information—call (415) 233-3737.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

ORTHO Vegetable & Garden Fungicide mixes readily with water and may be used in an ORTHO SPRAYETTE, ORTHO Lawn & Garden Sprayer, tank type or power sprayer. Shake bottle well before using. Mix appropriate amount with water and use at once.

Apply at listed rate per crop for disease to be controlled () in 1 gallon of water to cover an area of 200 sq. ft. (10' x 20') planted with listed crop. 1 TBS—1 Tablespoonful, 1 tsp—1 teaspoonful. Repeat at 7 to 14 day intervals. Use the 7 day schedule during severe disease conditions or wet weather unless otherwise stated. Note: Avoid spraying plants during extremely hot and sunny weather. Do not apply to wilted plants.

Do not graze treated areas or feed treated plant parts to livestock.

VEGETABLES—Numbers in () indicate days required between last application and harvest.

Beans (14) (Snap Beans): Rust [1 TBS.]—Begin in early bloom stage or when disease first threatens. Repeat at weekly intervals or as necessary.

Broccoli (0), Brussels Sprouts (0), Cabbage (0), Cauliflower (0), Alternaria Leafspot (Southeast only) [1½ tsp.], Downy Mildew [1½ tsp.]—Begin after transplants are set in the field or when conditions favor disease development. Repeat at 7 to 10 day intervals. **Ring Spot on Brussels Sprouts (California only) [2 tsp.]**—Begin at time of early sprout development or when conditions favor disease development. Repeat at 7 to 10 day intervals.

Carrots (0): Early and Late Blight [2 tsp.]—Begin when disease threatens and repeat at 7 to 10 day intervals or as required to maintain control.

Celery (7): Early and Late Blight [1½ tsp.], Bacterial Stalk Rot [1½ tsp.]—Begin when plants are set in the field. Repeat on a 4 to 7 day interval.

Corn (14) (Sweet): Helminthosporium Leaf Blight [2 tsp.]—Begin when conditions favor disease development. Repeat at a 4 to 7 day interval or as needed. Do not apply to sweet corn for processing.



ORTHO

Makes up to
128 Gallons
Diluted Spray

Vegetable & Garden Fungicide

Contains DACONIL 2787*

For the Control of Diseases on Vegetables, Roses
and Flowers
(as listed on label)

CONTROLS BLIGHT ON TOMATOES

Active Ingredient	By Wt.
**Chlorothalonil (tetrachloroisophthalonitrile)	29.6%
Inert Ingredients	70.4%
*DACONIL 2787 is a registered trademark of Diamond Shamrock Corp.	
**Covered under U.S. Patent No. 3,290,353 and No. 3,331,735	

Keep out of reach of children

WARNING

See side panel for additional precautionary statements.

NET CONTENTS 1 QT.

Cucumbers (0), Melons (0), Pumpkins (0), Squash (0): Anthracnose [2 tsp.], Gummy Stem Blight [2 tsp.], Scab [2 tsp.], Downy Mildew [2 tsp.], Leaf Blight [2 tsp.], Cercospora Leafspot (except Cucumbers) [2 tsp.], Powdery Mildew (except Southwestern States) on Cucumbers and Melons [2 tsp.]. Also suppresses Fruit Rot (Rhizoctonia) [1 TBS.] on Cucumbers—Begin when plants are in the first true leaf stage or when conditions are favorable for disease. Repeat applications at 7 day intervals or more frequently if required.

Potatoes (0): Early and Late Blight [1½ tsp.], Botrytis Vine Rot [1½ tsp.]—Begin when plants are 6 inches or when disease threatens. Repeat at 7 to 10 day intervals or as needed.

Tomatoes (0): Anthracnose [1 TBS.], Early and Late Blight [2 tsp.], Gray Leafspot [2 tsp.], Gray Mold (Botrytis) [1 TBS.], Rhizoctonia Fruit Rot [1 TBS.]—Begin when disease threatens. Repeat at 7 to 10 day intervals or more frequently under severe conditions. **Alternaria Fruit Rot or Black Mold (California only) [1 TBS.]**—Make 3 applications on a 10-14 day schedule beginning 2-6 weeks before anticipated harvest.

ROSES, FLOWERS

Roses: Blackspot [1 tsp.], Blossom Blight [1 tsp.]—Apply when young leaves first appear. Continue spray program throughout the growing season. Spray weekly during rainy seasons.

Chrysanthemums: Ascochyta Ray Blight [1½ tsp.], Gray Mold [1½ tsp.]—Apply at first sign of disease. Repeat as needed through growing season.

Geraniums: Botrytis Blight [1½ tsp.]—Use the 7 day schedule during wet weather or severe disease conditions.

Iris: Botrytis Blight [1½ tsp.], Leafspot [1½ tsp.]—Begin applications when leaves are 4" to 6" high.

Zinnias: Powdery Mildew (Erysiphe) [1½ tsp.]—Begin applications at the first sign of disease.

STORAGE AND DISPOSAL

Do not contaminate feed or foodstuffs.
Keep pesticide in original container.
Do not put concentrate or dilute into food or drink containers.
Do not store diluted spray.
Do not reuse empty container.
Wrap container and put in trash collection.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Chevron Chemical Company

Ortho Consumer Products Division
San Francisco CA 94120-7144

Product 1000 Made in U.S.A.

Form 9067-F

EPA Reg. No. 239-2438

EPA Est. 239-CA-1, 239-NJ-1, 239-MO-1
EPA Reg. No. 239-2438

ACCEPTED
RICHMOND, CA 94806
0036

MAY 27 1983

THE R. L. B. GROUP

1235 2438