



ACCEPTED

APR - 5 1982

U.S. Environmental Protection Agency  
Registration and Pesticide Control  
Washington, D.C. 20460  
EPA Reg. No. 5481-130 AA

## Fruit Fix Super Concentrate 800

A hormone type product for control of  
preharvest drop of apples and pears.

ACTIVE INGREDIENT  
1-Naphthaleneacetic Acid, Ammonium salt\* ..... 21.40%  
INERT INGREDIENTS: ..... 78.60%  
Total 100.00%

\*Equivalent to 19.5% 1-Naphthaleneacetic Acid or 800 grams (1.76 lbs) per gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR ADDITIONAL  
PRECAUTIONARY STATEMENTS

NOTICE: This product conforms to its chemical description and is reasonably fit for the purpose stated on the label, when used in accordance with directions under normal conditions of use. Manufacturer is not responsible for the use of this product contrary to the label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

AMVAC CHEMICAL CORPORATION  
Los Angeles, California

EPA EST. NO. 5481-011      EPA REG. NO. 5481-130 AA

NET CONTENTS

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

#### CAUTION

May be harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes or clothing. In case of eye contact, immediately flush eyes with plenty of water. Get medical attention if irritation persists. Wash thoroughly after handling. Remove and wash contaminated clothing before reuse.

#### ENVIRONMENTAL HAZARDS

Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Avoid drift of spray vapors.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Storage:** Do not store in direct sunlight. Do not store at temperatures of 32 F or below.

**Pesticide Disposal:** Pesticide, spray mixture or rinse water that cannot be used according to label instructions may be disposed of according to Federal, State or local procedures.

**Container Disposal:** Triple rinse (or equivalent) and dispose of in a sanitary land fill or by other approved State and local procedures.

FRUIT FIX SUPER CONCENTRATE 800 serves to control preharvest drop of fruit from windfall and breakdown. FRUIT FIX SUPER CONCENTRATE 800 is highly variable and temperature sensitive. Heavy fruit set is expected. FRUIT FIX SUPER CONCENTRATE 800 is highly variable and temperature sensitive. Heavy fruit set is expected. FRUIT FIX SUPER CONCENTRATE 800 is highly variable and temperature sensitive. Heavy fruit set is expected.

#### Directions for Apples and Pears

Spray leaves thoroughly, wetting all fruit stems and adjacent leaf areas, 5 to 7 days before harvest for pears. Do not spray within 5 days of harvest and control drop for 10 days to 4 weeks (less for McIntosh) than 2 applications and do not delay picking beyond optimum maturity. **GROUND SPRAY**—For apples, thoroughly mix 3 fl. ozs. with 250 gal. of water. For pears, use a 100 ppm spray obtained by mixing the appropriate spray using up to 670 gallons per acre of this 1-Naphthaleneacetic Acid per acre. Apply to fruit 5 to 7 days before harvest. **AIRPLANE SPRAY**—Mix 5 1/2 fl. ozs. with 3 gallons of water and apply to fruit 5 to 7 days before harvest. Apply 5 gallons of emulsion per acre of apples. For pears, apply 5 gallons of emulsion per acre.

#### Directions for Thinning Olive Trees

**TIMING IS CRITICAL.** Spray when young fruits are 1/8" to 3/16" in diameter. Thinning is effective on all varieties except Seville. **WITHOUT OIL.** Apply a solution of 100 ppm 1-Naphthaleneacetic Acid in 100 gallons of water. Emulsify by agitation with 1 1/2 gallons of oil. Spray only the heavily loaded trees until the leaves are thoroughly wet. **CAUTION:** Unless you wish to remove all your fruit, do not apply.

#### Directions to Prevent Fruiting of Ornamental Citrus

For best results, apply when the olive trees are in full bloom but before the fruit are formed. Apply a second spray on 7 to 14 days later. Mix 8 ozs. Fruit Fix Super Concentrate 800 and a wetting agent until leaves are thoroughly wet. Application should not be made when temperatures are over 85 F. Use a paraffin sheet while spraying and the cover left on until possible injury to these plants.

#### Directions for Fruit Thinning Oranges, Tangerines

Spray at the rate of 100 to 500 ppm (6 fl. ozs. per 100 gals. water). FRUIT FIX SUPER CONCENTRATE 800 is highly variable and temperature sensitive. Heavy fruit set is expected. Do not apply to nonbearing trees. Time of application for his culture and location. Concentration made when the fruit are in an early stage of development. In Florida the period of natural drop occurs in 3/4. Heavy fruiting activity should be expected during hot weather. Heavy fruiting activity should be expected during hot weather. Heavy fruiting activity should be expected during hot weather. Heavy fruiting activity should be expected during hot weather.

—DO NOT APPLY MORE THAN 1 APPLICATION PER YEAR

—DO NOT USE ON VARIETIES WHERE BOTH YOUNG FRUIT AND  
—DO NOT APPLY WITHIN 150 DAYS OF HARVEST.

MICROFICHE CREATED

DATE:-----