

Reg # 239-2479

PM-13

496

ORTHO Diazinon Soil & Turf
Insect Control
EPA Reg. No. 239-2479

COPY A

(FOR BIODAC FORMULATION ONLY)

SPREADER MAKE AND MODEL	APPLICATION RATE OF PRODUCT 2 LBS. PER 1000 SQ. FT.
ORTHO Broadcast Spreader	3.5
ORTHO Drop Spreader (Model 5000)	5.0
Craftsman (Model 671-19198)	6.0
Scotts Precision Flow (Model PF 1)	5.0
Scotts Easy Green	25.5
Cyclone Rotary (Model 96-5)	5.0
True Temper (Model CB 1100)	3.0
Red Devil Lawn Drop Spreader (Model FH-22N)	9.0
Red Devil Broadcast Spreader (Model FH KD 2N)	6.0

HAND HELD SPREADERS

NO. HOPPERS TO DELIVER
2 LBS. PER 1000 SQ. FT.

ORTHO WHIRLYBIRD®	1.0
Handy Spreader (Pennington)	0.375
Handy Green (Scotts)	0.75
Red Devil	0.375

EPAREG.RAM

ACCEPTED
JAN 13 1993

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, this product is registered under EPA Reg. No. 239-2479

BEST AVAILABLE COPY

ORTHO Diazinon Soil & Turf
Insect Control
EPA Reg. No. 239-2479

246

COPY B

(FOR BIODAC FORMULATION ONLY)

Ants: Apply 1 teaspoon of ORTHO Diazinon Soil & Turf Insect Control over and around each ant hill.

COPY C

(FOR BIODAC FORMULATION ONLY)

Imported Fire Ants: To aid in the control of Imported Fire Ants in lawns and other outdoor home recreational areas, apply 1/3 cup in a 2 ft. radius over and around each fire ant mound. Immediately after application, water in thoroughly. A minimum of 1 gallon of water should be applied to each mound area. Do not use more than 2 gallons of water. Apply water gently to avoid disturbing ants. Use of high pressure watering may disturb the ants and cause migration, thus reducing effectiveness of treatment. Treat new mounds as they appear. For best results, apply in cool weather, 65 to 80° F, or in early morning or late evening hours.

COPY D

(FOR BIODAC FORMULATION ONLY)

Digger Wasps: Apply 1 teaspoon of ORTHO Diazinon Soil & Turf Insect Control in and around each nest opening. Apply at dusk when wasps are not active.

COPY E

(FOR BIODAC FORMULATION ONLY)

NOTE: One level measuring cup (8 fl. oz.) holds 5 oz. of granules.

ACCEPTED
JAN 13 1993
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the purpose of
registered under
EPA Reg. No. 239-2479

**ORTHO Diazinon
Soil & Turf Insect Control
02591A**

3 7 6

TO OPEN
CUT SPOUT HERE



ORTHO Diazinon Soil & Turf Insect Control

CAUTION

PRECAUTIONARY STATEMENTS

Directions: For use on lawns and golf courses only. For use on golf courses, apply to fairways and tees only. For use on lawns, apply to all areas except flower beds, vegetable gardens, and areas where children and pets play. For use on golf courses, apply to all areas except greenhouses, tennis courts, and areas where children and pets play. For use on lawns, apply to all areas except flower beds, vegetable gardens, and areas where children and pets play.

First Aid: If swallowed, give water and call a doctor. If inhaled, get fresh air. If on skin, wash with soap and water.

Use Precautions: Do not use in enclosed areas. Do not use on areas where children and pets play. Do not use on areas where food is stored. Do not use on areas where livestock are kept. Do not use on areas where crops are grown. Do not use on areas where birds are kept.

Storage and Disposal: Store in original container. Do not store in areas where children and pets play. Do not store near food or feed. Do not store near livestock. Do not store near crops. Do not store near birds.

Other Precautions: Do not use on areas where children and pets play. Do not use on areas where food is stored. Do not use on areas where livestock are kept. Do not use on areas where crops are grown. Do not use on areas where birds are kept.

**ORTHO Diazinon
Soil & Turf Insect Control
02591A**

Covers up to 5,000 Sq. Ft.



ORTHO

**Diazinon
Soil & Turf
Insect Control**

ACCES
JAN
Under the Federal
Fungicide, and
as amended, is
registered under
EPA Reg. No.

This product must not be used on
golf courses and sod farms.

**Ready-to-use Granules For Use When Seeding or
Transplanting Vegetables**
**Kills: Wireworms, Rootworms, Root Maggots, Cutworms
and Other Listed Soil Insects**

Contains Diazinon

Ready-to-use granules for use only on:


- Home Lawns to Kill Billbugs, Chinch Bugs, Sod
Webworms and White Grubs, as listed.
- Home Vegetable Gardens to Kill Cutworms,
Wireworms, Root Maggots on listed crops.
- Nuisance Pests Outside the Home to Kill Ants, Fleas,
Brown Dog Ticks, Crickets, Clover Mites and other
listed pests.

Covers up to 5,000 Sq. Ft.

Aids in the Control of Imported Fire Ants

**SATISFACTION
GUARANTEED**

**FASTER ACTING...
CONTAINS MICRO-CEL® E**



Covers up to 5,000 Sq. Ft.

Keep out of reach of children

CAUTION
See additional precautionary
statements on back of bag

NET WT. 10 LBS.

**ORTHO Diazinon
Soil & Turf Insect Control
02591A**

MONEY BACK GUARANTEE
If you are not completely satisfied with this product, return the unused portion to your dealer for a full refund.

Diazinon
Soil & Turf
Insect Control
02591A

DESIGN

76h

ORTHO Diazinon Soil & Turf Insect Control 02591A

TO OPEN
CUT SPOUT HERE



ORTHO Diazinon Soil & Turf Insect Control

CAUTION
Keep out of reach of children.
Avoid contact with skin and clothing.
Do not inhale dust or spray.
Do not eat, drink, or smoke while using this product.
Wash hands thoroughly after use.
Do not use in areas where children play or where animals graze.
Do not use on lawns or in areas where people walk.
Do not use on lawns or in areas where people walk.

Target Pest	Application Rate
Ants	1/2 lb. per 1,000 sq. ft.
Beetles	1/2 lb. per 1,000 sq. ft.
Billbugs	1/2 lb. per 1,000 sq. ft.
Chinch Bugs	1/2 lb. per 1,000 sq. ft.
Crickets	1/2 lb. per 1,000 sq. ft.
Cutworms	1/2 lb. per 1,000 sq. ft.
Fleas	1/2 lb. per 1,000 sq. ft.
Grubs	1/2 lb. per 1,000 sq. ft.
Ground Squirrels	1/2 lb. per 1,000 sq. ft.
House Flies	1/2 lb. per 1,000 sq. ft.
Leaf Beetles	1/2 lb. per 1,000 sq. ft.
Measurers	1/2 lb. per 1,000 sq. ft.
Moles	1/2 lb. per 1,000 sq. ft.
Root Maggots	1/2 lb. per 1,000 sq. ft.
Scavenger Beetles	1/2 lb. per 1,000 sq. ft.
Spiders	1/2 lb. per 1,000 sq. ft.
Stink Bugs	1/2 lb. per 1,000 sq. ft.
Termites	1/2 lb. per 1,000 sq. ft.
Worms	1/2 lb. per 1,000 sq. ft.
Yellowjackets	1/2 lb. per 1,000 sq. ft.

STORAGE AND DISPOSAL
Store in original container in a cool, dry place.
Do not store in areas where children play or where animals graze.
Do not store in areas where people walk.
Do not store in areas where people walk.

Chemical Company
1000 North 10th Street
P.O. Box 1000
St. Paul, MN 55101
Tel: 612-291-1000



BEST AVAILABLE COPY



ORTHO


ACCEPTED
JAN 13 1993
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 239-247

Diazinon Soil & Turf Insect Control

This product must not be used on golf courses and sod farms.
Ready-to-use Granules For Use When Seeding or Transplanting Vegetables
Kills: Wireworms, Rootworms, Root Maggots, Cutworms and Other Listed Soil Insects
Contains Diazinon
Ready-to-use granules for use only on:
• Home Lawns to Kill Billbugs, Chinch Bugs, Sod Webworms and White Grubs, as listed.
• Home Vegetable Gardens to Kill Cutworms, Wireworms, Root Maggots on listed crops.
• Nuisance Pests Outside the Home to Kill Ants, Fleas, Brown Dog Ticks, Crickets, Clover Mites and other listed pests.
Covers up to 5,000 Sq. Ft.
Aids in the Control of Imported Fire Ants

SATISFACTION GUARANTEED

**FASTER ACTING
CONTAINS MICRO-CEL® E**



Covers up to 5,000 Sq. Ft.

CAUTION
Keep out of reach of children
See additional precautionary statements on back of bag

NET WT. 10 LBS.

ORTHO Diazinon Soil & Turf Insect Control 02591A

Diazinon Soil & Turf Insect Control
Covers up to 5,000 Sq. Ft.

MONEY-BACK GUARANTEE
If you are not completely satisfied with this product, we will refund your money.

9/6
5

BEST AVAILABLE COPY



ORTHO Diazinon Soil & Turf Insect Control

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye injury. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Do not store near food or feed products. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides. Do not contaminate ornamental fish ponds. Do not allow children or pets on treated areas until granules have been watered into the soil and the grass or soil is dry.

STATEMENT OF PRACTICAL TREATMENT

If swallowed, call a physician immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until the fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If inhaled, remove victim to fresh air and apply artificial respiration if indicated.

If on skin, wash promptly with plenty of soap and water. Rinse thoroughly. Get medical attention.

If in eyes, rinse eyes with plenty of water and call a physician immediately.

Note to Physicians: Emergency information: call 1-800-457-2022. This is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds, fish and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York between November 1 and May 20. Do not exceed maximum permitted application rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping granules. Where irrigation waterings is recommended in the directions or if waterfowl (ducks or geese), can be expected in the treated area after treatment, irrigate spot treatment, apply a minimum of 0.25 inch of water immediately after applying this product. Stop irrigation before puddling occurs. Keep out of lakes, streams, ponds, farm ditches and estuaries. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS

DIRECTIONS

Overall Lawn Application

ORTHO Diazinon Soil & Turf Insect Control controls the following lawn insects on both grass and dichondra lawns:

- Ants - Armyworms - Bermuda Mites - Brown Dog Ticks - Chiggers - Chinch Bugs - Clover Mites - Crickets - Cutworms - Earwigs - European Crane Fly (Larvae) - Fleas - Lawn Moths (Sod Webworms) - Leafhoppers - May Beetle Grubs - Mole Crickets - Millipedes - Sowbugs - Springtails (Collembola)

Apply ORTHO Diazinon Soil & Turf Insect Control with a fertilizer spreader. See spreader setting chart. Apply when grass is dry. Water lawn lightly after application. Repeat as necessary.

For Mole Crickets, apply in May or June when pests are small. Water lawn thoroughly after treatment.

For application with other tools, see amount you can treat with 1 lb. of product. For lawns less than 20' by 50' or 3'2" or per 100 sq. ft. in area 10' by 10'. For heavier lawns such as St. Augustine or for the heavier to medium insects such as Billbugs, Chinch Bugs, European Crane Fly, Mole Crickets and White Grubs, repeat application in 10 to 14 days if as necessary.

Billbugs, White Grubs of Japanese Beetle, European Chafer and Southern Chafer

Apply 2 lbs. per 1000 sq. ft. of lawn. Water grass thoroughly after application. For Billbug control, apply when ants are first observed or when chewed or brown grass indicates damage from these insects. Repeat application in 10 to 14 days or as necessary. For Chinch Bugs, apply any time between late July and early October.

European Crane Fly (Larvae) (Except California)

Apply 2 lbs. per 1000 sq. ft. of lawn. Water lawn thoroughly after application. Repeat application

SPREADER SETTING NUMBER

Spreader Make and Model	Application Rate of Product 2 lbs per 1000 sq. ft.
Central Imperial Mark No 1662-G	6 1/2
Central Suburban Lawn Spreader No 1622-0	6 1/2
Cyclone Model B1 Rotary	4 1/2
Greenfield Model 105	6 1/2
ORTHO Drop Spreader (Models 3000, 5000)	6
ORTHO Broadcast Spreader (Models 4000, 6000)	5
J.C. Penney Model 6005	7 1/2
J.C. Penney Rotary Spreader	5
Scott Precision Flow Spreader Model PF 4	4
Sears Craftsman Model 6719190	6

A Spot Treatment

Ants: Apply 1 1/2 teaspoons of ORTHO Diazinon Soil & Turf Insect Control over and around each ant hill.

B Imported Fire Ants

To aid in the control of Imported Fire Ants in lawns and other outdoor home recreational areas, apply 1/2 cup in a 2 ft. radius over and around each fire ant mound immediately after application. Water in thoroughly. A minimum of 1 gallon of water should be applied to each mound area. Do not use more than 2 gallons of water. Apply water gently to avoid disturbing ants. Use of high pressure watering may disturb the ants and cause migration, thus reducing effectiveness of treatment. Treat new mounds as they appear. For best results, apply in cool weather, 65 to 80°F or in early morning or late evening hours.

C Bigger Wasps

Apply 1 1/2 teaspoons of ORTHO Diazinon Soil & Turf Insect Control in and around each nest opening. Apply at dusk when wasps are not active.

D Bead Treatment Around House Foundation

The following pests are commonly found around or near foundations of houses as well as in lawns:

Cutworms (surface and subterranean)

Broccoli	Endive	*Pole Beans
Brussels	(Escarole)	Potatoes
Sprouts	Honeydews	Red Beets
Cabbage	Kale	*Snap Beans
Cantaloupes	Lettuce	Spinach
Carrots	*Lima Beans	Summer Squash
Casabas	Muskmelons	Sweet Corn
Cauliflower	Parsley	Swiss Chard
Celery	*Peas	Tomatoes
Collards	Peppers	Watermelons
Crenshaws	Persian	Winter Squash
Cucumbers	Melons	

Apply 10 oz. per 500 sq. ft. and mix into soil to a depth of 2 to 3 inches for surface Cutworms and 3 to 6 inches for subterranean Cutworms. For Potatoes, mix into soil 4 to 6 inches deep.

*NOTE: Fresh only. DO NOT use on dried bean or pea varieties.

Mole Crickets

Broccoli	Collards	Peppers
Brussels	Endive	Radishes
Sprouts	(Escarole)	Red Beets
Cabbage	Kale	Strawberries
Carrots	Lettuce	Tomatoes
Cauliflower	Mustard	Turnips

Apply 3 1/2 oz. per 500 sq. ft. and mix into soil to a depth of 1 to 2 inches.

Wireworms

Broccoli	Endive	Potatoes
Brussels	(Escarole)	Red Beets
Sprouts	Honeydews	*Snap Beans
Cabbage	Kale	Spinach
Cantaloupes	Lettuce	Summer Squash
Carrots	*Lima Beans	Sweet Corn
Casabas	Muskmelons	Sweet Potatoes
Cauliflower	Onions	Swiss Chard
Celery	Parsley	Tomatoes
Collards	*Peas	Turnips
Corn	Peppers	Watermelons
Crenshaws	Persian Melons	Winter Squash
Cucumbers	*Pole Beans	

Apply 10 1/2 oz. per 500 sq. ft. and mix into soil to a depth of 4 to 6 inches.

*NOTE: Fresh only. DO NOT use on dried bean or pea varieties.

Broccoli, Brussels, Sprouts, Cabbage, Cauli-

MONEY BACK GUARANTEE

If you are not satisfied after using this product, return this guarantee with your sales receipt and reason for dissatisfaction to:

Chevron Chemical Co., Ortho Consumer Affairs Dept., P.O. Box 5047, San Ramon, CA 94583-0947

ORTHO Diazinon Soil & Turf Insect Control 02591A

969

Ortho Diazinon kills over 100 insects on both grass and dicotyledonous lawns

Ants - Armyworms - Bermuda Mites - Brown Dog Ticks - Chiggers - Chinch Bugs - Clover Mites - Crickets - Cutworms - Earwigs - European Crane Fly (Larvae) - Fleas - Lawn Moths (Sod Webworms) - Leafhoppers - May Beetle Grubs - Mole Crickets - Millipedes - Sowbugs - Springtails (Collembola)

Apply ORTHO Diazinon Soil & Turf Insect Control with a fertilizer spreader. See spreader setting chart. Apply when grass is dry. **Water lawn lightly after application.** Repeat as necessary.

For **Mole Crickets**, apply in May or June when pests are small. **Water lawn thoroughly after treatment.**

For application with other types of equipment, apply at the rate of 2 lbs. per 1000 sq. ft. (an area 20' by 50' or 32' oz. per 100 sq. ft. (an area 10' by 10') for heavier lawns such as St. Augustine or for the harder to control insects such as Billbugs, Chinch Bugs, European Crane Fly, Mole Crickets, and White Grubs repeat application in 10 to 14 days or as necessary.

Billbugs, White Grubs of Japanese Beetle, European Chaler, and Southern Chaler

Apply 2 lbs. per 1000 sq. ft. of lawn. **Water grass thoroughly after application.** For Billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat application in 10 to 14 days or as necessary. For Grub control, apply any time between late July and early October.

European Crane Fly (Larvae) (Except California)

Apply 2 lbs. per 1000 sq. ft. of lawn. **Water lawn thoroughly after application.** Repeat application in 10 to 14 days or as necessary. For timing of application in your area, consult the State Agricultural Extension Service.

May Beetle Grubs (Except California)

Apply 2 lbs. per 1000 sq. ft. of lawn between mid April or early May or during August when activity is first observed or when brown grass indicates damage from this pest. **Water lawn thoroughly after application.** Repeat application in 10 to 14 days or as necessary.

Hyperodes Weevils (New York only)

Apply 2 lbs. per 1000 sq. ft. of grass. **Water grass thoroughly after application.** Treat problem areas in mid April and again in mid May.

NOTE
The spreader settings listed are approximate. Accurately weigh a few pounds into your spreader and apply to measured area. Then adjust spreader setting if necessary in order to assure correct rate of application.

STORAGE AND DISPOSAL

STORAGE Keep pesticide in original container. Do not put into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area.

DISPOSAL PRODUCT - Partially filled bag may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash. **CONTAINER** - Do not reuse empty bag. Discard bag in trash.

NOTICE Buyer assumes all responsibility for safety and use not in accordance with directions.

Manufactured for
Chevron Chemical Company © 1985
Ortho Consumer Products Division
P.O. Box 5047 San Ramon CA 94583-0947
Form R10311-J Product 5403
EPA Reg. No. 239-2479-AA



EPA Est. 239-MO-1, 44616-MO-01
Superscript corresponds to first letter of lot number on bottom of container

Ortho Model 105	6
ORTHODrop Spreader (Models 3000, 5000)	6
ORTHOBroadcast Spreader (Models 4000, 6000)	5
J.C. Penney Model 6005	7
J.C. Penney Rotary Spreader	5
Scott Precision Flow Spreader Model PF 4	4
Sears Craftsman Model 67119198	6

Spot Treatment
Ants Apply 1 1/2 teaspoons of ORTHO Diazinon Soil & Turf Insect Control over and around each ant hill.

Imported Fire Ants To aid in the control of Imported Fire Ants in lawns and other outdoor home recreational areas, apply 1 cup in a 2 ft. radius over and around each fire ant mound immediately after application. Water in thoroughly. A minimum of 1 gallon of water should be applied to each mound area. Do not use more than 2 gallons of water. Apply water gently to avoid disturbing ants. Use of high pressure watering may disturb the ants and cause migration thus reducing effectiveness of treatment. Treat new mounds as they appear. For best results, apply in cool weather 65 to 80°F or in early morning or late evening hours.

Digger Wasps Apply 1 1/2 teaspoons of ORTHO Diazinon Soil & Turf Insect Control in and around each nest opening. Apply at dusk when wasps are not active.

Band Treatment Around House Foundation The following pests are commonly found around or near foundations of houses as well as in lawns. Certain of these pests may enter houses:

- Ants
- Fleas
- Brown Dog Ticks
- Millipedes
- Clover Mites
- Sowbugs (Pillbugs)
- Crickets
- Springtails
- Earwigs
- Collembola

Apply a 5 foot band of ORTHO Diazinon Soil & Turf Insect Control next to the house foundation. Distribute granules uniformly, using a rate of 3/2 oz. for each 100 sq. ft. of area 15' by 20'. Repeat as necessary.

Garden Insect Control
ORTHODiazinon Soil & Turf Insect Control controls certain garden insects on crops listed. Apply just prior to planting and immediately mix thoroughly into soil to the specified depth.

NOTE One level measuring cup (8 fl. oz.) holds 4 oz. of granules.

Peppers	Sweet Chard
Collards	Tomatoes
Crenshaws	Watermelons
Cucumbers	Winter Squash

Apply 10 oz. per 500 sq. ft. and mix into soil to a depth of 2 to 3 inches for surface Cutworms and 3 to 6 inches for subterranean Cutworms. For Potatoes, mix into soil 4 to 6 inches deep.

***NOTE** Fresh only. DO NOT use on dried bean or pea varieties.

Broccoli	Collards	Peppers
Brussels	Endive	Radishes
Sprouts	(Escarole)	Red Beets
Cabbage	Kale	Strawberries
Carrots	Lettuce	Tomatoes
Cauliflower	Mustard	Turnips

Apply 3 1/2 oz. per 500 sq. ft. and mix into soil to a depth of 1 to 2 inches.

Broccoli	Endive	Potatoes
Brussels	(Escarole)	Red Beets
Sprouts	Honeydews	*Snap Beans
Cabbage	Kale	Spinach
Cantaloupes	Lettuce	Summer Squash
Carrots	*Lima Beans	Sweet Corn
Casabas	Muskmelons	Sweet Potatoes
Cauliflower	Onions	Swiss Chard
Cherry	Parsley	Tomatoes
Collards	*Peas	Turnips
Corn	Peppers	Watermelons
Crenshaws	Persian Melons	Winter Squash
Cucumbers	*Pole Beans	

Apply 10 1/2 oz. per 500 sq. ft. and mix into soil to a depth of 4 to 8 inches.

***NOTE** Fresh only. DO NOT use on dried bean or pea varieties.

Broccoli, Brussels Sprouts, Cabbage, Cauliflower

Root Maggots Apply 10 oz. per 500 sq. ft. over an tire seedbed before planting or transplanting and mix into soil to a depth of 3 to 4 inches.

Onions
Onion Maggots Apply 7 oz. per 500 sq. ft. and mix into soil to a depth of 3 to 4 inches.

MICRO CEL® Reg. TM of Celite Holdings Corp.



By Wt 95%

Active Ingredient
Diazinon [O,O diethyl O (2 isopropyl 6 methyl 4 pyrimidinyl)phosphorothioate]
Inert Ingredients



A