

Reg # 239 - 2479

PM-13

135

ORTHO Diazinon Soil & Turf
Insect Control
EPA Reg. No. 239-2479

COPY A

- POWERFLO™ Granules
- ~~Improved Formula~~
- ~~Better Spreading Formula~~
- Less Dust & Drift
- Denser Granule Stays in Place
- For Uniform Coverage

WLC
3/10/93

TO OPEN
CUT SPOUT HERE

1 INODAC FORMULATION ONLY
2 CORNED FORMULATION ONLY

Imazinon Soil & Turf Insect Control



ORTHO

Diazinon Soil & Turf Insect Control

SECRET

PROVEY BACK GUARANTEE

If you are not satisfied after using this product, I will return your money. No questions asked. I will also provide you with a full refund of your money. I will also provide you with a full refund of your money. I will also provide you with a full refund of your money.

Provey Chemical Co. One Customer Affairs Dept.
P.O. Box 1001 San Rafael, CA 94501

ACCEPTED

239-2479

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No.

Diazinon
Ortho Soil & Turf Insect Control
Covers up to 5,000 Sq. Ft.
02591A

This product must not be used on golf courses and sod farms.

Ready-to-use Granules For Use When Seeding or Transplanting Vegetables

**Kills: Wireworms, Rootworms, Root Maggots, Cutworms
and Other Listed Soil Insects
Contains Diazinon**

Ready-to-use granules for use only on:

- Home Lawns to Kill Billbugs, Chinch Bugs, Sod Webworms and White Grubs - as listed.

- **Have Vegetable Gardens to Kill Cancer.**

Wireworms, Root Maggots on listed crops.


- **Nuisance Pests Outside the Home to Kill Ants, Fleas, Brown Dog Ticks, Crickets, Clever Mites and other**

listed posts. **S, SPD**

Covers up to 500 Sq. Ft.

Aids in the Control of Imported Fire Arms

**FASTER ACTING ...
CONTAINS MICRO-CEL®**



NET WT. 10 LBS.

ORTHO Diazinon
Soil & Turf Insect Control
02591A

Diazinon
Ortho Soil & Turf Insect Control
Covers up to 5,000 Sq. Ft.

BEST AVAILABLE COPY

MASTER LABEL—INCLUDES TEXT FOR VARIOUS FORMULATIONS

1 BIODAC FORMULATION ONLY
2 CORNCOB FORMULATION ONLY



ORTHO Diazinon Soil & Turf Insect Control

BEST AVAILABLE COPY

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

May irritate skin, eyes, or clothing. Causes moderate eye injury. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after use. Do not eat, drink, or smoke near food or feed products. Food utensils such as spoons or measuring cups must not be used for food or feed. Do not contaminate ornamental fish ponds. Do not allow children or pets on treated areas until granules have been watered into the soil at the grass line.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Do not induce vomiting. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit is clear. Do not eat or drink anything by mouth if person is unconscious or convulsing.

If inhaled: Remove from area. Apply artificial respiration if indicated.

If on skin: Wash thoroughly with soap and water. Rinse thoroughly. Get medical attention.

If in eyes: Wash thoroughly with water and a clean, moist cloth.

Note to Physicians: Emergency information: 1-800-457-2022. This is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate, 1 mg. per 100 lbs. body weight, may be administered, but only in conjunction with atropine.

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds and other wildlife. Birds, especially waterfowl feeding or drinking on treated areas may be killed. Because of the migratory nature of waterfowl, do not apply this product to lawns in Nassau County, New York between November 1 and May 20. Do not exceed maximum application rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping applications. After application, watering is recommended in the directions of flow. Ducks or geese can be expected in the treated area after treatment. Avoid application of 0.25 inch of water immediately after applying this product. Stop irrigation before puddling occurs. Keep out of lakes, streams, ponds, marshes, and estuaries. Do not apply directly to water to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

DIRECTIONS FOR USE

Read entire label carefully before using this product. Do not use if label is missing or torn.

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS

DIRECTIONS

Overall Lawn Application

ORTHO Diazinon Soil & Turf Insect Control controls the following lawn insects on both grass and mixed lawns:

Ants - Armyworms - Bermuda Mites - Brown Dog Ticks - Chiggers - Chinch Bugs - Clover Mites - Crickets - Cutworms - Earwigs - European Crane Fly - Fleas - Fleas - Lawn Moths - Sod Webworms - Leafhoppers - May Beetle Grubs - Mealybugs - Millipedes - Sowbugs - Springtails - Centipedes

Apply ORTHO Diazinon Soil & Turf Insect Control to the lawn in early evening. Water lawn thoroughly after application. Repeat application as necessary.

For Mole Crickets: Apply ORTHO Diazinon Soil & Turf Insect Control to the lawn in early evening. Water lawn thoroughly after treatment.

For Billbugs, White Grubs of Japanese Beetle, European Chafers and Southern Chafers: Apply ORTHO Diazinon Soil & Turf Insect Control to the lawn in early evening. Water lawn thoroughly after application. Repeat application as necessary.

For Sod Webworms: Apply ORTHO Diazinon Soil & Turf Insect Control to the lawn in early evening. Water lawn thoroughly after application. Repeat application as necessary.

PRECAUTIONARY STATEMENTS

(CORNCOB FORMULATION ONLY)

Spot Treatment

Ants: Apply 1/2 teaspoon of ORTHO Diazinon Soil & Turf Insect Control over and around each ant hill.

Imported Fire Ants: To aid in the control of imported fire ants in lawns and other outdoor home recreational areas, apply 1/2 cup in a 2 ft. radius over and around each fire ant mound. Immediately after application, water in thoroughly. A minimum of 1/2 gallon of water should be applied to each mound area. Do not use more than 2 gallons of water. Apply water gently to avoid disturbing ants. Use of high pressure watering may disturb the ants and cause migration, thus reducing effectiveness of treatment. Treat new mounds as they appear. For best results, apply on cool weather (65 to 80°F) in early morning or late evening hours.

Digger Wasps: Apply 1/2 teaspoon of ORTHO Diazinon Soil & Turf Insect Control in and around each nest opening. Apply at dusk when wasps are not active.

(BIODAC FORMULATION ONLY)

Spreaders, Make and Model

ORTHO Broadcast Spreaders: 1/2 lb. per 1000 sq. ft.

ORTHO Drop Spreaders: 1/4 lb. per 1000 sq. ft.

Craftsman (Model 6711000): 1/2 lb. per 1000 sq. ft.

Scotts Precision Flow (Model PF-1): 1/4 lb. per 1000 sq. ft.

Scotts Easy Green: 1/2 lb. per 1000 sq. ft.

Cycle-Retrol (Model 96-5): 1/2 lb. per 1000 sq. ft.

True Runner (Model CB-1100): 1/2 lb. per 1000 sq. ft.

Garden Insect Control

ORTHO Diazinon Soil & Turf Insect Control controls certain garden insects on crops listed. Apply just prior to planting and immediately mix thoroughly into soil to the specified depth.

(CORNCOB FORMULATION ONLY)

NOTE: One level measuring cup (8 fl. oz.) holds 1/2 lb. of granules.

(BIODAC FORMULATION ONLY)

NOTE: One level measuring cup (8 fl. oz.) holds 1/2 lb. of granules.

Cutworms (surface and subterranean)

Broccoli	Endive	*Pea Beans
Brussels	Escarole	Potatoes
Cauliflower	Horseradish	Radishes
Cabbage	Kale	*Snap Beans
Cantaloupes	Lettuce	Squash
Carrots	* Lima Beans	Sweet Corn
Celery	Mustard Greens	Tomatoes
Collards	Peas	Winter Squash
Cress	Peppers	
Cucumbers	Parsnips	
	Spinach	

Apply 10 oz. per 500 sq. ft. and mix into soil to a depth of 2 to 3 inches for surface cutworms and 3 to 6 inches for subterranean cutworms. For Potatoes, mix into soil 4 to 6 inches deep.

*NOTE: Fresh only. DO NOT use on dried bean or pea varieties.

Mole Crickets

Broccoli	Collards	Peppers
Brussels	Endive	Radishes
Cauliflower	Escarole	Radishes
Cabbage	Kale	Strawberries
Carrots	Lettuce	Tomatoes
Cauliflower	Mustard	Turnips

ACCEPTED

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No.

MONEY BACK GUARANTEE

If you are not satisfied after using this product, return this guarantee with your sales receipt and reason for dissatisfaction to:

Chevron Chemical Co., Ortho Consumer Affairs Dept., P.O. Box 5047, San Ramon, CA 94583 0947

ORTHO Diazinon Soil & Turf Insect Control 02591A

By Wt
isopropyl 6-methyl
cyclohexanecarboxylate

5%
95%

