



BRUSH-B-GON™ ORTHO Brush Killer

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid contact with eyes. Avoid breathing spray mist. Wash thoroughly after handling. **STATEMENT OF PRACTICAL TREATMENT:** In case of eye contact, immediately flush eyes with fresh water for 15 minutes. If irritation continues, see a doctor. If swallowed, induce vomiting by administration of syrup of ipecac or under medical supervision. Call a physician. **Note to Physicians:** Emergency Information: call (415) 233-3737.

ENVIRONMENTAL HAZARDS: Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

BRUSH-B-GON Brush Killer is formulated to control unwanted woody vegetation in areas around homes, cabins, buildings, fence lines, trails, rights-of-way and other non-crop areas.

CONTROLS

Arkansas Rose
Big Leaf Maple
Black Cottonwood
Black Willow
Box Elder
California Rose
Choke Cherry
Common Ragweed
Elderberry
Elm
Evergreen Blackberry
Giant Ragweed

Himalayan Blackberry
Honeysuckle
Kudzu
Mackenzie Willow
Mesquite
Mimosa
Maple
Narrow Leaved Willow
Oak
Persimmon
Poison Ivy
Poison Oak

Raspberry
Red Alder
Sassafras
Scouler Willow
Sweet Gum
Trumpet Blackberry
Trumpet Creeper
Virginia Creeper
Wild Blackberry
Wild Grape
Wild Rose
Willow

WHEN TO APPLY:

Apply as a full coverage foliar spray when vegetation is fully leafed out and actively growing. Apply also to trunks and limbs of trees and stems of vines. Wilting or other symptoms will occur in 2 to 6 weeks. Application to mature foliage is preferred. Apply only when the air is calm to prevent drift to desirable vegetation. Rainfall within 24 hours may reduce effectiveness of treatment. Do not use on or around edible feed or food crops.

HOW TO APPLY FOLIAGE SPRAY:

To apply with ORTHO hose-end sprayers (ORTHO SPRAY ETTE 4, ORTHO Super Six SPRAY-ETTE or ORTHO Lawn and Garden Sprayer on "FOLIAGE" setting), tank sprayer or power sprayer, mix 4 fl. oz. per gal. water and apply to thoroughly wet all leaves, stems and trunks. Use low pressure and coarse spray pattern to reduce spray drift which may possibly damage desirable vegetation. Certain hard-to-control species, such as Poison Oak, Poison Ivy, Blackberries and Honeysuckle may require retreatment the following season for complete control.

IMPORTANT:

BRUSH-B-GON Brush Killer is non-selective. Do not apply to or clean equipment around desirable vegetation or ornamental plants as injury may result. Do not apply when drift may occur.

STORAGE AND DISPOSAL

Avoid contamination of food, feed or water. Do not reuse empty container. Wrap container and put in trash collection.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Chevron Chemical Company

Ortho Consumer Products Division
San Francisco CA 94120 7144 Richmond CA 94804 0036
Form 10125-C Product 5467 Made in U.S.A.
EPA Reg. No. 239 2491 AA

EPA Est. 239-CA-1, 239-NJ-1, 239-MG-1
Superscript used is first letter of lot number



Wash thoroughly
Immediately flush eyes
...containing by ad-
...system. Emer

...minate water

...consistent with

...STATEMENTS

...round homes.

...berry
...d Alder
...Kudzu
...Willow
...Gum
...Blackberry
...Creeper
...Creeper
...Blackberry
...Grape
...Rose
...

WHEN TO APPLY

Apply as a full coverage foliar spray when vegetation is fully leafed out and actively growing. Apply also to trunks and limbs of trees and stems of vines. Withering or other symptoms will occur in 2 to 6 weeks. Application to mature foliage is preferred. Apply only when the air is calm to prevent drift to desirable vegetation. Rainfall within 24 hours may reduce effectiveness of treatment. Do not use on or around edible feed or food crops.

HOW TO APPLY FOLIAGE SPRAY

To apply with ORTHO hose end sprayers (ORTHOSpray ETTIE 4 ORTHO Super Six Spray ETTIE or ORTHO Lawn and Garden Sprayer on "FOLIAGE" setting), tank sprayer or power sprayer, mix 4 fl. oz. per gal. water and apply to thoroughly wet all leaves, stems and trunks. Use low pressure and coarse spray pattern to reduce spray drift which may possibly damage desirable vegetation. Certain hard-to-control species, such as Poison Oak, Poison Ivy, Blackberries and Honeysuckle may require retreatment the following season for complete control.

IMPORTANT:

BRUSH-B-GON Brush Killer is non-selective. Do not apply to or clean equipment around desirable vegetation or ornamental plants as injury may result. Do not apply when drift may occur.

STORAGE AND DISPOSAL

Avoid contamination of food, feed or water. Do not reuse empty container. Wrap container and put in trash collection.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Chevron Chemical Company

Ortho Consumer Products Division
San Francisco CA 94120-7144 Richmond CA 94604-0036
Form 10125 C Product 5467 Made in U.S.A.
EPA Reg. No. 239-2491-AA

EPA Est. 239-CA-1 239-NJ-1 239-MO-1
Superscript used is first letter of lot number



ORTHO

Makes
16 Gals
Spray

**BRUSH-B-GON™
Brush Killer**

KILLS WOODY PLANTS



**POISON OAK, POISON IVY,
WILLOWS, OAK, KUDZU,
BLACKBERRIES, & OTHER
LISTED WOODY PLANTS**

BRUSH-B-GON Brush Killer
239-2491

COPY A

STUMP KILLING: To kill stumps and prevent regrowth from stumps of trees such as Alder, Box Elder, Black Cherry, Choke Cherry, Elm, Manzanita, Maple, Oak, Persimmon, Peach and Willow, apply product undiluted with a paint brush to the entire freshly cut stump, including the cut surface, remaining trunk and any exposed root surfaces. Do not reuse brush. Wrap brush in newspaper and put in trash collection.

COPY B

VINE KILLING: When brushy vines such as poison oak, poison ivy, etc. grows on or around desirable plants, do not allow spray to contact desirable vegetation. Cut the vine first and treat as directed for stump killing.

