



ACCEPTED

DEC 9 1983

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended for the pesticide registered under EPA Reg. No. 235-2496



ORTHO

# DIFOLATAN® 80

## Water Dispersible Granules

A WATER DISPERSIBLE GRANULAR FUNGICIDE FOR PLANT DISEASE CONTROL

Active Ingredient  
Captafol  
Inert Ingredients  
cis-N-[(1,1,2,2-tetrachloroethyl)thio]-4-cyclohexene-1,2-dicarboximide

By Wt.  
80%  
20%

KEEP OUT OF REACH OF CHILDREN  
DANGER-PELIGRO  
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

Chevron Chemical Company

Richmond, CA 94804-0035

NET WEIGHT 25 POUNDS

In the dormant period, or beginning when first petals fall of the 25 lb. rate of DIFOLATAN 80 Water Dispersible Granules was applied in the dormant period. Do not grow trees following DIFOLATAN 80 applications.

**BLUEBERRIES:** For Control of Fusicoccum Canker, Mummyberry and Phomopsis Twig Blight, Anthracnose Fruit Rot — Apply 2 1/2 to 5 lbs. (2 to 4 lbs. active) in 20 to 250 gals. spray mix per acre. Thorough coverage is essential. Do not apply within 21 days of harvest.

**Fusicoccum Canker:** Begin application at first outbreak and repeat at 4 to 6 week intervals. For the application immediately after harvest and before through leaf drop.

**Mummyberry and Phomopsis Twig Blight:** Begin application at early green tip of fruit and repeat at 10 to 14 day intervals until petal fall.

**Anthracnose Fruit Rot:** Begin application at first petal fall and repeat at 10 to 14 day intervals.

**SOUR CHERRIES:** Montmorency Variety Only — Mechanically Harvested Only. Monilinia Brown Rot Blossom Blight — Apply 1 to 100 gals. spray mix per acre. Begin application at first petal fall and repeat at 10 to 14 day intervals until petal fall.

**Coccomyces Leaf Spot:** a. Seasonal Program — Apply 1 to 4 lbs. per 100 gals. spray mix per acre. b. Reduced Application Program — Apply 1 to 4 lbs. per 100 gals. spray mix per acre.

**Monilinia Fruit Brown Rot:** Apply 1 to 100 gals. spray mix per acre. Begin application at first petal fall and repeat at 10 to 14 day intervals until petal fall.

**CRANBERRIES:** Fruit Rot — Apply 1 to 100 gals. spray mix per acre. Begin application at first petal fall and repeat at 10 to 14 day intervals until petal fall.

**CUCUMBERS:** Downy Mildew, Scab, Gummy Stem Blight, Anthracnose, Alternaria Leaf Blight — Apply 1 to 100 gals. spray mix per acre.

**MACADAMIA NUTS:** Hawaii Race of Blossom Blight (Botrytis) — Apply 1 to 100 gals. spray mix per acre.

**MELONS:** Cantaloupe, Honey Dew, Muskmelon and Others: Alternaria Leaf Blight, Downy Mildew, Anthracnose, Gummy Stem Blight — Apply 1 to 100 gals. spray mix per acre.

**ONIONS:** Purple Blotch — Apply 1 to 100 gals. spray mix per acre.

**ORANGES, GRAPEFRUIT, TANGERINES, LEMONS AND LIMES IN FLORIDA ONLY:** Ground Application — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Downy Mildew, Scab, Gummy Stem Blight, Anthracnose, Alternaria Leaf Blight — Apply 1 to 100 gals. spray mix per acre.

**PEARS:** Downy Mildew, Scab, Gummy Stem Blight, Anthracnose, Alternaria Leaf Blight — Apply 1 to 100 gals. spray mix per acre.

**PEPPERS:** Downy Mildew, Scab, Gummy Stem Blight, Anthracnose, Alternaria Leaf Blight — Apply 1 to 100 gals. spray mix per acre.

**POTATOES:** Early Blight, Late Blight — Apply 1 to 2 lbs. per acre in water to cover. Begin treatment when plants are 6" high and continue at 7 to 10 day intervals.

**PRUNES:** Brown Rot Blossom Blight — (California and Oregon only) — 1 to 2 lbs. per 100 gals. spray mix per acre, depending upon tree size.

**SWEET CORN:** (Florida Only) For Fresh Vegetable Use only. Do Not Apply to Sweet Corn to be Processed — Northern and Southern Corn Leaf Blight — Apply 1 to 100 gals. spray mix per acre.

**TOMATOES:** (Mechanically Harvested Only): Anthracnose, Early Blight, Late Blight, Gray Leaf Spot (Stemphylium spp.), Nail Head Spot, Septoria Leaf Spot, Fruit Rot — Apply 1 to 100 gals. spray mix per acre.

**WATERMELONS:** Gummy Stem Blight, Anthracnose, Downy Mildew — Apply 1 to 100 gals. spray mix per acre.

**WEEDLAND TARO (HAWAII ONLY):** Phytophthora Leaf Blight — 1 to 2 lbs. per 100 gals. spray mix per acre.

**TEMPLE ORANGES:** Florida Only — Scab and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

when mature fruit are on the trees. DIFOLATAN 80 Water Dispersible Granules may cause minor leaf spotting on new flush growth. DIFOLATAN 80 Water Dispersible Granules may cause minor spotting when applied to young citrus fruit. Do not apply to young citrus fruit grown for the fresh market.

**PEANUTS:** For Control of Early (Cercospora) Leafspot in all U.S. Peanut Growing Areas, Suppression of Web Blotch in Texas and Oklahoma and Control of Rust in Florida — Apply 2 lbs. in 10 to 60 gals. of spray mix per acre with ground equipment or in 5 to 10 gals. with aerial equipment. Begin application when disease first appears, usually 25 to 45 days after planting. Repeat at 10 to 14 day intervals to maintain control. Under severe disease conditions, apply at 10 day intervals. Do not apply within 14 days of digging. Do not graze treated areas or feed treated vines or hay to livestock. Do not hogdown treated areas.

**PINEAPPLES:** Pineapple Heart Rot — Dip Treatment — 1 1/2 lbs. per 10 gals. mixture. Dip immediately before planting using 100 to 250 gals. dip per acre. Spray Treatment — 10 lbs. in 400 gals. spray per acre. Apply first application immediately after planting; repeat at monthly intervals for a maximum of 8 applications.

**POTATOES:** Early Blight, Late Blight — 1 to 2 lbs. per acre in water to cover. Begin treatment when plants are 6" high and continue at 7 to 10 day intervals. Under severe disease conditions, use higher listed dosage or shorter spray interval.

**PRUNES:** Brown Rot Blossom Blight — (California and Oregon only) — 1 to 2 lbs. per 100 gals. spray mix per acre, depending upon tree size.

**SWEET CORN:** (Florida Only) For Fresh Vegetable Use only. Do Not Apply to Sweet Corn to be Processed — Northern and Southern Corn Leaf Blight — Apply 1 to 100 gals. spray mix per acre.

**TOMATOES:** (Mechanically Harvested Only): Anthracnose, Early Blight, Late Blight, Gray Leaf Spot (Stemphylium spp.), Nail Head Spot, Septoria Leaf Spot, Fruit Rot — Apply 1 to 100 gals. spray mix per acre.

**WATERMELONS:** Gummy Stem Blight, Anthracnose, Downy Mildew — Apply 1 to 100 gals. spray mix per acre.

**WEEDLAND TARO (HAWAII ONLY):** Phytophthora Leaf Blight — 1 to 2 lbs. per 100 gals. spray mix per acre.

**TEMPLE ORANGES:** Florida Only — Scab and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.

**PEACHES:** Florida Only — Alternaria Brown Spot and Rust Mite — Apply 1 to 100 gals. spray mix per acre.