



ORTHO

Treats up to
4,000 Sq. Ft.

Crab Grass Killer

FORMULA II

Also Kills Dallis Grass and Nutgrass

Active Ingredient	By Wt
Calcium acid methanearsonate	8.4%
Inert Ingredients	91.6%
Total arsenic as elemental	
All water soluble - 4.1%	

Bahia grass

Keep out of reach of children

CAUTION

See side panel for additional precautionary statements.

BEST AVAILABLE COPY

Chevron Chemical Company © 1985

Ortho Consumer Products Division

San Francisco CA 94120-7144

Product 5517

Made in U.S.A.

R Form 10584-B

EPA Reg. No 239-2510

EPA Est. 239-MO-1

NET CONTENTS 1 QT.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Causes eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Do not allow children or animals on treated area until after rainfall or when normal irrigation practice is resumed 2 days after treatment. Wear protective clothing when handling or applying this product, including long pants, long-sleeved shirt and rubber gloves and boots. Do not graze treated areas of feed livestock treated area. **STATEMENT OF PRACTICAL TREATMENT:** In case of eye contact, wash eyes immediately with fresh water for at least 15 minutes. See a doctor if irritation persists. Note to Physicians: Emergency Information - call (415) 233-3737.

ENVIRONMENTAL HAZARDS: Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

DIRECTIONS

ORTHO Crab Grass Killer Formula II can be used to control Crabgrass, Dallis Grass and Nutgrass on all of the major ornamental turf grasses including Dichondra, Bent and Fescue grasses. Do not use on St. Augustine, Carpet and Centipede grasses. Make new dilution for each use.

For best results, apply in the morning (after dew has dried) on a sunny, calm day to allow maximum absorption of weed killer. Late afternoon application may be diluted or washed off leaves by dew or rainfall during the night resulting in unsatisfactory control of weeds. Do not apply when temperatures are expected to exceed 90° F. or rainfall is expected within 24 hours after application.

CRABGRASS (Hairy Crabgrass or Smooth Crabgrass): See Dilution Table for dosages. Soil should be moist before treatment. Do not water for two days after treatment; then resume watering particularly during hot, dry weather. One application may be sufficient if applied when seedlings are less than 2 inches tall. Two applications four to seven days apart are generally needed for control of medium to mature crabgrass plants. Follow-up spot sprays may be necessary for newly germinated crabgrass. Lawns may be reseeded when crabgrass is dead. On newly-seeded lawns, do not commence applications until after two mowings. Bent, Fescue and Dichondra lawns are more sensitive and may be temporarily discolored. Additional wetting agents are not needed.

APPLICATION: Use tank type sprayer. After using sprayer with ORTHO Crab Grass Killer Formula II, clean sprayer thoroughly with water before spraying trees, flowers or vegetables. Do not apply with hose-end sprayer.

DILUTION TABLE

ORTHO Crab Grass Killer Formula II to Use	Add water to make	To cover
3 Tablespoonfuls (1½ oz.)	1 gal.	200 sq. ft.
6 Tablespoonfuls (3 oz.)	2 gals.	400 sq. ft.
8 oz. (1 cup)	5 gals.	1,000 sq. ft.
2 oz. (1 qt.)	20 gals.	4,000 sq. ft.

SPECIAL COMBINATION SPRAY: Use ORTHO Crab Grass Killer Formula II in combination with WEED-B-GON Lawn Weed Killer at recommended rates for both. Use 3 Tablespoonfuls of ORTHO Crab Grass Killer Formula II and 4 teaspoonfuls of WEED-B-GON Lawn Weed Killer per gallon water on 200 sq. ft. area. For larger areas, increase rate proportionately. Avoid drift to ornamentals when using this combination or injury may result. Do not use this combination on Dichondra. Observe all precautions, directions and limitations on both product labels.

DALLIS GRASS in Bermuda Lawns, Bluegrass, Dichondra and Zoysia Lawns: Use 2 ounces (4 Tablespoonfuls) per gallon of water and cover 200 sq. ft. (20' x 10'). Make 2 or 3 applications at weekly intervals starting in late June. Follow cautions given on label for control of crabgrass.

NUTGRASS in Bermuda Lawns and Zoysia Lawns: Use 3 ounces (6 Tablespoonfuls) per gallon of water and cover 200 sq. ft. (20' x 10'). Make 3 to 4 applications at monthly intervals starting in June. Follow cautions given on label for control of crabgrass.

STORAGE AND DISPOSAL

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area. Do not store diluted spray.

DISPOSAL: PRODUCT - Partially filled bottle may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash. **CONTAINER** - Do not reuse empty bottle. Rinse thoroughly before discarding in trash.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

