

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0080, Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

237733

Application for Pesticide - Section I

1. Company/Product Number 239-2522	2. EPA Product Manager Theresa J. Stone	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) ORTHO Daconil 2787 Multipurpose Fungicide	PM# 22	
5. Name and Address of Applicant (Include ZIP Code) The SOLARIS Group of Monsanto Company P. O. Box 5006, San Ramon, CA 94583-0906 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other, Explain _____

NOTIFICATION
JUN 14 1996

Explanation: Use additional page(s) if necessary. (For section I and Section II.)
Submit five copies of finished labeling for APN: ORTHO Daconil Multipurpose Fungicide Plant Disease Control revising the modified Environmental Hazards statement as requested in the Agency's letter of March 28, 1996; also add a modified homeowner exclusionary statement as approved by John Barnin in a 4/15/96 telephone conversation with F. X. Kamienski. This is to certify that this notification meets every requirement of Pr Notice 88-6.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	If "Yes" Unit Packaging wgt. No. per container		<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
If "Yes" Unit Packaging wgt. No. per container		If "Yes" Package wgt No. per container			
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	


Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name F. X. Kamienski	Title Registration Specialist	Telephone No. (Include Area Code) (510) 355-3335
2. Signature <i>Francis X. Kamienski</i>		3. Title Registration Specialist
4. Typed Name Francis X. Kamienski		5. Date May 29, 1996
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received 8/16/96 SIC 8/16/96 REC'D EPA/OPP/DP/DA

ORTHO®

Makes Up To
64 Gallons Diluted Spray

Multi-Purpose Fungicide
Daconil 2787®
Plant Disease Control



CONTROLS DISEASES -
Leaf Spots, Rust, Blights,
Fruit Rots, Mildews,
Scab, Molds, etc.



**ON: Lawns, Vegetables,
Fruit Trees, Flowers,
Shrubs and Shade Trees**

Active Ingredient
Chlorothalonil (tetrachloro-
phthalonitrile) 29.6%
Inert Ingredients 70.4%

KEEP OUT OF REACH OF CHILDREN
WARNING See back panel booklet for
additional precautionary statements.
NET CONTENTS 16 FL OZ (1 PT) 473 mL



NOTIFICATION
JUN 14 1996

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING: Causes eye and skin irritation. This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Do not get in eyes, on skin, or on clothing. Wear long sleeve shirt, long pants, and rubber gloves while mixing and applying this product. Wash skin and hands thoroughly after using. Remove and wash contaminated clothing before reuse.

Keep children and pets off treated area until dry.
FIRST AID: In case of eye contact, immediately flush eyes with fresh water for 15 minutes and see a doctor. In case of skin contact, wash skin with soap and water. Get medical attention if irritation persists. **NOTE:** In Physicians: Emergency information call 1-800-454-2333.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and other aquatic organisms. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply when weather conditions favor drift of spray.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Medical information call 1-800-454-2333
Product information call 1-800-225-2843

ORTHO® SOLARIS®, DIAZINON PLUS®, ISOTOX®,
MALATHION 50 PLUS®, ORTHENE®, SPRAY-ETTE®
Registered trademarks of Monsanto Company
SEVIN® is the Registered trademark of Rhone-Poulenc for
carbamyl insecticide.

DACONIL 2787® trademark of ISK Biotech Corp.
© Monsanto Company 1995
Manufactured for The SOLARIS Group of Monsanto Company
P.O. Box 5008 San Ramon CA 94583-0808
Form 0-730-04 / Product 1000
EPA Reg. No. 239-2522 / EPA Est. 239-1A-31, 58996-MO-1A
Superscript is first letter of lot number
Made in USA

CONIFER DISEASES

PINES, DOUGLAS FIR, SPRUCES -
Scleroderris Canker (Pines), Swiss Needlecast, Rhabdochline and Needlecast (Douglas Fir): Mix 1 fl oz (2 Tbs) / 4 gal water and apply as full coverage spray to the point of runoff.

Rhizosphaera Needlecast (Spruces), Scirrhia Brown Spot (Pine): Mix 6 fl oz (12 Tbs) / 4 gal water and apply as above. Make first application in spring when new shoot growth is 1/2 to 2 inches in length. Make additional applications at 3 to 4 week intervals until dry conditions return.

Lophodermium Needlecast: Mix 1 fl oz (2 Tbs) / 4 gal water and apply full coverage spray to the point of runoff. North Central and Northeastern States: Begin applications in mid-July to early August before infection occurs. Make additional applications at 3 to 4 week intervals until dry conditions return.

Combination Sprays with Insecticides - on Ornamentals

ORTHO Multi-Purpose Fungicide DACONIL 2787® may be used together with the following insecticides on ornamentals (at the

label rates for both products): ORTHO Liquid SEVIN® Brand Carbaryl Insecticide, or ORTHO DIAZINON PLUS® Insect Spray, or ORTHO MALATHION 50 PLUS® Insect Spray, or ORTHENE® Systemic Insect Control, or ISOTOX® Insect Killer Formula IV. Apply fungicide on a regular schedule to control disease and add insecticide only when insect control is desired. Do not apply more than three consecutive applications of combination sprays.

STORAGE AND DISPOSAL

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area.

DISPOSAL: PRODUCT - Partially filled bottle may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash.

CONTAINER - Do not reuse empty bottle. Rinse thoroughly before discarding in trash.

ORTHO

**Multi-Purpose Fungicide
Daconil 2787® Plant Disease Control**

For effective control of a broad spectrum of lawn, vegetable, fruit, and ornamental plant diseases.
CONTROL: Leaf Spots, Rust, Blights, Fruit Rots, Mildews, Scab, Molds, etc. on Lawns, Vegetables, Fruit Trees, Flowers, Shrubs and Shade Trees.
Specially formulated for residential use.

Pool open for use directions and further information.
WARNING: Causes eye and skin irritation. This product may produce temporary allergic reactions or rash on exposed skin areas. Do not get in eyes, on skin or on clothing. SEE ADDITIONAL PRECAUTION STATEMENTS AND DIRECTIONS INSIDE BOOKLET.

Medical Information call 1-800-454-2233
Product Information call 1-800-225-2823

© Monsanto Company 1985
Manufactured for
The SOLARIS Group of
Monsanto Company
PO Box 5300
San Ramon CA 94583-0302
EPA Reg. No. 230-2522
EPA Est. 230-14-3
50236-MO-1*
Superscript is first letter of lot number
Made in USA



PRESS TO RESEAL



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

1sp = teaspoon / Tbs = Tablespoon

DIRECTIONS: Readily mixes with water and can be used in an ORTHO® SPRAY-ETTE® or tank-type sprayer. Shake well before using. Fine spray required for proper coverage; use an ORTHO SPRAY-ETTE for best results.

LAWN DISEASES

Place a small amount of water in sprayer, add fungicide, then add water to desired level and agitate gently to mix. Apply spray evenly. See below for rate and coverage. Apply after mowing. Withhold watering until spray has dried. Do not apply when temperature is above 90°F to prevent temporary discoloration of grass lawns. Do not graze treated areas or feed treated clippings to animals.

Important: Proper identification of lawn disease is essential for control. Take a specimen of the disease in a professional garden center or contact the County Extension Service for positive identification before treatment.

GRASS LAWN DISEASES

Copper Spot (Gleocercospora), Curvularia Leaf Spot, Dollar Spot (Sclerotinia), Gray Leaf Spot (Panicum), Gray Snow Mold (Typhula), Helminthosporium Leaf Spot and melting out.

AMOUNT TO USE

GALLONS OF SPRAY DESIRED	TO COVER LAWN AREA (SQ FT)	FOR ESTABLISHED DISEASE** UNDER SEVERE CONDITIONS	FOR LIGHT DISEASE** UNDER MILD CONDITIONS
2	333	4 fl oz (8 Tbs)	2 1/2 fl oz (5 1/2 Tbs)
4	667	8 fl oz (16 Tbs)	5 1/2 fl oz (10 1/2 Tbs)
6	1,000	12 fl oz (24 Tbs)	8 fl oz (16 Tbs)

*During periods of very wet or damp weather, make 3 successive applications at 7 day intervals. Continue treatments following rates for Light Disease incidence and interval listed below.

**Apply at a 10 to 14 day interval depending on severity of disease and conditions favoring disease development until lawn has recovered.

VEGETABLE DISEASES

Apply at listed rate per crop for disease to be controlled in one gal of water to cover 200 sq ft (10' x 20').

Large Brown Patch (Rhizoctonia), Red Thread (Corticium), Stem Rust of Bluegrass (Puccinia), Dichondra Diseases-(CA only), Alternaria Leaf Spot, and Anthracnose (Gleosporium).

Repeat at 7 to 14 day intervals. Use the 7 day schedule during severe disease conditions or wet weather unless otherwise stated. Note: Avoid spraying plants during extremely hot and

ORNAMENTAL DISEASES (continued)

Species	Diseases Controlled	Suggested First Applications
BULBS AND FLOWERING PLANTS		
Note: Use 1 1/4 fl oz (2 1/4 Tbs) / 4 gal	Black Spot Botrytis Blight	Spring bud break
Species	Anthracnose Cercospora Alternaria Botrytis Leaf Blight	Spring bud break
Zinnia	Powdery Mildew	First sign of disease

*Discoloration of plants has been noted on certain varieties when applications are made during flowering.

ORNAMENTAL DISEASES (continued)

Species	Diseases Controlled	Suggested First Applications
FOLIAGE PLANTS		
Ornamentals	Fusarium Leaf Spot	Pre-vegetation
Pachyandra - Use 3 1/2 fl oz (6 1/2 Tbs) / 4 gal	Botrytis Leaf Blight	Spring bud break
Leatherleaf fern	Ascochyta Blight Cercospora Leaf Spot Cylindrocladium Leaf Spot Rhizoctonia Blight	Spring bud break
Portulaca (Chamaedorea)	Botrytis Leaf Spot (Phoma/Phoma)	Cool, moist conditions
Prayer plant (Maranta)	Helminthosporium Leaf Spot	Early propagation
Oyster plant (Pilea)	Tan Leaf Spot	Early propagation
Platanus	Phytophthora Blight Dactyloctenium Leaf Spot	Moist conditions
Sparganium	Cercospora Leaf Spot	Warm, moist conditions

Blight (Blast), Purple Blotch and Downy Mildew (Suppression) (1 Tbs). Apply at first sign of disease and repeat at 7 to 10 day intervals. Do not apply more than 3 times per season on green onions, leeks or shallots.

• Potato (0): Early Blight, Late Blight, Botrytis Vine Rot (1½ tsp). Begin when plants are 6 inches high or when disease threatens and continue at 7 to 10 day intervals or as needed to maintain disease control.

• Tomato (0): Early Blight, Late Blight, Gray Leaf Spot, Gray Leaf Mold, Septoria Leaf Spot, Anthracnose, Alternaria Fruit Rot or Black Mold, Rhizoctonia Fruit Rot, Botrytis Gray Mold (1 Tbs). Begin when conditions favor disease development. Repeat at 7 to 10 day intervals or more frequently under severe conditions.

FRUIT DISEASES

Spray to thoroughly cover tree canopy including undersides of leaves. For best results, use an ORTHO® SPRAY-E-TE® Hose-End sprayer.

• Peach, Nectarine, Apricot, Cherry, Plum, Prune: Peach Leaf Curl; Coryneum Blight (Shothole) on Peach, Nectarine, Apricot: Mix

1½ fl oz (3 Tbs) in 4 gal of water (3 fl oz [6 Tbs] / 8 gal water) and spray foliage to point of runoff. Make one application at leaf fall in late autumn. Make one or two additional applications in mid to late winter before buds begin to swell. Where Coryneum Blight (Shothole) occurs, apply an additional application one to two weeks after petal fall or at shucksplit (tiny fruit just beginning to form) to prevent fruit infections. Brown Rot Blossom Blight – Apply at same rates. Make one application at popcorn (pink, red or early white bud stage when first color shows) and a second application at full bloom. If cool, moist weather continues, make an additional application at petal fall. Cherry Leaf Spot on Cherries: Scab on Peach, Nectarines and Apricot – In addition to the bloom applications listed above, make one application at shucksplit. Do not apply ORTHO Multi-Purpose Fungicide DACONIL 2787® after shucksplit or before harvest. For control of Cherry Leaf Spot after harvest, make one application to foliage within 7 days after fruit is removed. In trees with a history of high leaf spot incidence, make a second application 10 to 14 days later.

• Papaya: Alternaria Fruit Spot, Anthracnose and Stem End Rot: Apply full coverage spray including fruit at the rate of 1 fl oz (2 Tbs) / 4 gal water. Begin treatment when fruit spots appear (April to July) and continue treatments at 14 day intervals until dry weather conditions return.

• Passion Fruit (Hawaii only): Alternaria Fruit and Leaf Spot (Passion Fruit Brown Spot): Apply full coverage spray, including fruit, at the rate of 1/2 fl oz (1 Tbs) / 4 gal water. Begin treatment before spots appear on fruit (April to July) and continue treatments at 14 day intervals until dry weather conditions return.

ORNAMENTAL DISEASES

Mix 2½ tsp / gal or 3 Tbs (1½ fl oz) / 4 gal of water and spray foliage to point of runoff unless other directions are given in the table at right. Begin applications, as directed for each species and disease condition cited, and repeat on a 7 to 14 day schedule until conditions are no longer favorable for disease development. During periods when conditions favor severe disease incidence, generally cloudy or wet weather, use the shorter interval between applications. For outdoor use only.

ORNAMENTAL DISEASES

Species	Diseases Controlled	Suggested First Applications
BROADLEAF SHRUBS AND TREES		
Ash (Fraxinus)	Cercospora, Cercosporidium, Cylindrosporium Leaf Spots	Spring bud break
Azalea, Rhododendron	Phytophthora Die-back; Oudemans Flower Blight	New leaf emergence, Early bloom
Buckeye, Horsechestnut	Leaf Blotch Anthracnose	Spring bud break
Cherry-Laurel	Cercospora Leaf Spot	Petal fall
Crabapple	Scab, Cedar-apple Rust, Sphaeria Leaf Spot	Spring bud break
Dogwood	Anthracnose (Discalis sp.) Septoria Leaf Spot	Spring bud swell, Early bloom
Euonymus	Anthracnose	Spring bud break
Firethorn (Pyracantha)	Scab	Spring bud break
Flowering Almond, Quince and Cherry Hawthorn	Melanconia Blossom/Branch Blight, Rust, Fabraea Leaf Spot	Early bloom, Pre-bloom
Holly	Rhizoctonia Web Blight	Warm, moist conditions
Mountain Laurel	Cercospora Leaf Spot	Spring bud break

ORNAMENTAL DISEASES (continued)

Species	Diseases Controlled	Suggested First Applications
BROADLEAF SHRUBS AND TREES		
Oak (Red group only)	Tapinoma Blister, Actinopeltis Leaf Spot, Anthracnose	Dormant bud swell
Oregon-Grape (Mahonia)	Rust	Spring bud break
Photinia	Fabraea (Helminthosporium) Leaf Spot	Spring bud break
Pieris (Andromeda)	Phytophthora Die-back	New leaf emergence
Poplar	Marssonina Leaf Spot	Spring bud break
Privet	Cercospora Leaf Spot	Prolonged wet conditions
Sycamore, Planetree	Anthracnose	Spring bud break
Viburnum	Powdery Mildew	Mid-summer

ORNAMENTAL DISEASES (continued)

Species	Diseases Controlled	Suggested First Applications
BULBS AND FLOWERING PLANTS		
Carnation	Alternaria Leaf Spot/ Branch Rot, Botrytis	Transplant of cuttings, Cool, moist conditions
Chrysanthemum/ Daisy	Mycosphaerella Ray Blight, Septoria Leaf Spot	Transplant of cuttings
Geranium	Botrytis Flower Blight (Gray Mold), Rust	Pre-bloom, Cool, moist conditions
Gladolus	Curvularia Leaf/ Flower Spot, Botrytis Leaf/ Flower Spot	Early propagation
Hollyhock	Rust	Early seedling stage
Hydrangea (Hollige only)	Cercospora and Septoria Leaf Spots, Rust	Early propagation
Iris	Botrytis Blossom Blight, Didymella Leaf Spot	Cool, moist conditions
Lily	Botrytis Gray Mold (Polar phase), Phytophthora Blight	Pre-bloom
Petunia	Botrytis Gray Mold (Polar phase), Botrytis Blight	Pre-bloom

sunny weather. Do not apply to wilted plants. Do not graze treated areas or feed treated plant parts to animals.

VEGETABLES

Numbers in () indicate days required between last application and harvest.

• Beans (Green) (7): Rust and Botrytis Blight (Gray Mold) (1 Tbs)-Begin applications during early bloom stage or when disease first threatens. Repeat at weekly intervals or as necessary.

• Broccoli (0), Brussels Sprouts (0), Cabbage (0), Cauliflower (0): Alternaria Leaf Spot and Downy Mildew (1½ tsp). Begin after transplants are set in the garden or when conditions favor disease development. Repeat at 7 to 10 day intervals.

Ring Spot in Brussels Sprouts (CA only) (2 tsp). Begin at time of early sprout development or when conditions favor disease development. Repeat at 7 to 10 day intervals.

• Carrots (0): Cercospora (Early) Blight and Alternaria (Late) Blight (2 tsp). Begin when disease threatens and repeat at 7 to 10 day

intervals or as required to maintain control.

• Celery (7): Cercospora (Early) Blight and Septoria (Late) Blight (1½ tsp). Basal Stalk Rot and Pink Rot (Suppression) (1 Tbs). Begin when plants are set in the garden. Repeat on a 3 to 5 day spray schedule for lower rate (1½ tsp) or a 7 day schedule for the higher rate (1 Tbs).

• Corn (Sweet) (14): Helminthosporium Leaf Blight and Rust (2 tsp). Begin when conditions favor disease development. Repeat at 4 to 7 day intervals or as needed. Do not apply to sweet corn for processing.

• Cucumbers (0), Melons (0), Pumpkins (0), Squash (0): Anthracnose, Downy Mildew and Target Spot (2 tsp); Scab, Gummy Stem Blight (Black Rot), Leaf Blight, Cercospora Leaf Spot, Powdery Mildew (except Southwestern states) (1 Tbs). Also suppresses Fruit Rot (Rhizoctonia) on Cucumbers (1 Tbs). Begin when plants are in the first true leaf stage or when conditions are favorable for disease. Repeat applications at 7 day intervals or more frequently if required.

• Onion (Green) (14) and Dry Bulb (7), Leek (14), Shallot (14), Garlic (7): Botrytis Leaf