

ACCEPTED

DEC 23 1985

239-2526



ORTHO

MAKE.
32 GALLONS
OF DILUTED
SPRAY

Home Orchard Spray Formula II

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER: Causes irreversible eye damage. Harmful if swallowed. May cause allergic skin reactions. Do not get in eyes or on clothing. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Remove contaminated clothing and wash before reuse. Avoid breathing of spray mists or vapors. Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT: In case of eye contact, immediately flush eyes with fresh water for at least 15 minutes and see a doctor. If swallowed, do not make person vomit. Call a doctor immediately. If on skin, wash thoroughly with soap and water. If inhaled, move person from exposure area. Note to Physicians: Emergency Information - call (415) 233-3737. This product contains light petroleum solvent and may present an aspiration problem. Malathion is a cholinesterase inhibitor. If signs and symptoms of cholinesterase inhibition develop, atropine is antidotal. 2-PAM is also antidotal and may also be given in conjunction with atropine but should never be used alone.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS: Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

HOW TO USE: SHAKE WELL FOR 1 TO 2 MINUTES BEFORE MEASURING. Mix at the rate of 2 Tablespoons per gallon of water unless otherwise indicated. Spray thoroughly, contacting all upper and lower leaf surfaces and fruits until entire tree is dripping wet. For medium size trees (6 to 8 ft. high) apply 2 to 3 gals. of diluted spray per tree. Large trees may require 6 gals. or more. If any of the following insects or diseases attack any of the listed crops, follow the application directions for the particular crop. Numbers in () indicate the number of days required between last application and harvest. Make new dilution for each use. Do not apply during periods of hot-humid weather as injury may occur.

INSECTS: Aphids, Apple Maggot, Cankerworm, Codling Moth, Curculios, Flea Beetles, Japanese Beetle, Leafhoppers, Leafrollers, Oriental Fruit Moth, Pear Psylla, Pear Slug, Spider Mites, Spittlebug, Strawberry Leaf Beetle, Strawberry Weevil, Sucking Plant Bugs, Tent Caterpillars, Thrips.

DISEASES: Black Rot (fruit & leaf spots), Bitter Rot, Botryosphaeria, Botrytis Rot, Brooks Fruit Spot, Brown Rot, Fly Speck, Scab.

- A Multi-Purpose Spray to Control Many Fruit Insects and Diseases
- Controls: Aphids, Brown Rot, Codling Moth, Scab, Apple Maggot, Tent Caterpillar, Japanese Beetle and many other listed pests
- EZ-E to Use Liquid Formula

SHAKE
VIGOROUSLY
BEFORE
USING


Active Ingredients	By Wt.
*Captan	14.2%
Related Derivatives	0.8%
**Malathion	7.5%
***Methoxychlor Technical	15.0%
Inert Ingredients	62.5%
*N-(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide	
**O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate	
***Equivalent to 13.2%, 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 1.8% of other isomers and related compounds.	

Keep out of reach of children

DANGER

See side panel for additional precautionary statements.

NET CONTENTS 1 QT.



Crops
*Apples** (7),
*Pears** (7)

When To Apply

Spray when new spring growth starts and repeat at 7 day intervals up to bloom (blossoms fully open). After petal fall, repeat at 10 day intervals. NOTE: DO NOT SPRAY DURING BLOOM.

If Apple Maggots are a serious recurring problem in your area, add ORTHO Liquid SEVIN (1 lbs./gal.) to your regular spray program beginning in late June and continue 2 to 3 more applications through July and early August.

Spray at petal fall, 2 weeks after petal fall and again 2 weeks before harvest.

Begin spraying at pink bud stage and repeat application every 10 days until bloom. After petal fall, repeat sprays at 10 day intervals. NOTE: Use 3 Tablespoonfuls per gallon of water if brown rot incidence is a serious problem.

Begin spraying when pests appear or at first sign of new blooms and fruit. Repeat at 10 to 14 day intervals.

**Cherries (7),
**Apricots (21)
**Peaches (21)

Strawberries (3)

*Excessive sprays may cause some russetting to fruit of Golden Delicious Apple and Bartlett Pear.

**To prevent Brown Rot Blossom Blight infection during full bloom, use ORTHOCIDE (Captan) Garden Fungicide (alone) to prevent damage to pollinating insects.

NOTE: For spraying fruit trees in dormant season use ORTHO Dormant Insect and Disease Control.

STORAGE AND DISPOSAL

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area. Do not store diluted spray. **DISPOSAL: PRODUCT**—Partially filled bottle may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash. **CONTAINER**—Do not reuse empty bottle. Rinse thoroughly before discarding in trash.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Chevron Chemical Company © 1985

Ortho Consumer Products Division
San Francisco CA 94120-7144 Richmond CA 94804-0036
Product 5535 Made in U.S.A.
Form 10514-C
EPA Reg. No. 239-2526 EPA Est. 239-MO-1