

VEGETABLE GARDEN INSECT CONTROL

ORTHO-KLOR® 1% DURSBN Lawn & Soil Granules controls certain garden insects on crops listed below. Apply just prior to planting and immediately mix thoroughly into soil to the specified depth, except where applied with the seed in the seed furrow. Do not make more than one application per season.

Note: One level measuring cup (8 fl. oz.) holds 4 oz. of granules.

CROP	INSECTS CONTROLLED	AMOUNT TO APPLY
Sweet Corn	Cutworms, Wireworms, Flea Beetles, Beet Beetles, Flea Beetles, Flea Beetles	Apply (a, b) 8 oz. of granules per 100 sq. ft. evenly over the soil surface and work into the top 4-6 inches of soil.
Sweet Potato	Cutworms, Flea Beetles, Flea Beetles, Flea Beetles	Apply 5 1/2 oz. per 100 linear feet in the furrow with the seed at the time of planting.
Onions, Dry Bulb	Cutworms, Flea Beetles, Flea Beetles, Flea Beetles	Apply 5 1/2 oz. per 100 linear feet in the furrow with the seed at the time of planting.
Vegetables: Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Chinese Cabbage, Cauliflower, Kohlrabi, Lettuce, Peas, Beans, Turnips	Cutworms, Flea Beetles, Flea Beetles, Flea Beetles	Apply 8 oz. per 100 linear feet in the furrow with the seed or transplants at the time of planting. Note: Do not use Rotabag® bags for food or feed.
Radishes	Cutworms, Flea Beetles, Flea Beetles, Flea Beetles	Apply 5 oz. per 100 linear feet in the furrow with the seed at the time of planting.

HOME INVADING PESTS

PERIMETER TREATMENT AROUND FOUNDATION (OUTSIDE)

The following pests are commonly found around or near the foundation of houses. Many of these insects will enter home seeking food and shelter.

Ants	Crickets	Fleas	Brown Dog Ticks
Earwigs	Cloves Mites	Sawflies	Mites
Spiders (In Black)	Cockroaches	Fleas	Silverfish
Widow Spiders			

To control and prevent entry of the above pests, apply this product in a band 2-4 feet wide around and adjacent to the building using an ORTHO Whirllybird Hand Spreader when pests are active and may feed entrance. Thoroughly treat garbage can storage areas, and on porches, patios, around dog houses and pet enclosures. Do not treat rats with this product. See table below for coverage.

Note: ORTHO Whirllybird Hand Spreader (One level, hopperful) holds 1 1/2 cups of product.

Band Width is	One Level, Whirllybird Hand Spreader (1 1/2 cups of product)
2 ft	90 ft
4 ft	45 ft
6 ft	23 ft

HOW TO USE: Apply uniformly to home lawns and other ornamental and recreation turf grass areas using an ORTHO Drop or Broadcast Spreader, ORTHO WHIRLYBIRD Spreader, hand or power operated gravity or rotary type spreader designed for application of granular materials. Treat when insects or signs of feeding damage first appear. Retreat as necessary if reinfestation occurs. For Mole Crickets, apply in May or June when pests are small. Keep children and pets off treated areas until this material is washed into turf and grass is dry. Consult your local Agricultural Extension Service Specialist for proper time to treat in your area.

To Control Surface and Foliage Feeding Pests: Ants, Chinch Bugs, Cutworms, Flea Armyworms, Earwigs, Crickets, Brown Dog Ticks, Fleas, Leafhoppers, Mole Crickets, Lawn Moth (Soil Webworms), Chiggers, Clover Mites, Grasshoppers, Spittlebugs, Sowbugs, Pillbugs.

Apply 2 1/2 lbs. of ORTHO-KLOR® 1% DURSBN Lawn & Soil Granules per 1,000 sq. ft. (40' x 25'). Immediately after application, give the lawn a light to moderate watering. See table below for suggested spreader settings.

To Control Soil Insects — Mole Crickets and White Grubs (larvae of European Chafer, Southern Chafer and Japanese Beetle): Use double the rate for foliage feeding insects, i.e. 5 lbs./1,000 sq. ft. (see table). Note: An easy way to apply the higher rate is to spread the ORTHO-KLOR® 1% DURSBN Lawn & Soil Granules at the regular rate (2 1/2 lbs./1,000 sq. ft.) using the spreader settings below in one direction, then immediately re-apply a second application at the same rate (2 1/2 lbs./1,000 sq. ft.) at right angles to the first direction. This helps assure a more uniform distribution for better control. Thoroughly water immediately after application to wash the insecticide into the soil. For best results, treat when the grubs are young and actively feeding near the soil surface, usually during late July and August.


Weevils (Turfgrass): Use the higher rate (5 lbs./1,000 sq. ft.) described above under White Grubs. Treat suspected areas in mid-April and again in mid-May.

Suggested Spreader Settings

Make of Spreader	2 1/2 lbs./1,000 sq. ft.	5 lbs./1,000 sq. ft.
ORTHO Drop Spreader	7 1/2	9
ORTHO Broadcast Spreader	4 1/2	6 1/2
Scotts FF-1	7	9 1/2
Scotts FF-1	6	8
Cyclone	5 1/2	7
ORTHO WHIRLYBIRD Spreader		

An ORTHO WHIRLYBIRD Spreader will hold 1 1/2 cups of product when the hopper is filled level to the top. This will cover 90 square feet at the low rate or 250 square feet at the high rate.

Note: Above spreader settings are approximate. Accurately weigh a few pounds into your spreader and apply to measured area. Then adjust spreader setting if necessary to assure correct rate of application.



STORAGE AND DISPOSAL

STORAGE: Keep inside in original container. Do not put into food or drink containers. Avoid contamination of feed and feed-stuffs. Store in a cool, dry place, preferably in a locked storage area.

DISPOSAL: PRODUCT — Partially filled bag may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash. **CONTAINER** — Do not reuse empty bag. Discard bag in trash.

NOTICE: Buyer assumes all responsibility for safety of this product in accordance with directions.

Chevron Chemical Company 1986

Ortho Consumer Products Division
P.O. Box 5047 San Ramon CA 94583 0947
Product 5602 Made in U.S.A.
Form 10870 C
EPA Reg. No. 239-2570
EPA Est. 239-MO-1

NET WT. 10 LBS.

(
ORTHO-KLOR 1% DURSBA~~N~~awn & Soil Granules
EPA Reg. No. 239-2570

3

COPY A

CONTROLS SURFACE AND FOLIAGE FEEDING PESTS

Sod Webworms, Chinch Bugs, Grasshoppers, Armyworms, Crickets, Chiggers,
Fleas, Ticks

SOIL INSECTS


Grubs, Mole Crickets

HELPS CONTROL HOME INVADING PESTS

Ants, Cockroaches, Earwigs, Sowbugs, Silverfish, Spiders

VEGETABLE GARDEN INSECTS

Wireworms, Cutworms, Root Maggots, Billbugs



CONFIDENTIAL

PROPERTY OF
CROVON CHEMICAL COMPANY
UNTIL ACCEPTED BY EPA