


DRAFT #7
JAC:kt/TBL3
3/23/88

Note: EPA Reg. Label
Includes Text for all Sizes

BRUSH-B-GON® Poison Ivy & Poison Oak Killer

- o Kills Woody Plants, Weeds, Vines and Brambles - Roots and All.
- o Kills Poison Ivy, Poison Oak, Blackberries, Kudzu, Willows and Other Listed Woody Plants.
- o Kills Weeds - Dandelion, Knotweed Plantains, Oxalis, Spurge, Thistles, Wild Morningglory, and other listed weeds.
- o Ready-to-Use, No Mixing.



CONFIDENTIAL
PROPERTY OF
CHEVRON CHEMICAL COMPANY
UNTIL ACCEPTED BY EPA

(Illustration)

Active Ingredient	By Wt.
*Triclopyr (3,5,6-trichloro-2 pyridinyl - oxyacetic acid) as the triethylamine salt.....	0.70%
Inert Ingredients	99.30%


*Equivalent to 0.50% acid

Keep out of reach of children.

CAUTION

See back panel for additional precautionary statements.

NET CONTENTS 1/2 GAL.



CONFIDENTIAL

2
3

PROPERTY OF

CHEVRON CHEMICAL COMPANY

Back Panel

UNTIL ACCEPTED BY EPA
BRUSH-B-GON® Poison Ivy & Poison Oak Killer

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS.

CAUTION: Harmful if swallowed. Avoid contact with eyes. Avoid breathing spray mist. Wash thoroughly after handling. **STATEMENT OF PRACTICAL TREATMENT:** If swallowed, induce vomiting by administering syrup of ipecac or under medical supervision. Call a physician. In case of eye contact, immediately flush eyes with fresh water for 15 minutes. If irritation continues, see a doctor. If inhaled, remove person from exposure area. **Note to Physicians:** Emergency Information - call (415) 233-3737.

ENVIRONMENTAL HAZARDS: Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Formulated for "spot" application to kill unwanted vegetation, vines, brambles and weeds in areas around homes, cabins, buildings, fences, trails, and in non-crop areas. Do not use on or around edible feed or food crops.

Kills Woody Plants, Weeds, Vines and Brambles

o Woody Plants, Vines and Brambles

- | | | | |
|-------------------|----------------------|---------------------|---------------------|
| Arkansas Rose | Evergreen Blackberry | Maple* | Scouler Willow* |
| Big Leaf Maple* | Field Horsetail Rush | Narrow Leaf Willow* | Sweet Gum* |
| Black Cottonwood* | Giant Ragweed | Oak* | Trailing Blackberry |
| Black Willow* | Himalayan Blackberry | Persimmon* | Trumpet Creeper |
| Box Elder | Honeysuckle | Poison Ivy | Virginia Creeper |
| California Rose | Kudzu | Poison Oak | Wild Blackberry |
| Choke Cherry* | Mackensie Willow* | Raspberry | Wild Grape |
| Common Ragweed | Mesquite* | Red Alder* | Wild Rose |
| Elderberry* | Mimosa* | Sassafras* | Willow* |
| Elm* | | | |

*small volunteer plants or "suckers" only

o Weeds (Not in Lawn Areas)

- | | | |
|-------------------------------|---------------------|---------------------|
| Bindweed (Wild Morning-Glory) | Creeping Woodsorrel | Narrowleaf Plantain |
| Broadleaf Plantain | Curly Dock | Oxalis |
| Buckhorn Plantain | Dandelion | Purple Nutsedge |
| Buttercup | English Daisy | Purslane |
| Canada Thistle | Florida Pusley | Spurge |
| Carpetweed | Ground Ivy | Tansy Ragwort |
| Chickweed | Horsetail Rush | White Clover |
| | Knotweed | Wild Garlic |

WHEN TO APPLY: Apply when vegetation is fully leafed out and actively growing. Wilting or other symptoms (i. e. yellowing) will occur in two to six weeks. For best results, apply to mature leaves. Apply only when air is calm to prevent drift to desirable vegetation. Rainfall within 24 hours may reduce effectiveness of treatment.

HOW TO USE: Shake before using. Apply with hand trigger sprayer. Fold back protective hinged tab exposing nozzle and snap in place on sprayer top. Adjust sprayer nozzle to allow a coarse spray for applying to leaves and stems of unwanted vegetation. Stand close to weed (one to two feet). Aim spray nozzle at weed and away from face. Be careful when applying this product not to spray surrounding desirable lawn grasses, groundcovers, or nearby plants as they may also be killed. If desirable plants are accidentally sprayed, wash them immediately with water. Hard-to-control plants, such as Poison Oak, Poison Ivy, Blackberries, and Honeysuckle may require retreatment the following season if any regrowth occurs.

IMPORTANT: BRUSH-B-GON Poison Ivy & Poison Oak Killer is non-selective. When treating brushy vines such as Poison Ivy, Poison Oak, Kudzu, etc. growing on or around desirable plants, do not allow spray to contact desirable vegetation or ornamental plants as injury may result. Do not apply when drift may occur.

NOTE: For treating mature plants in larger areas, use BRUSH-B-GON Brush Killer - a concentrated product.

STORAGE AND DISPOSAL

STORAGE: For storage, reseal by replacing tab closure over spray nozzle. Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area. Keep from freezing. **IMPORTANT:** After each use, store sprayer nozzle under handle of container above liquid level of contents to prevent leakage and siphoning from nozzle.

DISPOSAL: PRODUCT - Partially filled bottle may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash. **CONTAINER** - Do not reuse empty bottle. Rinse thoroughly before discarding in trash.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Chevron Chemical Company @ 1987
ORTHO Consumer Products Division
P.O. Box 5047 San Ramon CA 94583-0947
Product 5609 Made in USA
Form _____ EPA Reg. No. 239-_____
EPA Est. 239-IA-3

CONFIDENTIAL

PROPERTY OF

CHEVRON CHEMICAL COMPANY

UNTIL ACCEPTED BY EPA

JAC:kt/TBL3