

239-2640

09/22/2006

1/8

Master Label

GreenSweep Weed & Feed 20-0-0
EPA Reg. NO. 239-2640

[sticker]

Greens
Your Lawn
In 24 HOURS
AND Kills Weeds!

ACCEPTED

SEP 22 2006
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

239-2640

[FRONT]

FEEDS And TREATS
UP TO
6,000
SQUARE FEET

GreenSweep Weed & Feed

[Note to PM: Optional Front Panel Claims]

- Kills dandelions, clover and other listed lawn weeds
- Easy to use! Just attach hose and spray!
- Contains patented slow-release nitrogen for extended feeding.
- Gives your lawn a quick boost of nitrogen while controlling weeds.
- Quickly feeds your lawn while controlling weeds.
- Fertilizes lawn and kills broadleaf weeds.
- Connect & Spray

NET 1 QT (32 FL OZ) 946 mL, 2.3 lb

Active Ingredients

2,4-D, Dimethylamine Salt*	2.29%
Mecoprop-p, Dimethylamine Salt**	1.15%
Dichlorprop-p, Dimethylamine Salt***	1.13%
Other Ingredients	95.43%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel booklet for additional precautionary statements.

XX-X-XX

UPC

[Place this form number vertically on side] LF07630X000

[BACK COVER PAGE]

[corner flap]

OPEN Resealable Label for Directions and Precautions

PRESS TO RESEAL [on base of label]

OPEN

GreenSweep Weed & Feed

XX-X-X

GUARANTEED ANALYSIS

F643

Total Nitrogen (N)	XX%
XX% Urea Nitrogen	
XX5% Other Water Soluble Nitrogen*	
Available phosphate(P ₂ O ₅)	X%
Soluble potash (K ₂ O)	X%

Derived from: triazone, urea, and potassium pyrophosphate.

*Slowly Available Water Soluble Nitrogen from triazone.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-sc.com>

[Optional]Specially formulated for residential use.

The Ortho Group
14111 Scottslawn Rd.
Marysville, OH 43041
Form LB07622T000
EPA Reg. No. 239-2640
EPA Est. 239-IA-3^I, 58996-M0-1^A
Superscript is first letter of lot number
Made in USA

[Note to PM: Optional back panel ³ claims]

GreenSweep Weed & Feed fertilizes lawns and kills broadleaf weeds in one easy step.

A beautiful, weed-free lawn is just a spray away.

Controls broadleaf weeds and gives your lawn a quick boost of nitrogen that lasts for weeks.

Use on bluegrass, fescue, ryegrass, bahiagrass, common Bermudagrass, centipedegrass and zoysiagrass. When used as directed, Scotts Liquid Turf Builder supplements your regular fertilizer program.

To spot treat lawn weeds try Ortho® Weed-B-Gon®. It Kills Weeds, Not Lawns. (image of WBG 24 oz. RTU)

[phone icon] Questions, Comments or Medical Information call 1-800-543-TURF (8873) www.scotts.com

WEEDS CONTROLLED

Alder, Artichoke, Aster, Austrian fieldcress, Beggartick, Bider, Bitterweed, Black medic, Box elder, Broomweed, Buckhorn, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chickweed, Chicory, Clover, Dandelion, Dock, Elderberry, English daisy, Florida pusley, Goldenrod, Ground ivy, Healall, Henbit, Honeysuckle, Horsetail, Ironweed, Knotweed, Lambsquarters, Mallow, Morning glory, Musk thistle, Mustard, Nettle, Nutgrass, Oxalis, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison Ivy, Poison Oak, Ragweed, St. Johnswort, Sheep sorrel, Shepherdspurse, Smartweed, Sowthistle, Speedwell, Spurge, Spurweed, Stinkweed, Sumac, Sunflower, Thistle, Tumbleweed, Velvet leaf, Veronica, Vetch, Virginia buttonweed, Virginia creeper, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Woodsorrel, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

[END OF OPTIONAL CLAIMS]

This product contains:

- *2,4-Dichlorophenoxyacetic Acid 1.9%, .18 lb/gal
- ***+(R)-2-(2-Methyl-4-Chlorophenoxy)propionic Acid .96%, .18 lb/gal
- ***+(R)-2-(2,4-Dichlorophenoxy)propionic Acid.. .96%, .18 lb/gal
- Isomer Specific AOAC Method No. 6.321.

[UPC]

4/8

[BACK]

[NOTE TO PM: DIRECTIONS FOR USE OPTION 1]

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

HOW TO APPLY:

- Shake well before using. Connect sprayer to hose. Turn on water.
- To begin spraying, point nozzle in the direction you want to spray. Bend small yellow plastic tab on side of sprayer back and turn knob clockwise to ON position.
- Spray evenly over measured area.
- To stop spraying, bend small yellow plastic tab back and turn knob counter clockwise to OFF position. Turn off water.
- Relieve water pressure by bending plastic tab back and turning knob to ON position until water slows to a drip. Then, turn knob back to OFF position.
- Disconnect sprayer from hose.

AMOUNT TO APPLY

Apply only the recommended amount for the area to be covered. Exceeding recommended rates may damage turf.

- **WARM SEASON LAWNS** - bahia, common Bermuda, centipede and zoysia.
1 quart covers 6,000 sq ft.

NOTE: Some temporary injury may occur with these lawns. In spring, do not apply until grass has reached full green-up. In fall, do not use when grass is going into dormancy. Some hybrid Bermuda grasses may be sensitive to this product.

- **COOL SEASON LAWNS** - bluegrass, fescue and ryegrass
1 quart covers 4,000 sq ft.

5/8

FOR BEST RESULTS

- If lawn needs watering, water thoroughly 24 hours before application.
- Do not cut grass 1 to 2 days after application.
- Do not water for 48 hours after application as watering will wash off weed killing material.
- Spray when air is calm to avoid spray drift that might injure desirable ornamental plants.
- Do not apply when temperatures exceed 85°F.
- Do not reseed, sod or plug lawns, or plant stolons for 3 to 4 weeks after use. Do not apply to newly seeded lawns until they have been mowed three times.
- Spray when weeds are actively growing.
- [RESULTS:] Weeds will begin to curl within a few days. Some hard-to-control weeds may require a repeat application in 3 to 4 weeks.

IMPORTANT

For grass lawns only. Do not spray St. Augustinegrass, Buffalograss, dichondra, lippia or desirable clovers. Do not use this product on flower or vegetable beds, or around shrubs, trees or ornamental plantings. The maximum number of broadcast applications per treatment site is 2 per year.

[NOTE TO PM: END OF DIRECTION FOR USE OPTION 1]

6/8

[NOTE TO PM: DIRECTIONS FOR USE OPTION 2 for Private Label Branding]

For grass lawns only. Do not use this product on flower or vegetable beds, or around shrubs, trees or ornamental plantings.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

HOW TO USE:

- Make sure ON/OFF valve is in OFF position. Shake well.
- Connect sprayer nozzle to hose. Turn on the water at the tap.
- When ready to treat, point in the direction away from face and body and pull toward the hose end connection.
- Turn the ON/OFF valve to the ON position. The water mixes automatically with this product at the proper ratio.
- Walk and spray at steady pace.
- Apply only the recommended amount for the area to be covered. Exceeding recommended rates may damage turf.

For COOL SEASON LAWNS such as Bluegrass, Fescue and Ryegrass: 1 quart covers 3,300 sq ft. When applying to Bentgrass, grass height should 1½ inches or more.

For WARM SEASON LAWNS such as Bahia, Common Bermuda, Common St. Augustine, Centipede and Zoysia. 1 quart covers 5000 sq ft. Do not use on Floratam or other improved varieties of St. Augustinegrass. Some temporary injury may occur with these lawns. In spring, do not apply until grass has reached full green-up. In fall, do not use when grass is going into dormancy. Note: Some hybrid Bermuda grasses may be sensitive to this product.

RESULTS: Weeds will begin to curl within a few days. Some hard-to-control weeds may require a repeat application in 3 to 4 weeks. The maximum number of broadcast applications per treatment site is 2 per year.

IMPORTANT USE PRECAUTIONS: Spray when weeds are actively growing. If lawn needs watering, water thoroughly 24 hours before application. Do not cut grass 1 to 2 days before and after application. Do not water for 48 hours after application as watering will wash off weed killing material. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Do not apply when temperatures exceed 85°F. Do not apply to newly seeded grasses until they have been mowed three times. Do not reseed, sod, or plus lawns or plant stolons for 3 to 4 weeks after use. Do not spray Buffalograss, dichondra, lippia or desirable clovers.

[NOTE TO PM: END OF DIRECTION FOR USE OPTION 2]
STORAGE AND DISPOSAL

STORAGE: Store product in original container in a safe place. Protect container from freezing.

DISPOSAL: **If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin or clothing. Wear long pants, long-sleeved shirt, socks, shoes and chemical resistant gloves when handling or applying this product. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

[Reentry icon] Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

FIRST AID:

If swallowed: A poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. **If in eyes** eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling poison control center or doctor, or going for treatment.

ENVIRONMENTAL HAZARDS: This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters nor pour the washwaters on the ground. Spray or drain

over a large area away from wells, other water sources and desirable plants. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes. If spills occur, collect the material and dispose of by following disposal instructions on this label.

NOTICE: Buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

-OR-

Scotts No-Quibble Guarantee®

If for any reason you, the consumer, are not satisfied after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

[phone icon] **Questions, Comments or Medical Information**
call 1-800-543-TURF (8873) www.scotts.com

©2003, Licensed by The Scotts Company. World rights reserved.

The Scotts Company
14111 Scottslawn Rd.
Marysville, OH 43041

Form LB07622T000
EPA Reg. No. 239-2640-538
EPA Est. 239-IA-3^I, 58996-M0-1^A
Superscript is first letter of lot number
Made in USA