

239-2664

04/11/2000

1/5

ORTHO® WEED B GON® READY-SPRAY® EPA Reg. No: 239-2664
ORTHO® READY-SPRAY® WEED-B-GON® LAWN WEED KILLER

Now Treats Northern and Southern Lawns

- Kills Weeds Not Lawns
- Kills Dandelion, Clover & Other Broadleaf Weeds
- Rain-Proof in Hours
- Treats up to 12,800 sq. ft


(See Appendix for additional claims)

NEW National Formula

Ready-Spray®
Connect / Attach & Spray

Active Ingredients

Mecoprop, dimethylamine salt	10.60%
2,4-D, dimethylamine salt	3.05%
Dicamba, dimethylamine salt	1.30%
Other Ingredients	85.05%



KEEP OUT OF REACH OF CHILDREN

CAUTION See back panel booklet for additional precautionary statements.

NET 32 FL OZ (1 QT) 946 m L

[icon] **WATER BASED**

OPEN

**Resealable Label
for Directions & Precautions
PRESS TO RESEAL**

Ready-Spray makes lawn and garden care easy. Simply attach the hose to the attached sprayer. Eliminates mixing, measuring and messy clean up. Automatically mixes the right amount of product and water.

OR

Kills weeds in your lawn - won't harm lawn grasses. Weed-B-Gon singles out weeds enters through leaves and moves inside the weed to the roots.

Makes lawn weed control easy. Simply connect hose to attached sprayer. Eliminates mixing, measuring and messy clean-up.

PRODUCT FACTS	
Treats up to 12,800 sq ft (approx. size of 6 tennis courts)	
KILLS WEEDS	Dandelion, Clover, Chickweed <i>[illustration]</i> plus over 80 additional broadleaf weeds
WHERE TO USE	ON LAWNS <i>[illustration]</i> without harming lawn grasses
AMOUNT TO SPRAY	Bent, Buffalo, Centipede & St. Augustine (EXCEPT Floratam): 8 oz (1/4 bottle) treats 3,200 sq ft
	Bahia, Bermuda, Fescues, Kentucky Blue, Perennial Rye & Zoysia 8 oz (1/4 bottle) treats 1,600 sq ft
†	Questions, Comments or Medical Information call 1-800-225-2883 www.ortho.com

THIS PRODUCT CONTAINS:

0.75 lb. 2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 8.76%
 0.22 lb. 2,4-Dichlorophenoxyacetic acid equivalent per gallon or 2.53%
 0.093 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 1.08%
 Isomer Specific by AOAC Method 6.DO1-5.

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

AMOUNT TO SPRAY

- Measure lawn. Calculate square feet by multiplying length times width. *[walking man icon]*
- Use the amount indicated for your lawn variety:
 [FOR SENSITIVE LAWNS:] **Bentgrass, Buffalograss, Centipedegrass & St. Augustinegrass (EXCEPT Floratam, a variety of St. Augustinegrass widely grown in Florida.):** 8 oz (1/4 bottle) treats 3,200 sq ft

[FOR HARDY LAWNS:] **Bahiagrass, Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass & Zoysiagrass:** 8 oz (1/4 bottle) treats 1,600 sq ft

HOW TO USE THE READY-SPRAY®	
CONNECT <i>[illustration]</i>	Connect sprayer to hose. Be sure on/off switch is set to "OFF". Turn on water.
<i>[illustration]</i>	Use coin or key to rotate small (color) button to "OPEN".
SPRAY <i>[illustration]</i>	To BEGIN spraying, point nozzle towards lawn. Turn on/off switch forward to "ON". Spray evenly over measured area.
FINISH <i>[illustration]</i>	• To STOP spraying, turn on/off switch back to "OFF". Rotate small (color) button to "CLOSED". Turn off water.

- Relieve water pressure by turning switch to "ON" until water slows to a drip. Turn switch back to "OFF". Disconnect from hose.

[Color of button is green.]

FOR BEST RESULTS	
MOW [illustration/ mower]	Do not mow for 1 to 2 days after application.
WATER [illustration /hose]	<ul style="list-style-type: none"> • If soil is dry, water 1 to 2 days before application. • Rain or watering 6 hours after application will not wash away effectiveness.
WHEN TO APPLY [illustration / weather]	<ul style="list-style-type: none"> • Spray when: <ul style="list-style-type: none"> - weeds are small and actively growing - temperature is below 90°F - air is calm to avoid spray drift onto landscape plants as contact with foliage may injure flowers, shrubs, trees and other desirable plants • Reapplication can be made in 2 to 3 weeks if necessary. • Newly seeded lawn can be sprayed after first mowing. • Bare spots can be planted with grass seed 1 week after application.

OR

WHEN TO [MOW & - optional] WATER [icon]

- Do not mow for 1 to 2 days before or after application. [optional]
- If soil is dry, water 1 to 2 days before application.
- Rain or watering 6 hours after application will not wash away effectiveness.

WHEN TO APPLY [icon]

- Spray when weeds are small and actively growing.
- [Spray when] temperature is below 90°F.
- [Spray when] air is calm to avoid spray drift onto landscape plants as contact with foliage may injure flowers, shrubs, trees and other desirable plants
- Reapplication can be made in 2 to 3 weeks if necessary.
- Newly seeded lawn can be sprayed after first mowing.
- Bare spots can be planted with grass seed 1 week after application.

IMPORTANT

If an ornamental shrub or tree is in the lawn, keep spray off foliage and exposed roots. St. Augustinegrass, Centipedegrass or Bermudagrass lawns may have delayed green-up if this product is used when lawns are emerging from dormancy or when temperature drops below 45° F within 72 hours of application. Do not spray Carpetgrass, Dichondra or desirable clovers. Do not use on or around fruits, vegetables or flowers. The maximum number of broadcast applications per treatment site is 2 per year. Do not exceed specified dosages for any area.

HOW IT WORKS

Because broadleaf weeds and grasses grow differently, Weed-B-Gon® kills broadleaf weeds without harming lawns. Weed-B-Gon singles out weeds, enters through leaves and moves inside the weed to the roots.

WEEDS [icon]

Annual Aster, Bedstraw, Beggarweed, Bindweed, Bittercress, Black Medic, Brassbuttons, Broadleaf Plantain, Buckhorn, Bull Thistle, Burdock, Bur Clover, Burweed, Buttercup, Buttonweed, Carolina Geranium, Carpetweed, Catsear, Chicory, Chickweed, Cinquefoil, Clover, Cocklebur, Common Chickweed, Cudweed, Curly Dock, Dandelion, Dock, English Daisy, False Dandelion, Florida Pusley, Galinsoga, Geranium, Ground Ivy, Hairy Fleabane, Hawkweed, Healall, Henbit, Knawel, Knotweed, Lambsquarters, Lespedeza, Mallow, Matchweed, Morningglory, Mouseear Chickweed, Musk Thistle, Mustard, Narrowleaf Plantain, Oxalis, Pennsylvania Smartweed, Pennywort (Creeping Charlie), Peppergrass, Pigweed, Piney Woods Bedstraw, Plantain, Poison Ivy, Poison Oak, Prickly Lettuce, Prostrate Spurge, Puncture Vine, Purslane, Ragweed, Red Clover, Red Henbit, Scarlet Pimpernel, Sheep Sorrel, Shepherdspurge, Sowthistle, Speedwell, Spotted Spurge, Spurge, Tansy Mustard, Thistle, Velvetleaf, White Clover, Wild Carrot, Wild Garlic, Wild Lettuce, Wild Onion, Wild Strawberry, Yarrow & other broadleaf weeds

STORAGE AND DISPOSAL [icon]

STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

PRECAUTIONARY STATEMENTS [icon]

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. Wear face shield or goggles when mixing, loading or applying this product. When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves.

Personal Hygiene Statements: After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

[Reentry icon] Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried.

FIRST AID: IF IN EYES: Immediately flush eyes with plenty of water. Get medical attention if irritation persists. **IF ON SKIN:** Wash with plenty of soap and water. Get medical attention. **Note to Physician:** Emergency information call 1-800-225-2883.

ENVIRONMENTAL HAZARDS: This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. ~~Do not contaminate water when disposing of equipment washwaters.~~ Do not apply when weather conditions favor drift from target area. ~~When~~

