239-	-266	5
------	------	---

5-22-2003

i	K

Please read instructions on reverse before	pleting form.	Form Appro	ved, OMB No.	2070-0060	). Approvel expires 2-28-95				
United States  Environmental Protection Agency Washington, DC 20460		<u></u>	Registra Amenda Other		OPP Identifier Number				
Application for Pesticide - Section I									
1. Company/Product Number 239-2665		Joanna Millor		posed Classification  None Restricted					
4. Company/Product (Name) ORTHO Weed-B-Gon lawn Weed Killer3		PM# 23							
5. Name and Address of Applicant (Include ZIP Code) The Scotts Company d/b/a The ORTHO Group 14111 Scottslawn Rd Marysville, OH 43041		6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to:  EPA Reg. No.							
Check if this is a new address		Product Name							
	Sec	ction - II							
Amendment - Explain below.  Resubmission in response to Agency le  Notification - Explain below.	tter dated	Final printed I. Agency letter "Me Too" App Other - Explai	dication.	NO	TIFICATION y 2 2 2003				
The ORTHO Group is submitting a notification o Notice 98-10 and EPA regulations at 40 CFR 15 product. I understand that it is a violation of 18 l consistent with the terms of PR Notice 98-10 and penalties under sections 12 and 14 of FIFRA.	2.46, and no other changes J.S.C. Sec. 1001 to willfully r d 40 CFR 152.46, this produ	have been made to the lab make any false statement t ct may be in violation of FIf	eling or the confi o EPA. I further	dential state understand	ement of formula of this that if this notification is not				
	Sec	tion - III							
1. Material This Product Will Be Packaged In:				^					
Child-Resistant Packaging  Yes  No  No  Unit Packaging  Water  Yes  No		No Pier Gla		Metal Plastic Glass	al ric ss				
* Certification must Unit Packaging v	No. per   If "Ye vgt. container   Packs	s" No. per ge wgt container	No. per Container Paper Other (Specify)		pecity)				
3. Location of Net Contents Information 4. Size(s) Retail Container 5. Location of Label Directions				ns					
6. Manner in Which Label is Affixed to Product Paper glued Stenciled  Other									
	Sect	tion - IV			, tenne no				
1. Contact Point (Complete items directly belo	w for identification of indiv	vidual to be contacted, if .	necessary, to pr	ocess this	application.)				
Name Title Charles T. Levey Manager, Federal		er, Federal Registrations	3	Telephone 937-644-7	No. (include Area Code) 696				
I certify that the statements I have made I acknowledge that any knowlingly false both under applicable law,		-			6. Date Application Received (Stamped)				
2. Signature/ Nan h auly		r, Federal Registrations							
4. Typed Name ( / // Charles T. Levey	5. Date	5. Date May 16, 2003							



May 16, 2003

Joanne Miller
U.S. Environmental Protection Agency
Office of Pesticide Programs
Registration Division, H7505C
Ariel Rios Building
1200 Pennsylvania Ave., N.W.
Washington, DC 20460

Subject:

Notification of additional claims in accordance with PR Notice 98-10 ORTHO Weed-B-Gon Lawn Weed Killer3, EPA Reg. No. 239-2665 and Weed-B-Gon Ready-Spray Isomer Formula, EPA Reg. No. 239-2666

#### Dear Joanne Miller:

The ORTHO Group is submitting into application additional marketing claims for the subject products. I have highlighted the changes on each of the enclosed final printed labels. Please find enclosed the following:

- 1) Application Form 8570-1 for each of the product registrations.
- 2) One (1) copy of the final printed label for each of the product registrations.

Please call me at 937-644-7696, if you have any questions regarding this submission. Thank you in advance for your experation.


Sincerely

The Scotts Company

Charles T. Levey


Manager, Federal Registrations





NOTIFICATION
MAY 2 2 2003

2.025



# \*WEEDS CONTROLLED (continued)

12

corn spurry, swinecress, bull thistle, Canada thistle, Flodman thistle, cotton thistle, distaff thistle, Drummonds thistle, elk thistle, gray thistle, Italian thistle, leafy thistle, meadow thistle, milk thistle, Platte thistle, plumeless thistle, Russian thistle, Scotch thistle, slenderflower thistle, tall thistle, wavyleaf thistle, yellowspine thistle, stiff verbena, camplipoda Veronica, common vetch, hairy vetch, winter vetch, wooley vetch, violet, field violet, wild carrot, wild lettuce, woodsorrel, creeping woodsorrel, yellow woodsorrel, Florida yellow woodsorrel, wooleyleaf bursage, common yarrow, western yarrow, yellow yarrow.

# STORAGE & DISPOSAL



STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

#### PRECAUTIONARY STATEMENTS



#### **HAZARDS TO HUMANS & DOMESTIC AHIMALS**

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. Wear face shield or goggles when mixing, loading or applying this product. When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves. Personal Hygiene Statements: After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.



Do not allow people (other than applicator) or pets on treatment area during application. People and pets may enter treated area after spray has dried.

# ORTHO

**OPEN** Resealable Laber for Directions & Precautions

Won't Harm Lawns - Guaranteed! Contains the special Ortho LawnGuard™ formula. How it works — Weed-B-Gon LawnGuard formula specifically targets weeds, killing them while leaving the grass unharmed – guaranteed. It's Rain-Proof™ Brand. Rain two hours after spraying won't wash away the weed killing power!

# PRODUCT FACTS

Treats up to 12,800 sq ft (approximate size of 6 tennis courts)

**WEEDS** 

Dandelion, Clover, Chickweed plus over 200 additional broadleaf lawn weeds\*



WHERE TO USE

ON LAWNS without harming lawn grasses

**AMOUNT** TO USE

Bentgrass, buffalograss, centipedegrass & St. Augustinegrass (DO NOT USE on floratam): 1 floz (2 Tbs) / gal of water / 400 sq ft Bahiagrass, Bermudagrass, fescues, Kentucky bluegrass, perennial ryegrass & zoysiagrass: 2 fl oz (4 Tbs) / gal of water / 400 sq ft



Questions, Comments or Medical Information? Call 1-800-225-2883 uww.ortho.com 🖳

For weeds outside the lawn (on driveways, sidewalks, etc.) — use Ortho GroundClear® Complete Vegetation Killer. It kills weeds and grasses and prevents new growth for up to 1 year.

Manufactured for The ORTHO Group P.O. Box 190 Marysville, OH 43040 Form LBXXXXXXXXXX EPA Reg. No. 239-2665 EPA Est. 239-1A-31, 58996-MO-1A Superscript is first letter of lot number Made în USA





1026

3

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

# MIXING INSTRUCTIONS

1 Tablespoon (Tby): 3 teaspoons (tsp)



FOR SENSITIVE LAWNS: Bentgrass, buffalograss, centipedegrass & St. Augustinegrass (DO NOT USE on floratam\*)

Amount to Use: 1 fl oz (2 Tbs) per gallon of water for each 400 sq ft. When using Ortho® Dial 'N Spray®: Set dial to 1 oz. Add 2 ½ fl oz (5 Tbs) to sprayer jar. DO NOT ADD WATER. Spray evenly over 1,000 sq ft until jar is empty.

Floratam is a variety of St. Augustinegrass widely grown in Florida.

FOR HARDY LAWNS: Bahiagrass, Bermudagrass, fescues, Kentucky bluegrass, perennial ryegrass & zoysiagrass

Amount to Use: 2 fl oz (4 Tbs) per gallon of water for each 400 sq ft. When using Ortho® Dial 'N Spray®: Set dial to 2 oz. Add 5 fl oz (10 Tbs) to sprayer jar. DO NOT ADD WATER. Spray evenly over 1,000 sq ft until jar is empty.

#### REFILL DIRECTIONS

To refill only an empty 1.33 Gallon Weed-B-Gon Ready-to-Use Pull 'N Spray® Bottle Amount to Use: Add 7 fl oz of this product to the empty 1.33 gallon container. Slowly fill with 160 fl oz of clean tap water.

Specially formulated for residential use.

### FOR BEST RESULTS

MEASURE

- To treat entire lawn (broadcast application): Use a tank or hose-end sprayer. Measure lawn. Calculate square feet by multiplying length times width.
- Spray evenly over measured area.
- · To spot treat individual weeds in lawns: Use a tank sprayer.

MOW

Do not mow for 1 to 2 days before or after application.



- If soil is dry, water 1 to 2 days before application.
- Rain or watering 2 hours after application will not wash away effectiveness.



- Spray when:
- weeds are small and actively growing
- temperature is below 90°F
- air is calm to avoid spray drift onto landscape plants as contact with foliage may injure flowers, shrubs, trees and other desirable plants
- · Reapplication can be made in 2 to 3 weeks if necessary.
- Newly seeded lawn can be sprayed after first mowing.
- Bare spots can be planted with grass seed 1 week after application.

# **IMPORTANT**

If an ornamental shrub or tree is in the lawn, keep spray off foliage and exposed roots.

St. Augustinegrass, centipedegrass or Bermudagrass lawns may have delayed green-up if this product is used when lawns are emerging from dormancy or when temperature drops below 45°F within 72 hours of application.

Do not spray carpetgrass, dichondra or desirable clovers. Do not use on floratam, a variety of St. Augustinegrass. Do not use on or around fruits, vegetables or flowers. The maximum number of broadcast applications per treatment site is 2 per year. Do not exceed specified dosages for any area.

#### **HOW IT WORKS**

Because broadleaf weeds and grasses grow differently, Weed-B-Gon® kills broadleaf weeds without harming lawns. Weed-B-Gon singles out weeds, enters through leaves and moves inside the weed to the roots.

#### WEEDS CONTROLLED



White heath aster, bushy aster, heath aster, purple aster, white prairie aster, catchweed bedstraw, marsh bedstraw, northern bedstraw, hairy beggarticks, creeping beggarweed, threeflower beggarweed, Florida beliflower, Florida betony, field bindweed, hairy bittercress, whitehead broom, burclover, common burdock, great burdock, lawn burweed, smallflower buttercup, hairy buttercup, bushy buttonweed, Virginia buttonweed, black medic, Carolina geranium, carpetweed, wild carrot, cat's-ear, smooth chaff-flower, chicory, common chickweed, mouseeared chickweed, sticky chickweed, chicory, common cinquefoil, oldfield cinquefoil, Alyce-clover, crimson clover, hop clover, white clover, wandering cudweed, purple cudweed, narrowleaf cudweed, shiny cudweed, cupids shaving brush, English daisy, oxeye daisy, common dandelion, faise dandelion, catsear dandelion, dayflowers, purple deadnettle, dichondra, dogfennel, doveweed, broadleaf dock, curly dock, dollarweed, doveweed, heartleaf drymary, eclipta, facelis, oakleaf fleabane, rough fleabane, wild garlic, Carolina geranium, clammy ground cherry, smooth ground cherry, ground ivy, groundsel, yellow hawkweed, healalf, henbit, sprawling horseweed, innocense (blue eyed Mary), knawel, prostrate knotweed, tufted knotweed, kochia, lambsquarters, lespedeza, field madder, alkali mallow, Venice mallow, common mallow, little mallow, dwarf mallow, bristly mallow, marcela (yellow joyweed), marestail, horseweed, mat lippia, milkweed, moneywort, mugwort, common mullein, moth mullein, Mexican morningglory, bigroot morningglory, ivyleaf morningglory, tall morningglory, red morningglory, ball mustard, black mustard, blue mustard, indian mustard, tumble mustard, wild mustard, tansy mustard, white mustard, wormseed mustard, bitter nightshade, eastern black nightshade, silverleaf nightshade, cutleaf nightshade, hairy nightshade, wild onion, old world diamond flower, birdseye pearlwort, Florida pellatory, pennywort, Asiatic pennywort, peppergrass, Virginia pepperweed, prostrate pigweed, redroot pigweed, smooth pigweed, tumble pigweed, pineappleweed, broadleaf plantain, buckhorn plantain, blackseeded plantain, poison ivy, poison oak, cutleaf evening primrose, showy evening primrose, purple cudweed, purslane, Brazil pusley, largeflower pusley, Florida pusley, giant ragweed, common ragweed, western ragweed, lanceleaf ragweed, redweed, yellow rocket, field sandbur, longspine sandbur, Mexican or Texas sandbur, southern sandbur, sibara, red sorrel, sheep sorrel, shepardspurse, Pennsylvania smartweed, swamp smartweed, ladysthumb smartweed, pule smartweed, bitter sneezeweed, bilobed speedwell, common speedwell, com speedwell, ivyleaf speedwell, Persian speedwell, purslane speedwell, slender speedwell, snow speedwell, thymeleaf speedwell, winter speedwell, common spiderwort, spotted spurge, prostrate spurge,

(continued

1 6 F

6/6

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

Note to Physician: Emergency information call 1-800-225-2883

# **ENVIRONMENTAL HAZARDS:**

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants.

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target area. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP-p have been associated with mixing/loading and disposal sites.

Caution should be exercised when handling 2,4-D and MCPP-p pesticides at such sites to prevent contamination of groundwater supplies.

Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills.

Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

**NOTICE:** Buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

# The ORTHO Guarantee

If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.



© 2003, Licensed by The ORTHO Group. World rights reserved.

Manufactured for The OATHO Group P.O. Box 190 Marysville, OH 43040

Form LBXXXXXXXXX EPA Reg. No. 239-2665 EPA Est. 239-IA-3<sup>I</sup>, 58996-MO-1<sup>A</sup> Superscript is first letter of lot number Made in USA PRESS TO RESEAL