

239-2666

5-22-2003

115

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 239-2666	2. EPA Product Manager Joanne Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Weed-B-Gon Ready-Spray Isomer Formula	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) The Scotts Company d/b/a The ORTHO Group 14111 Scottslawn Rd Marysville, OH 43041 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

Amendment - Explain below.

Resubmission in response to Agency letter dated _____

Notification - Explain below.

Final printed labels in response to Agency letter dated _____

"Me Too" Application.

Other - Explain below.

NOTIFICATION
MAY 22 2003

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

The ORTHO Group is submitting a notification of "addition of marketing claims" per PR Notice 98-10. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	2. Type of Container <input checked="" type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container

3. Location of Net Contents Information
 Label Container

4. Size(s) Retail Container

5. Location of Label Directions

6. Manner in Which Label is Affixed to Product
 Lithograph Paper glued Stenciled Other _____

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Charles T. Levey	Title Manager, Federal Registrations	Telephone No. (include Area Code) 937-644-7696
--------------------------	---	---

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

2. Signature 	3. Title Manager, Federal Registrations	6. Date Application Received (Stamped)
4. Typed Name Charles T. Levey	5. Date May 16, 2003	

2/5



3.47"

NOTIFICATION
MAY 22 2003

ORTHO®

WEED@GON®
Ready-Spray

**KILLS WEEDS,
NOT LAWNS**

KEEP OUT OF REACH OF CHILDREN
See back panel booklet for additional
precautionary statements.

NET 32 FL OZ (1 QT) 946 mL

FPO

WEEDS CONTROLLED (continued)

mouseeared chickweed, sticky chickweed, chicory, common cinquefoil, oldfield cinquefoil, Alyce-clover, crimson clover, hop clover, white clover, wandering cudweed, purple cudweed, narrowleaf cudweed, shiny cudweed, cupids shaving brush, English daisy, oxeye daisy, common dandelion, false dandelion, catsear dandelion, dayflowers, purple deadnettle, dichondra, dogfennel, doveseed, broadleaf dock, curly dock, dollarweed, doveseed, heartleaf drymary, eclipta, facelis, oakleaf fleabane, rough fleabane, wild garlic, Carolina geranium, clammy ground cherry, smooth ground cherry, ground ivy, groundsel, yellow hawkweed, healall, henbit, sprawling horseweed, innocense (blue eyed Mary), knawel, prostrata knotweed, tufted knotweed, kochia, lambsquarters, lespedeza, field madder, alkali mallow, Venice mallow, common mallow, little mallow, dwarf mallow, bristly mallow, marcela (yellow joyweed), marestail, horseweed, mat lippia, milkweed, moneywort, mugwort, common mullein, moth mullein, Mexican morningglory, bigroot morningglory, ivyleaf morningglory, tall morningglory, red morningglory, ball mustard, black mustard, blue mustard, Indian mustard, tumble mustard, wild mustard, tansy mustard, white mustard, wormseed mustard, bitter nightshade, eastern black nightshade, silverleaf nightshade, cutleaf nightshade, hairy nightshade, wild onion, old world diamond flower, birdseye pearlwort, Florida pellatory, pennywort, Asiatic pennywort, peppergrass, Virginia pepperweed, prostrate pigweed, redroot pigweed, smooth pigweed, tumble pigweed, pineappleweed, broadleaf plantain, buckhorn plantain, blackseeded plantain, poison ivy, poison oak, cutleaf evening primrose, showy evening primrose, purple cudweed, purslane, Brazil pusley, largeflower pusley, Florida pusley, giant ragweed, common ragweed, western ragweed, lanceleaf ragweed, redweed, yellow rocket, field sandbur, longspine sandbur, Mexican or Texas sandbur, southern sandbur, sibara, red sorrel, sheep sorrel, shepardspurse, Pennsylvania smartweed, swamp smartweed, ladythumb smartweed, pale smartweed, bitter snajzeweed, bilobed speedwell, common speedwell, com speedwell, ivyleaf speedwell, Persian speedwell, purslane speedwell, slender speedwell, snow speedwell, thymeleaf speedwell, winter speedwell, common spiderwort, spotted spurge, prostrate spurge, corn spurry, swinecress, bull thistle, Canada thistle, Flodman thistle, cotton thistle, distaff thistle, Drummonds thistle, elk thistle, gray thistle, Italian thistle, leafy thistle, meadow thistle, milk thistle, Platte thistle, plumeless thistle, Russian thistle, Scotch thistle, slenderflower thistle, tall thistle, wavyleaf thistle, yellowspine thistle, stiff verberna, campipoda Veronica, common vetch, hairy vetch, winter vetch, wooley vetch, violet, field violet, wild carrot, wild lettuce, woodsorrel, creeping woodsorrel, yellow woodsorrel, Florida yellow woodsorrel, wooleyleaf bursage, common yarrow, western yarrow, yellow yarrow

STORAGE & DISPOSAL

STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.
DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.



ORTHO WEED & GON
 KILLS WEEDS, NOT LAWNS

OPEN
 Resealable Label
 for Directions &
 Precautions

Makes lawn weed control easy. Simply connect hose to attached sprayer. Eliminates mixing, measuring and messy clean-up. Kills dandelion, clover & other broadleaf weeds. Won't harm lawns – guaranteed! Rain-proof!™ brand. Rain two hours after spraying won't wash away the weed killing power!

PRODUCT FACTS

Treats up to 12,800 sq ft (approximate size of 6 tennis courts)

KILLS WEEDS	Dandelion, Clover, Chickweed plus over 200 additional lawn weeds†	
WHERE TO USE	ON LAWNS without harming lawn grasses	
AMOUNT TO SPRAY	Bentgrass, buffalograss, centipedegrass and St. Augustinegrass (DO NOT USE on floratam) 8 oz (1/4 bottle) treats 3,200 sq ft	
	Bahiagrass, Bermudagrass, fescues, Kentucky bluegrass, perennial ryegrass and zoysiagrass 8 oz (1/4 bottle) treats 1,600 sq ft	

 Questions, Comments or Medical Information?
 Call 1-800-225-2883  www.ortho.com

Specially formulated for residential use.
 For weeds outside the lawn (on driveways, sidewalks, etc.) – use Ortho GroundClear® Complete Vegetation Killer. It kills weeds and grasses and prevents new growth for up to 1 year.

* (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent to 4.38%
 ** 2,4-Dichlorophenoxyacetic acid equivalent to 2.53%
 *** 3,6-dichloro-o-anisic acid equivalent to 1.08%
 Isomer Specific by AOAC Method 6.D01-5.

Made in USA
 The ORTHO Group
 P.O. Box 190
 Mansfield, OH 43040
 EPA Reg. No. 239-2666
 EPA Est. 239-1A-31, 58996-MQ-1A
 Superscript is first letter of lot number
 Made in USA



031403

0 71549 03992 1026

3/5

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

AMOUNT TO SPRAY



- Measure lawn. Calculate square feet by multiplying length times width.
- Use the amount indicated for your lawn variety.
- FOR SENSITIVE LAWNS:** 8 oz (1/4 bottle) treats 3,200 sq ft. Bentgrass, buffalograss, centipedegrass and St. Augustinegrass (DO NOT USE on floratam, a variety of St. Augustinegrass widely grown in Florida)
- FOR HARDY LAWNS:** 8 oz (1/4 bottle) treats 1,600 sq ft. Bahiagrass, Bermudagrass, fescues, Kentucky bluegrass, perennial ryegrass and zoysiagrass

HOW TO USE THE READY-SPRAY SYSTEM

CONNECT



1. Shake well before using.
2. Connect sprayer to hose.
3. Turn on water.

SPRAY

1. To begin spraying, point nozzle in the direction you want to spray.
2. Bend small yellow plastic tab on side of sprayer back and turn knob clockwise to ON position.
3. Spray evenly over measured area.

FINISH

1. To stop spraying, bend small yellow plastic tab back and turn knob counter clockwise to OFF position.
2. Turn off water.
3. Relieve water pressure by bending plastic tab back and turning knob to ON position until water slows to a drip. Then turn knob back to OFF position.
4. Disconnect sprayer from hose.



FOR BEST RESULTS

MOW



Do not mow for 1 to 2 days after application.

WATER



- If soil is dry, water 1 to 2 days before application.
- Rain or watering 2 hours after application will not wash away effectiveness.

WHEN TO APPLY



- **Spray when:**
 - weeds are small and actively growing
 - temperature is below 90°F
 - air is calm to avoid spray drift onto landscape plants as contact with foliage may injure flowers, shrubs, trees and other desirable plants
- Reapplication can be made in 2 to 3 weeks if necessary.
- Newly seeded lawn can be sprayed after first mowing.
- Bare spots can be planted with grass seed 1 week after application.

IMPORTANT

If an ornamental shrub or tree is in the lawn, keep spray off foliage and exposed roots. St. Augustinegrass, centipedegrass or Bermudagrass lawns may have delayed green-up if this product is used when lawns are emerging from dormancy or when temperature drops below 45°F within 72 hours of application. Do not spray carpetgrass, dichondra or desirable clovers. **Do not use on floratam, a variety of St. Augustinegrass.** Do not use on or around fruits, vegetables or flowers. The maximum number of broadcast applications per treatment site is 2 per year. Do not exceed specified dosages for any area.

HOW IT WORKS

Because broadleaf weeds and grasses grow differently, the Weed-B-Gon® LawnGuard™ Formula kills broadleaf weeds without harming lawns. Weed-B-Gon singles out weeds, enters through leaves and moves inside the weed to the roots.

↑ WEEDS CONTROLLED



White heath aster, bushy aster, heath aster, purple aster, white prairie aster, catchweed bedstraw, marsh bedstraw, northern bedstraw, hairy beggarticks, creeping beggarweed, threeflower beggarweed, Florida bellflower, Florida betony, field bindweed, hairy bittercress, whitehead broom, burclover, common burdock, great burdock, lawn burweed, smallflower buttercup, hairy buttercup, bushy buttonweed, Virginia buttonweed, black medic, Carolina geranium, carpetweed, wild carrot, cat's-ear, smooth chaff-flower, chicory, common chickweed,

515

PRECAUTIONARY STATEMENTS



HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. Wear face shield or goggles when mixing, loading or applying this product. When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves. **Personal Hygiene Statements:** After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.



Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried.

FIRST AID



If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **Note to Physician:** Emergency information call 1-800-225-2883.

ENVIRONMENTAL HAZARDS: This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target area. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCP-p have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCP-p pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

NOTICE: Buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

The ORTHO Guarantee

If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.



Questions, Comments or Medical Information?
Call 1-800-225-2883



www.ortho.com

© 2013, Licensed by The ORTHO Group. World rights reserved.

Manufactured for The ORTHO Group

P.O. Box 190

Marietta, OH 43040

Form LBXXXXXXYY

EPA Reg. No. 239-2666

EPA Est. 239-1A-31, 58996-MD-1A

Superscript is first letter of lot number

Made in USA

PRESS TO RESEAL