



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (H7505C)  
401 "M" St., S.W.  
Washington, D.C. 20460

EPA Reg. Number:  
239-2692

Date of Issuance:  
SEP 21 2004

Term of Issuance:  
Conditional

NOTICE OF PESTICIDE:

Registration  
 Reregistration

Name of Pesticide Product:  
Weed-Be-Gone  
Concentrate Formula II

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):  
The Scotts Company d/b/a The Ortho Group  
P.O. Box 190  
Marysville, OH 43040

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- 1. Add the phrase "EPA Registration No. 239-2692" to the label before you release the product for shipment.

COMMENTS CONTINUED ON PAGE 2 OF THIS NOTICE OF REGISTRATION

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product under the enclosed stamped copy of the label constitutes acceptance of these conditions.

Enclosure

Joanne I. Miller  
Product Manager (23)  
Herbicide Branch  
Registration Division (7505C)

Signature of Approving Official:

ISI

Date:

SEP 21 2004

2. The name "Weed-B-Gone Concentrate Formula II" as it appears on your application, EPA Form 8570-1, must be used on the labeling of this product. Use the entire product name where-ever mentioned on the label (Do not shorten to: "Weed-B-Gon®").
  3. Revise the ingredient statement by adding a "Total.....100.00% line following "Other Ingredients.....90.89%".
  4. Under "Product Facts", "Kills Weeds" delete "(many other)". Restrict number of weeds to those listed on this label under "Weeds Controlled" section of this label.
  5. Under the "Amount To Use" correct the dosage for "Southern Lawns (grasses) to read: "3 fl oz (6 Tbs)/gal of water / 1000 sq ft".
  6. In the chart "For Best Results" revise the "Mowing" block to read: [Before and After You Apply" and revise the statement: "Mowing before application is OK." to read: "Mow before application if needed".
  7. Under "Amount To Spray" correct the statement: "(For floratam lawns, use Weed-B-Gon for St. Augustine Lawns)" to read: (Do not use Weed-B-Gon on Floratam variety of St. Augustine lawns". As written the statement does not follow the statement (This product kills floratam grass.).
  8. Delete the line: "EPA Reg. No 239-NEW", and correct the spelling of the word "away" in the "Environmental Hazards" section.
  9. Submit one (1) copy of the final printed labeling before you release this product for shipment.
  10. Submit and/or cite all data required for the registration of this product when the Agency requires all registrants of similar products to submit data; and submit acceptable responses required for reregistration of this product under FIFRA, section 4. A Stamped copy of the label is enclosed for your records.
-

**WEED B GON® Concentrate Formula II**

**Active Ingredients**

- 2,4-D, dimethylamine salt\* .....4.55%
- Mecoprop-p, dimethylamine salt\*\* .....2.30%
- Dichlorprop-p\*\*\* .....2.26%
- Other Ingredients .....90.89%

**Isomer Specific Method Equivalent to:**

- \*2,4—dichlorophenoxyacetic acid 3.8%, 0.32 lbs./gal.  
CAS# 2008-39-1
  - \*\*(+)-R-2-(2-methyl-4-chlorophenoxy)propionic acid 1.9%, 0.16 lbs./gal.  
CAS# 16484-77-8
  - \*\*\*(+)-R-2-(2,4-Dichlorophenoxy)propionic acid 1.9%, 0.16 lbs./gal.  
CAS# 15165-67-0
- Contains the single isomer for of MCPP-p and 2,4-DP-p

For Use On Residential Lawns Only

**KEEP OUT OF REACH OF CHILDREN**

**WARNING** See back panel booklet for additional precautionary statements.

**NET 16 FL OZ (1 PT) 473 mL**  
**32 FL OZ (1 QT) 946 mL**


ACCEPTED  
 with COMMENTS  
 in EPA Letter Dated  
**SEP 21 2004**

Under the Federal Insecticide,  
 Fungicide, and Rodenticide Act  
 as amended, for the pesticide  
 registered under EPA Reg. No.

239-2692

**OPEN**  
**Resealable Label**  
**for Directions & Precautions**  
**PRESS TO RESEAL**

Kills weeds in your lawn - won't harm lawn grasses. Weed-B-Gon singles out weeds, enters through leaves and moves inside the weed to the roots.

<b>PRODUCT FACTS</b>	
Treats up to 8,000 sq ft [pf]	
Treats up to 16,000 sq ft [qf]	
<b>KILLS WEEDS</b>	Dandelion, Clover, Chickweed [illustration] plus (and) over 200 other (many other) (broadleaf) (lawn) weeds*
<b>WHERE TO USE</b>	ON LAWNS [illustration] without harming lawn grasses
<b>AMOUNT TO USE</b>	(For (on) Northern Lawns (grasses)): Fescues, Kentucky bluegrass, annual & perennial rye & zoysia 5 fl oz (10 Tbs) / gal of water / 1000 sq ft
	(For (on) Southern Lawns (grasses)) Bermuda, bahia, bent, buffalo & paspalum 3 fl oz (9 Tbs) / gal of water / 1000 sq ft
((If you are unsure of your lawn type) Use the map to determine which lawn types are most common in your area) [map icon]	
([color] = Fescues, Kentucky bluegrass, annual & perennial rye & zoysia; [color] = Bermuda, bent, buffalo & paspalum; [color] = St. Augustine & centipede)	
[-or-]	
(If you are unsure of your lawn type, contact your local county extension service before applying).	
	<b>Questions, Comments or Medical Information</b> call 1-800-225-2883 <a href="http://www.ortho.com">http://www.ortho.com</a>

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Shake Well Before Use

<b>MIXING INSTRUCTIONS</b>	
<b>(For (on) Northern Lawns (grasses))</b>	
<ul style="list-style-type: none"> <li><u>Fescues, Kentucky bluegrass, annual &amp; perennial rye &amp; zoysia</u>  <b>Amount to Use:</b> When using a tank sprayer, use 5 fl oz (10 Tbs) per 1 gallon of water for each 1000 sq ft  <i>[icon]</i> When using <b>Ortho® Dial 'n Spray®</b>: Set dial to 5 1/3 oz. Add 26 fl oz (52 Tbs) to sprayer jar. <b>DO NOT ADD WATER.</b> Spray evenly over 5,000 sq ft. There's no waste! Any unused product can be poured back into its original container.</li> </ul>	
<b>(For (on) Southern Lawns (grasses))</b>	
<ul style="list-style-type: none"> <li><u>Bermuda, bahia, bent, buffalo &amp; paspalum:</u>  <b>Amount to Use:</b> When using a tank sprayer, use 3 fl oz (6 Tbs) per 1 gallon of water for each 1000 sq ft  <i>[icon]</i> When using <b>Ortho® Dial 'n Spray®</b>: Set dial to 3 oz. Add 15 fl oz (30 Tbs) to sprayer jar. <b>DO NOT ADD WATER.</b> Spray evenly over 5,000 sq ft. There's no waste! Any unused product can be poured back into its original container.</li> </ul>	

1 Tablespoon (Tbs) = 3 teaspoons (tsp)

1 fl oz = 2 Tbs

<b>FOR BEST RESULTS</b>	
<b>HOW TO APPLY</b> <i>[icon / walking man]</i>	<ul style="list-style-type: none"> <li><b>To treat entire lawn (For Broadcast Application):</b> Use a tank or [Ortho® Dial-N-Spray®] hose-end sprayer. Measure lawn. Calculate square feet by multiplying length times width. Spray evenly to lightly wet measured area of lawn.</li> <li><b>For spot treatment:</b> use a tank sprayer: Hold tip of sprayer wand 12 inches from weed, aim at center of weed and spray until lightly wet. To reduce spray drift, adjust sprayer nozzle to deliver a coarse spray.</li> </ul>
<b>MOWING [Before You Apply]</b> <i>[illustration]</i>	<ul style="list-style-type: none"> <li>Do not mow for 1 to 2 days after application. Mowing before application is OK.</li> </ul>
<b>WATERING</b> <i>[illustration]</i>	<ul style="list-style-type: none"> <li>If lawn needs watering, water thoroughly before application.</li> <li>Do not water 48 hours after application. Watering will wash away weed killing material.</li> </ul>
<b>WHEN TO APPLY</b> <i>[illustration]</i>	<ul style="list-style-type: none"> <li>Spray when weeds are small and actively growing.</li> <li>[Spray when] temperature is above 45°F and daytime</li> </ul>

	<p>temperature is below 90°F.</p> <ul style="list-style-type: none"> <li>• [Spray when] air is calm to avoid spray drift onto landscape plants as contact with foliage may injure flowers, shrubs, trees and other desirable plants.</li> <li>• Avoid fine mists.</li> <li>• Reapplication can be made in 2 to 3 weeks if necessary.</li> <li>• In the event further applications are needed, use as a spot treatment.</li> <li>• Newly seeded lawn can be sprayed after mowing 3 times.</li> <li>• Do not reseed for 3 to 4 weeks after use.</li> <li>• (Fall Application – Many weeds germinate in the fall. Using Weed-B-Gon to eliminate fall weeds will help reduce the number of weeds you see the following spring.)</li> </ul>
--	--

**Optional Directions for Hose-End Sprayer)**

<b>AMOUNT TO SPRAY</b>	
<i>[walking man icon]</i>	<ul style="list-style-type: none"> <li>• Measure lawn. Calculate square feet by multiplying length times width.</li> <li>• Use the amount indicated for your lawn variety.</li> </ul>
<b>(For (on) Northern Lawns (grasses))</b>	
<ul style="list-style-type: none"> <li>• <u>Fescues, Kentucky bluegrass, annual &amp; perennial rye &amp; zoysia</u> 16 fl oz (1/2 bottle) treats 3,200 sq ft.</li> </ul>	
<b>(For (on) Southern Lawns (grasses))</b>	
<ul style="list-style-type: none"> <li>• <u>Bermuda, bahia, bent, buffalo &amp; paspalum:</u> 16 fl oz (1/2 bottle) treats 5,333 sq ft.</li> </ul>	
(Floritam is a variety of St. Augustinegrass widely grown in Florida. (This product kills floritam grass.) (For floritam lawns, use Weed-B-Gon for St. Augustine Lawns.)	

**HOW TO USE THE READY-SPRAY® SYSTEM (Hose-End Sprayer)**

<b>CONNECT</b>	<ol style="list-style-type: none"> <li>1. Connect sprayer to hose.</li> <li>2. Turn on water.</li> </ol>
<b>SPRAY</b>	<ol style="list-style-type: none"> <li>3. To begin spraying, point nozzle in the direction you want to spray.</li> <li>4. Bend small (color) plastic tab on side of sprayer back and turn knob clockwise to ON position.</li> <li>5. Spray evenly to lightly wet measured area of lawn.</li> </ol>
<b>FINISH</b>	<ol style="list-style-type: none"> <li>6. To stop spraying, bend small (color) plastic tab back and turn knob counter clockwise to OFF position.</li> <li>7. Turn off water.</li> <li>8. Relieve water pressure by bending plastic tab back and turning knob to ON position until water slows to a drip. Then, turn knob back to OFF position.</li> <li>9. Disconnect sprayer from hose.</li> </ol>

**IMPORTANT**

Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. If an ornamental shrub or tree is in the lawn, keep spray off foliage and exposed roots. Do not spray Carpetgrass, Dichondra or desirable clovers. DO NOT USE ON floratam - a variety of St. Augustinegrass widely grown in Florida. This product kills floratam grass. St. Augustinegrass, Centipedegrass or Bermudagrass lawns may have delayed green-up if this product is used when lawns are emerging from dormancy or when temperature drops below 45° F within 72 hours of application. The maximum number of broadcast applications per treatment site is 2 per year. Do not exceed specified dosages for any area.

**\*WEEDS CONTROLLED [icon]**

Alder, Annual yellow sweet clover, Artichoke, Aster, Austrian fieldcress, Bedstraw, Beggartick, Bident, Bindweed, Bird vetch, Bitterweed, Bitter wintercress, Black-eyed Susan, Black medic, Black mustard, Blackseed plantain, Blackseed plantain, Blessed thistle, Blue lettuce, Blue vervain, Box elder, Bracted plantain, Brassbuttons, Bristly oxtongue, Broadleaf dock, Broadleaf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull nettle, Bull thistle, Burdock, Burning nettle, Bur ragweed, Burweed, Buttercup, Canada thistle, Carolina geranium, Carpetweed, Catchweed bedstraw, Catsear, Catnip, Chickweed, Chicory, Cinquefoil, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common Chickweed, Common mullein, Common sowthistle, Corn Chamomile, Creeping jenny, Crimson clover, Croton, Cudweed, Curly dock, Curly indigo, Dandelion, Dead nettle, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Fall dandelion, False dandelion, False flax, False sunflower, Fiddleneck, Field bindweed, Field pansy, Flea bane (daisy), Flixweed, Florida betony, Florida pusley, Frenchweed, Galinsoga, Garlic mustard, Goathead, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hairy bittercress, Hairy fleabane, Hawkweed, Healall, Heartleaf drymary, Hedge bindweed, Hedge mustard, Hemp, Henbit, Hoary cress, Hoary plantain, Hoary vervain, Honeysuckle, Hop clover, Horsenettle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knawel, Knotweed, Lambsquarter, Lespedeza, Locoweed, Lupine, Mallow, Marshelder, Matchweed, mexicanweed, Milk vetch, Milkweed bloodflower, Mugwort, Morningglory, Mousear chickweed, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf vetch, Nettle, Orange hawkweed, Oxalis, Oxeye daisy, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineywoods bedstraw, Plains coreopsis, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, povertyweed, Prairie, Prickly lettuce, Prickly sida, Primrose, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Puncture vine, Purslane, Ragweed, Red clover, Redroot pigweed, Red sorrel, Redstem filaree, Rough cinquefoil, Rough fleabane, Roundleafed marigold, Rush, Russian pigweed, Russian thistle, St. Johnswort, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherdspurse, Slender plantain, Smallflower galinsoga, Smartweed, Smooth dock, Smooth pigweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spiny, Spiny amaranth, Spiny cocklebur, Spotted catsear, Spotted knapweed, Spotted spurge, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Strawberry clover, Sumac, Sunflower, Sweet clover, Tall nettle, Tall vervain, Tansy mustard, Tansy ragwort, Tanweed, Tarweed, Thistle, Tick trefoil, Toadflax, Trailing Crownvetch, Tumble mustard, Tumble pigweed, Tumbleweed, Velvet leaf, Venice mallow, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Virginia pepperweed, Wavyleaf bullthistle, Western clematis, Western salsify, White clover, White mustard, Wild mustard, Wild aster, Wild buckwheat, Wild carrot, Wild four-o'clock, Wild garlic, Wild geranium, Wild lettuce, Wild marigold, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild vetch, Willow, Witchweed, Woolly morningglory, Woodsorrel, Woolly croton, Woolly plantain, Wormseed, Yarrow, Yellow rocket, Yellow flower pepperweed, and other broadleaf weeds.

**STORAGE AND DISPOSAL [icon]**

**STORAGE:** To be stored in original container and placed in an area inaccessible to children. Keep from freezing.

**DISPOSAL:** **If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** If product cannot be used as directed, or offered to others for their use, call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

**PRECAUTIONARY STATEMENTS [icon]****HAZARDS TO HUMANS & DOMESTIC ANIMALS**

**WARNING:** Causes substantial but temporary eye injury. Harmful if swallowed, inhaled or absorbed through the skin. Do not get in eyes or on clothing. Avoid breathing spray mist and contact with skin. Wear long pants, long-sleeved shirt, socks shoes, rubber gloves and eye protection when mixing, loading or applying this product. It is recommended that safety glasses include front, brow and temple protection. After using this product, wash non-disposable gloves thoroughly with soap and water before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

Do not allow people (other than applicator) or pets on treatment area during application. People and pets may enter treated area after spray has dried.

**FIRST AID:**

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**Note to Physician:** Emergency information call 1-800-225-2883.



**ENVIRONMENTAL HAZARDS:** This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters, nor pour the washwater on the ground; spray or drain over a large area away from wells, other water sources and desirable plants. Do not apply this product through any type of irrigation system. Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D, MCPP-p and 2,4-DP-p have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

**NOTICE:** Buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

**The ORTHO Guarantee**

If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.

**? Questions, Comments or Medical Information ?**  
call 1-800-225-2883      <http://www.ortho.com>

Manufactured for  
The ORTHO Group  
P.O. Box 190  
Marysville, OH 43040  
EPA Reg. No 239-NEW  
EPA Est. 239-IA-3<sup>I</sup>, 58996-MO-1<sup>A</sup>  
Superscript is first letter of lot number  
Made in USA