




198

**ORTHO**  
**Crab Grass Killer**  
**Formula II**



Front Panel

Treats up to  
16,000 Sq. Ft.

Also Kills Dallis Grass, Nut Grass, Goose Grass

**Active Ingredient**

	By Wt.
Calcium acid methanearsonate . . . . .	8.4%
Inert Ingredients . . . . .	91.6%

Total arsenic as elemental

All water soluble - 4.1%

**Keep out of reach of children**

**CAUTION**

See back panel for additional precautionary statements.

**NET CONTENTS 1 GAL.**

10/4/83

Back Panel

### Precautionary Statements

#### Hazards to Humans & Domestic Animals

##### CAUTION:

Causes eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Do not allow children or animals on treated area until after rainfall or when normal irrigation practice is resumed 2 days after treatment. Wear protective clothing when handling or applying this product, including long pants, long-sleeve shirt and rubber gloves and boots. Do not graze treated areas or feed livestock treated forage.

#### STATEMENT OF PRACTICAL TREATMENT

In case of eye contact, wash eyes immediately with fresh water for at least 15 minutes. See a doctor if irritation persists.

**Note to Physicians:** Emergency Information - call (415) 233-3737 (collect).

#### ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning or disposal of waste.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

#### DIRECTIONS

ORTHO Crab Grass Killer Formula II can be used to control Crabgrass, Dallis Grass and Nut Grass on all of the major ornamental turf grasses including Dichondra, Bent and Fescue grasses.

Do not use on St. Augustine, Carpet and Centipede grasses.

**CRABGRASS** (Hairy Crabgrass or Smooth Crabgrass): See Dilution Table for dosages. Soil should be moist before treatment. Do not water for two days after treatment, then resume watering, particularly during hot, dry weather. One application may be sufficient if applied when seedlings are less than 2 inches tall. Two applications four to seven days apart are generally needed for control of medium to mature Crabgrass plants. Follow-up spot sprays may be necessary for newly germinated Crabgrass. Lawns may be reseeded when Crabgrass is dead. On newly-seeded lawns, do not commence applications until after two mowings. Bent, Fescue and Dichondra lawns are more sensitive and may be temporarily discolored. Additional wetting agents are not needed.

**APPLICATION:** Use tank-type sprayer during summer months. After using sprayer with ORTHO Crab Grass Killer Formula II, clean sprayer thoroughly with water before spraying trees, flowers or vegetables. Do not apply with a hose-end sprayer.

**SPECIAL COMBINATION SPRAY:** Use ORTHO Crab Grass Killer Formula II in combination with WEED-B-GON Lawn Weed Killer at recommended rates for both. Use 3 Tablespoonfuls of ORTHO Crab Grass Killer Formula II and 4 teaspoonfuls of WEED-B-GON Lawn Weed Killer per gallon water on 200 sq. ft. area. For larger areas, increase rate proportionally. Avoid drift to ornamental when using this combination or injury may result. Do not use this combination on Dichondra. Observe all precautions, directions and limitations on both product labels.

**DALLIS GRASS** in Bermuda Lawns, Bluegrass, Dichondra and Zoysia Lawns: Use 2 ounces (4 Tablespoonfuls) per gallon of water and cover 200 sq. ft. (20' x 10'). Make 2 or 3 applications at weekly intervals starting in late June. Follow cautions given on label for control of crabgrass.

**Nut Grass** in Bermuda Lawns and Zoysia Lawns: Use 3 ounces (6 Tablespoonfuls) per gallon of water and cover 200 sq. ft. (20' x 10'). Make 3 to 4 applications at monthly intervals starting in June. Follow cautions given on label for control of crabgrass.

---

DILUTION TABLE For Temperatures 85°F and Above

---

ORTHO Crab Grass Killer Formula II to Use	Add water to make	To cover
3 Tablespoonfuls (1½ oz.)	1 gal.	200 sq. ft.
6 Tablespoonfuls (3 oz.)	2 gals.	400 sq. ft.
8 oz. (1 cup)	5 gals.	1,000 sq. ft.
32 oz. (1 qt.)	20 gals.	4,000 sq. ft.

---

For temperatures below 85°F.

Use double the recommended amounts given in above table.

---

**STORAGE AND DISPOSAL**

**STORAGE:** Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry

4

place, preferably in a locked storage area. Do not store diluted spray. Make a new dilution for each use.

**DISPOSAL: PRODUCT** - Partially filled bottle may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash. **CONTAINER** - Do not reuse empty bottle. Rinse thoroughly before discarding in trash.

**NOTICE:** Buyer assumes all responsibility for safety and use not in accordance with directions.

Chevron Chemical Company

Ortho Consumer Products Division

San Francisco CA 94120-7144 Richmond CA 94804-0036

Form 10320-B Product 5517 Made in U.S.A.

EPA Reg. No. 239-NEW

R S M  
EPA Est. 239-CA-1, 239-NJ-1, 239-MO-1

Superscript used in first letter of lot number.

CP #3

*[Faint, illegible handwritten text]*