


50 LBS. NET



10% SEVIN* DUST

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-Methylcarbamate)

Handwritten notes:
340-87
Comments

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-Methylcarbamate) ----- 10%

INERT INGREDIENTS: ----- 90%

TOTAL ----- 100%

Handwritten:
15 lbs per acre

GENERAL DIRECTIONS

Apply SEVIN DUST when first signs of insect infestation are noted, preferably before damage from feeding occurs. For best results, apply thoroughly to upper and lower leaf surfaces. If needed, repeat applications every 7 to 10 days.

DIRECTIONS FOR USE

BEANS: For control of Mexican Bean Beetle/Bean Leaf Beetle/Corn Earworm/Leafhopper/Spotted Cucumber Beetle: Apply 13 to 15 lbs per acre.
POTATOES: For control of Colorado Potato Beetle/Blister Beetle/Flea Beetle/Leafhopper: Apply 13 to 15 lbs. per acre.
SWEET CORN and FIELD CORN: To control Corn Earworm and Fall Army Worm: Apply at rate of 20 lbs. per acre. Begin dusting when first silks appear and repeat at 2 to 3 day intervals. Do not use within 7 days of harvest for forage.
CUCUMBERS, SUMMER SQUASH AND CANTALOUPE: to control Spotted Cucumber Beetles, Squash Beetle, Flea Beetle, Melon Worm and Pickleworm: Apply at rate of 10 lbs. per acre. Thorough coverage is essential. May be applied up to harvest. Make only one application on cucumbers because of possible foliar injury.
TOBACCO: Hornworms/Budworms/Flea Beetles/June Beetles/Japanese Beetles: Apply 10 to 20 lbs. per acre. Good coverage is essential, especially for Flea Beetle control. Pay particular attention to upper side of leaves over entire plant.
COTTON: Boll Weevil/Bollworm/Cotton Fleahopper/Thrips/Cotton Leafworm: At the 8-leaf stage apply 15 lbs. per acre. Repeat at 5-day intervals for 4 applications, July 1 to July 24 application may not be needed. Check field for Bollworm damage or increase of Boll Weevils. On July 24 resume applications at 5-day intervals, at the 15 lb. rate, until squaring stops or all bolls are at least 21 days old. Consult local agricultural authorities for additional dosage and timing.

Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

WARNING
KEEP OUT OF THE REACH OF CHILDREN
CAUTION!

Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Wear regular long sleeved work clothing. Change to clean clothing daily. Avoid contamination of feed and feed stuffs. Avoid spraying near food and food products.
Dispose of waste by burying in non-crop land, away from water supplies. Burn container. Keep out of sight.
Do not contaminate any body of water, by direct application, cleaning of equipment or disposal of wastes and containers.
This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your County Agricultural Extension Service.
Note for Physician: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.
Trade Mark of UCC for Carbaryl (7-naphthyl methylcarbamate) RE: U. S. Patent No. 2,903,478
USDA Reg. No. 240-89

NOTICE OF WARRANTY

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes all risk to persons or property arising out of use or handling and accepts the product on these conditions.

