25\$ SEVIN* DUST

For the Control of Insects on Catton

J. S. Pat. Nos. 2,903,478; 3,909,855

CAUTION

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

Harmful if inhaled or swallowed. Avoid prolonged or repeated breathing of dust. Do not take internally. Skin contact with any insecticide may be harmful. Avoid unnecessary contact. Wash hands and face before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

Note to Physician: Carbaryl is a moderate, reversible, cholinestersse inhibitor. Atropine is antidotal.

AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES.

DIRECTIONS FOR USE

For early-season thrips, flea beetles and cotton fleahopper. Also for cotton leafworm and striped blister beetle: Apply 2 to & pounds of D-H 25% Sevin Dust per acre on appearance of insects.

For Mid and late-season boll weevil, bollworm, fall armyworm, cotton leaf perforator, leaf rollers, leaf hoppers and tarnished plant bug. Also for light to moderate infestation of Lygus bugs in Western irrigated cottons. Apply 4 to pounds of 25% Sevin Dust per acre, depending on the size of the cotton and insect population level. Applications should be made every 5 to 7 days as long as control is necessary.

For stink bugs and salt-marsh caterpillar: Apply 8 lbs. of D-H 25% Sevin Dust per acre as needed.

For grasshoppers: Apply 2 to 6 pounds of D-H 25% Sevin Dust per acre on appearance of insects and their feeding damage. The highest dowage should be used when grasshoppers are mature or material is applied on partly defoliated plants.

Aphid populations will be suppressed by repeated applications of this insecticide.

<u>No. 175,</u>

- 1. Calibrate ground equipment or aerial dust apparatus accurately. After properly adjusting equipment to deliver the require low volume, maintain constant speed throughout application of concentrate Sevin Dust to insure even coverage. Strictly avoid excessive application.
- 2. To effectively apply Concentrate Sevin Dust by aircraft use standard venturi type equipment with spreader vanes or Swathmaster type spreaders. Fly at normal height (4 to 6 feed above the top of the plants is suggested) and maintain regular swath width (usually 30 to 35 feed is indicated). For application with ground equipment use tractor mounted power dusters with a metering device that feeds the Concentrate Sevin Dust through one nossle per row (with deflector) or similarly designed

- 3. For optimum results apply Concentrate Sevin Dust only when weather conditions are favorable. Wind and rising air currents may cause undesirable drift. Strictly avoid excessive dust drift away from crop being treated.
- 4. Sevin does not control spider mites. Where infestations are encountered, use a recommended miticide.
- 5. Sevin may be applied after bolls open. Allow 7 days between last application and grazing treated fields or harvesting crop residue for use as feed or bedding for dairy or meat animals.
- 6. For protection of honeybees avoid use, if possible, during periods when honeybees are visiting the crop or neighboring crops. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance (bey and bee flight range) until one week after apply. **ion.

13POSAL WARNING: Dispose of waste by burning in non-crop lands away from water supplies. Burn bag. Keep out of smoke.

WARRANTY

- 1. Daly-Herring Company guarantees and warrants (a) that the active ingredient and the minimum active ingredient content and the net weight of the contents are as stated within lawful limits and (b) that the directions, warnings, representations and other statements on this label are based upon experts evaluation of tests of effectiveness, of toxicity to plants and to laboratory animals, and of residue remaining on food crops. Tests have not been made on all varieties or in all states.
- 2. Daly-Herring Company makes no other express or implied guarantee, warranty, or representation whatever, of a similar or different sort and NO EXPRESS OR IMPLIED WARRANTY OR MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, and make is to be inferred by a purchaser or user. No modification of this paragraph is authorized, except by specific reference to this paragraph in writing by an employee of the formulator.

U. S. D. A. REG. No. 2110-

Net Weight: 50 pounds

DALY-HERRING COMPANY

Ahoskie, N. C.

Dillon, S. C.

Kinston, N. C.

*Trade mark of Union Carbide Corporation for the active ingredient carbaryl insecticide.