

UNDER THE FEDERAL INSECTICIDE ACT  
 FOR THE YEAR 1977  
 240-197  
 TO AVOID

**WARNING:** May be fatal if swallowed. Do not breath spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly after using. Avoid contamination of food, food containers and feedstuffs.

Do not use this product on dairy animals, in dairy barns or on feed or forage to be fed to dairy animals or other animals being finished for slaughter.

**WARNING:** Do not apply to cucurbits. Do not use treated crops for feed or pasturing of poultry, dairy animals, or animals being finished for slaughter.

Highly toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not contaminate any body of water, by direct application, cleaning of equipment or disposal of wastes and containers nor apply to any area not specified on the label. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not reuse container. Destroy when empty.

Not for use or storage in or around the home

FPA Reg. No. 240-197

**NOTICE OF WARRANTY:** Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

D-4

# 60% Toxaphene

## EMULSIFIABLE CONCENTRATE

Contains 6 lbs. of Toxaphene per gallon

### ACTIVE INGREDIENTS:

Toxaphene - Technical Chlorinated	
Camphene - (Chlorine Content 67-69%)	61.2%
Petroleum Distillate	34.8%
<b>INERT INGREDIENTS:</b>	<b>4.0%</b>

Total 100.0%

### KEEP OUT OF THE REACH OF CHILDREN

See side panel for other warnings.

#### Manufactured By

KINSTON, NORTH CAROLINA 28501

Ahoskie, N. C. 27910      Greensboro, N. C. 27408      Dillon, S. C. 29536  
 Teachey, N. C. 28464

For Aircraft application use a minimum of 3 to 10 gallons of water per acre.

**COTTON:** Cotton Fleahopper, Thrips, Cotton Leafworm: Apply 3 to 4 pints per acre when the first two leaves appear. Follow by another application 7 days later. At the 8-leaf stage apply 2 quarts per acre. Repeat at 5-day intervals for 4 applications. July 1 to July 24 application may not be needed. Check field for Bollworm damage or increase of Boll Weevils. On July 24 resume applications at 5-day intervals, at the 2 quart rate, until squaring stops or all bolls are at least 21 days old.

**CABBAGE:** Diamond Back Moth, Imported Cabbage Worm, Harlequin Bug, Cabbage Aphid: Use 1 to 2 quarts per acre. Do not apply after heads begin to form.

**TOMATOES:** Tomato Hornworm, Flea Beetle, Tomato Fruitworm: Use 1 to 2 quarts in 100 gallons of water per acre. Repeat at 2 week intervals or whenever needed. Do not apply within 5 days of harvest.

**PEANUTS, SOYBEANS:** Armyworm, Velvet Bean Caterpillar, Corn Earworm, Bean Leafbeetle, Grasshoppers: Mix 1 to 2 qts. with sufficient water to give good coverage per acre. Do not feed forage to dairy animals being finished for slaughter.

Do not make more than two applications after pods begin to form. Do not feed treated plants or ensilage made from treated plants to poultry, dairy animals or animals being finished for slaughter. Do not feed soybean mill trash to livestock or poultry.

#### SOYBEANS:

Do not apply within 21 days of harvest or grazing.

#### COTTON:

Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season.