

[241-31]  
1 of 4




(red)



Active Ingredient .....  
Trade Name .....  
EPA Reg. No. ....  
EPA Reg. No. 1-21

(18 pt. type)

(14 pt. type)



19/24/69

CAUTION

[241-31]  
2 of 4

HARMFUL IF SWALLOWED

- Avoid prolonged breathing of spray mist or dust
- Avoid prolonged or repeated contact with skin
- Wash thoroughly after handling
- Do not contaminate food or feed products.

Highly toxic to fish. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes and containers.

Do not use this product for any uses other than those specified on this label.

DIRECTIONS FOR USE

DISCLAIMER

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use.

Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

American Cyanamid Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY.

BUYER assumes the risk of any use contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable by American Cyanamid Company.

Consult your state agricultural experiment station or state agricultural extension service for additional information as to timing, number and rate of applications needed will vary with local conditions.

Unless otherwise indicated, dosages on this label are given in pounds of Malathion 25% Wettable Powder per acre. When applying by ground equipment, use recommended amount in sufficient water for good coverage; by aircraft, apply in 2 to 5 gallons of water. Application should be made when insects appear and repeated as necessary. Do not apply when weather conditions favor drift of spray from areas treated.

VEGETABLES

Pounds  
Per Acre

<u>CRUAS</u> Aphids .....	1
Japanese Beetle .....	5
Do not apply within 1 day of harvest.	

FIELD CROPS

[241-31]  
3 of 4

	<u>Pounds per Acre</u>
<u>PEANUTS:</u> Thrips, Potato Leafhopper .....	4
May be grazed or harvested on day of application.	
<u>SOYBEANS:</u> Mexican Bean Beetle.....	6-8
May be applied on day of harvest or forage use.	
<u>SUGAR BEETS:</u> Aphids, Spider Mite .....	5
Grasshoppers.....	8
Do not apply within 3 days of harvest.	

FRUIT AND NUTS

	<u>Lbs. per 100 gal 1% water</u>
<u>ALMONDS:</u> Aphids, Red Spiders .....	2-2½
Leaf TWIG Borer .....	3-4
Apply during the petal fall period or during the period appropriate for spraying larvae of the May brood. No more than 8 pounds of actual malathion per acre should be applied to almond trees. May be applied on day of harvest.	
<u>CHESTNUTS:</u> Mites.....	2
Apply when mites appear in numbers and 7-10 days later. No more than 5 pounds of actual malathion per acre should be applied to chestnut trees. May be applied on day of harvest.	
<u>FILBERTS:</u> Filbert Aphid, Apple Nymph, Eye-spotted Bud Moth, Scale (crawler stage), Tingids .....	2
No more than 3 pounds of actual malathion per acre should be applied to filbert trees. May be applied on day of harvest.	
<u>MACALANIA NUTS:</u> Green Stink Bug.....	4
Apply when insects start to feed on nuts and repeat as necessary. No more than 15 pounds of actual malathion per acre should be applied to Macalania nut trees. May be applied during harvest.	
<u>PAPAYAS:</u> Aphids, Nymph .....	5
May be applied on day of harvest.	

POULTRY (Chickens, Ducks, Geese, Turkeys)

[241-31]  
4 of 4

Use high-pressure sprayer for surface sprays.  
Avoid contamination of feed, feed troughs and water fountains.

	PESTS	DIRECTIONS
DIRECT APPLICATION	Northern fowl mite Poultry lice Chicken red mite* *(as a supplement to premise treatment for chicken red mite)	Use 1/2 ozs. per gal. of water per 10-150 birds. Apply as a full- coverage spray. Repeat application in 8 weeks or when necessary.
PREMISE TREATMENTS	Northern fowl mite Poultry lice Chicken red mite Flies	Use 1/2 ozs. per gal. of water. Apply liberally to litter, walls, ceilings, roosts, nests and adjacent areas taking care to force the spray into all cracks and crevices. Repeat application as necessary.

Destroy empty bags by burning. Stay away from smoke and fumes.

AMERICAN CYANAMID COMPANY  
AGRICULTURAL DIVISION  
PRINCETON, N. J.

ACCEPTED WITH COMMENTS