

USE CLAIM

THIMET 10-G Soil and Systemic Insecticide
 USDA Reg. No. 241-53

Crop	Pests Controlled	Rate THIMET 10-G	Application	Remarks
COTTON: At Planting Time	Early season thrips, aphids, mites and leaf miners	5 lbs. per acre (40-inch row spacing) or 6 ozs. per 1000 ft. of row (minimum 36-inch spacing)	Distribute the granules evenly in the furrow at planting time. Soil temperature, moisture and tilth should be favorable for good germination and emergence.	Suggested for areas where a limited period of protection against early season pests is necessary for good cotton growth. Use 3 lbs. THIMET 10-G per acre for hill-dropped cotton.
	Mites, thrips, aphids, black cutworms, leaf-hoppers, white flies and leaf miners	10-15 lbs. per acre (40-inch row spacing) or 12-18 ozs. per 1000 ft. of row (minimum 36-inch spacing)	Distribute the granules evenly in the furrow at planting time. Soil temperature, moisture and tilth should be favorable for good germination and emergence. Under adverse conditions plant stand may be affected.	Recommended for areas requiring an extended period of mite control as well as early season protection against other cotton pests. The higher rate is suggested for heavier soils.
COTTON: At Side-Dressing Time (Irrigated cotton only)	Mites from spring to early summer	15-20 lbs. per acre (40-inch row spacing) or 18-24 ozs. per 1000 ft. of row (minimum 36-inch spacing)	Incorporate granules into soil as a side-dress application deep enough to avoid disturbance by future cultivations.	Irrigate as soon as possible after treatment. Do not apply later than 60 days before harvest.

241-53

12/01/71

ACCEPTED

DEC 1 1971

241-53

11