

THIMET 10G Soil and Systemic Insecticide

EPA Reg. No. 241-53

USE CLAIM

CROP	PESTS CONTROLLED	RATE THIMET 10G	APPLICATION	REMARKS
Potatoes	Aphids Leafhoppers Leafminers Psyllids Wireworms Flea beetle larvae Reduction of flea beetle adults	<u>Light or Sandy Soils:</u> 20 lbs. per acre (38-inch row spacing) or 23 ozs. per 1,000 ft. of row for any row spacing (minimum 32-inch spacing) <u>Heavy or Clay Soils:</u> 30 lbs. per acre (38-inch row spacing) or 35 ozs. per 1,000 ft. of row for any row spacing (minimum 32-inch spacing)	Distribute the granules evenly in the furrow or granules may be banded on each side of the row at planting time.	Wait 90 days after treatment before harvesting potatoes.
	Colorado potato beetle (early season control)	<u>Heavy or Clay Soils:</u> 30 lbs. per acre (38-inch row spacing) or 35 ozs. per 1,000 ft. of row for any row spacing (minimum 32-inch spacing)		

ACCEPTED

JAN 11 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 241-53. SUBJECT
TO ATTACHED COMMENTS.

241-53

01/11/1973

1/1