

THIMET 10G Soil and Foliar Insecticide
EPA Reg. 241-53-AA

USE CLAIM

Crop	Pest Controlled	Rate of THIMET 10G	Application	Remarks
FIELD CORN: Application at Planting Time	Corn rootworms Mites	<div style="border: 1px solid black; padding: 5px; text-align: center;"> <p>ACCEPTED</p> <p>MAR 05 1973</p> <p>UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTER- ED UNDER NO. SUBJECT TO ATTACHED COMMENTS.</p> <p><i>241-53</i></p> </div> <p>10 lbs. per acre (40 inch row spacing) or 12 ozs. per 1000 ft. of row for any row spacing (minimum 30-inch row spacing)</p>	Place granules in a 7-inch band over the row directly behind the planter shoe in front of the press wheel.	Do not place THIMET granu in direct contact with so. One additional applicatio can be applied at cultiva or over corn later in the season.
	Seed corn beetle (Iowa and Illinois only)		Apply granules at time of cultivation in a band at base of plants just ahead of cultivator shovels so as to cover granules with soil. Apply in June as soon as rootworm activity is observed.	Do not make any later applications after cultivation treatment.
	Reduction of wireworms (North Central and Southeastern States only)		Application at Cultivation	Corn rootworms
Application Over the Plant	European corn borer (1st brood) (North Central States only)	Corn leaf aphids Mites		Apply granules into the whorl of the plant

241-53

03/05/1973

1/2

USE CLAIM

(page 2)

Crop	Pest Controlled	Rate of THIMET 10G	Application	Remarks
SWEET CORN: Application at Planting Time	Corn Rootworms Mites	10 lbs. per acre (40 inch row spacing) or 12 ozs. per 1000 ft. of row for any row spacing (minimum 30-inch row spacing)	Place granules in a 7-inch band over the row directly behind the planter shoe in front of the press wheel	Do not place THIMET granule in direct contact with seed. One additional treatment can be made at cultivation.
	Seed Corn Beetle (Iowa and Illinois only)			
	Reduction of wire- worms (North Central and Southeastern States only)			
Application at Cultivation	Corn rootworms		Apply granules at time of cultivation in a band at base of plants just ahead of cultivator shovels so as to cover granules with soil. Apply in June as soon as rootworm activity is observed.	Do not make any later applications after cultivation treatment.