

Net Weight: 50 Lbs.

CYPREX® 4% Dust
Fruit Fungicide



Active Ingredient:

Dodine (n-dodecylguanidine acetate) 4.0%
Inert Ingredients 96.0%

EPA Reg. No. 241-62-AA

A FUNGICIDE DUST FORMULATION FOR USE BY GROUND
DUSTER OR AIRPLANE, FOR CONTROL OF APPLE AND PEAR SCAB

CAUTION

KEEP OUT OF REACH OF CHILDREN

CONCENTRATED MATERIAL CAUSES EYE IRRITATION

In case of contact with eyes, flush eyes with plenty of water
for at least fifteen minutes; get medical attention.

PRECAUCION

AL USUARIO: Si usted no lee inglés, no use este producto hasta
que la etiqueta le haya sido explicada ampliamente.
(TO THE USER: If you cannot read English, do not use this product
until the label has been fully explained to you.)

This product is toxic to fish. Keep out of lakes, streams or ponds.
Do not apply when weather conditions favor drift from areas treated.
Do not contaminate water by cleaning of equipment or disposal of
wastes.

Destroy empty bags and cartons by burning. Stay away from smoke
and fumes.

SEE REVERSE SIDE FOR RECOMMENDATIONS FOR USE


DISCLAIMER

American Cyanamid Company warrants only that the material contained
herein conforms to the chemical description on the label and is
reasonably fit for the use therein described when used in accordance
with the directions for use.

Any damages arising from a breach of this warranty shall be limited
to direct damages, and shall not include consequential commercial
damages such as loss of profits or values, etc.

American Cyanamid Company makes no other express or implied warranty,
including any other express or implied warranty of FITNESS or of
MERCHANTABILITY.

BUYER assumes the risk of any use contrary to label instructions, or
under abnormal conditions, or under conditions not reasonably
foreseeable by American Cyanamid Company.



RECOMMENDATIONS FOR USE

Consult your state agricultural experiment station or state agricultural extension service for additional information as the timing, number and rates of applications needed will vary with local conditions.

Care should be taken to avoid application on plums since foliage or fruit injury may result.

Apples: Apple Scab - Apply 25-40 lbs. of CYPREX 4% Dust per acre at 5 to 7-day intervals or as needed to maintain scab control from prebloom through first cover applications. Use 40 lbs. CYPREX 2% Dust per acre in later cover applications.

Do not apply within 5 days of harvest.

Dodine-TREATED APPLES SHOULD NOT BE USED IN THE MANUFACTURE OF APPLE POMACE FOR USE IN LIVESTOCK FEEDS. SUCH USE MAY RESULT IN ILLEGAL RESIDUES IN MILK OR MEAT.

Pears: Pear Scab - Western States Only - Apply 25-40 lbs. of CYPREX 4% Dust per acre at 5 to 7-day intervals or as needed to maintain scab control from prebloom through first cover applications. Continue applications at 10 to 14-day intervals or as needed. Use 40 lbs. of CYPREX 2% Dust per acre in later cover applications.

Do not apply within 5 days of harvest.

DO NOT APPLY IN COMBINATION WITH LIME OR CHLOROBENZILATE.

Do not contaminate food or feed products.

Do not graze cover crops in treated orchards.

In case of an emergency endangering life or property involving this product, call collect, day or night, Area Code 201 - 835-3100.

AMERICAN CYANAMID COMPANY
AGRICULTURAL DIVISION
PRINCETON, N.J. 08540