

Front Panel

Pressurized Pet Spray
Kills Fleas, Ticks, Lice
and other insects

ACTIVE INGREDIENTS:

Malathion
Cyfluthrin
Permethrin
Deltamethrin
Other active ingredients
0.1% of 1,1,1-trichloro-2,2,2-trifluoroethane
petroleum distillates

INERT INGREDIENTS:

0.1% of 1,1,1-trichloro-2,2,2-trifluoroethane
Malathion Premium Grade Technical.
**Consists of 0.4% (butyl carbonyl) (C-propyl) propyl
0.1% other related compounds.

USDA Reg. No. 241-07

(1-gal. type)

CAUTION!

(1-gal. type)

KEEP AWAY FROM CHILDREN

See Below For Additional Cautions and Warnings

Net Contents:

DIRECTIONS FOR USE

Apply spray to the animal from a distance of 6 to 12 inches. Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered including the legs and under the body. As you spray, fluff the hair so that the spray will penetrate into the fur. For cats, spray on the basis of a 4 to 6-second application to a 5-pound animal. For dogs, spray on the basis of a 20-30 second application to a 25-pound animal. Repeat treatments, if necessary, should not be more often than once weekly. Make sure spray wets ticks. Do not spray into the eyes, face or scrotum. Do not overspray.

Thoroughly spray animal's living quarters and bedding; attempting to contact directly as many insects as possible.

Avoid use on birds or around fish bowls.

CAUTION!

Harmful if swallowed. Avoid prolonged or repeated breathing of spray mist. Do not contaminate food or feed products.

WARNING!

Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130° F. may cause bursting. Never throw container into fire or incinerator.

Company Signature
Address

AB:mm
10/5/66