



---

## CAUTION!

THIS PRODUCT IS HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN

**Avoid breathing dust or spray mist**

**Avoid contact with skin**

**Wash thoroughly after handling**

**Do not contaminate food or feed products.**

---

**WARNING:** This product may cause cholinesterase inhibition. Atropine is antidotal.

Pralidoxime chloride (2-PAM, PROTOPAM chloride) may be effective as an adjunct to atropine. Use according to label directions.

**This product is toxic to birds and other wildlife. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.**

This product is toxic to bees. Restrict application to the period after dark when bees are inside the hives or in the early morning before the bees are foraging in the fields. Protective information may be obtained from your cooperative agricultural extension service.

**BAG DISPOSAL:** Destroy empty bags by burning. Stay away from smoke and fumes.

### DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above.

Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values or any other special or indirect damages.

materials, or the manner of use or application all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above.

Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values or any other special or indirect damages.

American Cyanamid Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY.

#### DIRECTIONS FOR USE

Do not use this product for any uses other than those specified on this label. Repeat applications as necessary. Consult your state experiment station or state extension service for proper timing of applications.

CROP	PESTS CONTROLLED	RATE	INTERVAL (DAYS) Between Last Application and Harvest
Apples	Apple maggot† Codling moth*†	2 lbs./100 gallons of water	28 Do not apply during the bloom period. Apply at petal-fall and every 10 to 14 days thereafter until control is achieved. †Under heavy infestations some sting injury may occur. *Midwest and eastern states only.
Apples Pears	Aphids, Leafhoppers, Mites (except rust mites)	1 to 2 lbs./100 gals. water	28 Do not apply during the bloom period.
Grapes (Grown in California)  Raisin and Wine Grapes  Table and Eating Grapes	Grape leafhopper  Pacific spider mite  Thrips	Ground: 6-8 lbs./acre. Apply through concentrate or dilute spray equipment in 20-400 gallons of water per acre depending upon vine growth density. Repeat applications as necessary.  Air: 8 lbs./acre. Apply in 10-30 gallons of water per acre. Repeat applications as necessary.	28