


241-145

10/04/1973

1/2



ACCEPTED
 4 OCT 1973
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC PESTICIDE REGISTER-
 ED UNDER NO. 241-145 SUBJECT
 TO ATTACHED COMMENTS.



THIMET® 15-G

SOIL AND SYSTEMIC INSECTICIDE FOR USE ON CORN

CYANAMID

Active Ingredient:

Phorate (O,O-diethyl S-[(ethylthio) methyl] phosphorodithioate)..... 15.0%

Inert Ingredients..... 85.0%

EPA Reg. No. 241-145-AA
ORGANIC PHOSPHATE COMPOUND MIXTURE, DRY N.O.S.

DANGER!

KEEP OUT OF REACH OF CHILDREN



POISON

See Back Panel For Antidote And Other Warnings

AL USUARIO: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

Net Weight: 15 Lbs.

24641-05

DS4

DANGER!

POISONOUS BY SKIN CONTACT, INHALATION, OR SWALLOWING

RAPIDLY ABSORBED THROUGH SKIN

REPEATED INHALATION OR SKIN CONTACT MAY, WITHOUT SYMPTOMS,

PROGRESSIVELY INCREASE SUSCEPTIBILITY TO POISONING

DO NOT GET IN EYES, ON SKIN, ON CLOTHING. Wear freshly laundered, long-sleeved work clothing daily. While transferring from package to equipment, wear a clean cap and gloves (rubber or cotton). If cotton gloves are used, they must be laundered or discarded after each day's use. Rubber gloves should be washed with soap and water after each use. Do not wear the same gloves for other work. Destroy and replace gloves frequently.

In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and water. Launder clothing before reuse. Wash thoroughly with soap and water before eating or smoking. Bathe at the end of the work day and change outer clothing.

DO NOT BREATHE DUST. While emptying bags into equipment, pour downwind and allow as little free fall as possible. Do not pour at face level and do not allow dust to reach the breathing zone.

DO NOT CONTAMINATE FOOD OR FEED PRODUCTS. Sweep up and bury spillage, whether it occurs indoors or in the field. Once a bag has been opened, use it completely or bury the remainder. Make sure that the hoppers of equipment are emptied while still in the field. Keep all unprotected persons out of operating areas.

KEEP OUT OF REACH OF DOMESTIC ANIMALS.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME

Destroy empty bags and cartons by burning. Stay away from smoke and fumes.

This product is toxic to fish, birds and other wildlife. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DISCLAIMER

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use.

Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

American Cyanamid Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY.

BUYER assumes the risk of any use contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable by American Cyanamid Company.

DIRECTIONS FOR USE

IMPORTANT! READ PRECAUTIONARY STATEMENTS BEFORE USING THIS PRODUCT.

Do not use this product for any uses other than those specified on this label.

THIMET 15-G should be applied with a granular pesticide applicator properly calibrated to assure accurate placement and proper dosage.

Fields may be re-entered on day of treatment with soil applications of THIMET 15-G. Do not enter fields within 7 days after plants have been treated with foliar applications.

Crop	Pest Controlled	Rate of THIMET 15G	Application	Remarks
SWEET CORN: Application at Planting Time	Corn Rootworms Mites	6.5 lbs. per acre (40 inch row spacing) or 8 ozs. per row spacing (minimum 30-inch row spacing)	Place granules in a 7-inch band over the row directly behind the planter shoe in front of the press wheel.	Do not place THIMET granules in direct contact with seed. One additional treatment can be made at cultivation.
	Seed Corn Beetle (Iowa and Illinois only)			
	Reduction of wireworms (North Central States only)			
Application at Cultivation	Corn rootworms		Apply granules at time of cultivation in a band at base of plants just ahead of cultivator shovels so as to cover granules with soil. Apply in June as soon as rootworm activity is observed.	Do not make any later applications after cultivation treatment.
FIELD CORN: Application at Planting Time	Corn rootworms Mites	6.5 lbs. per acre (40 inch row spacing) or 8 ozs. per 1000 ft. of row for any row spacing (minimum 30-inch row spacing)	Place granules in a 7-inch band over the row directly behind the planter shoe in front of the press wheel.	Do not place THIMET granules in direct contact with seed. One additional application can be applied at cultivation or over corn later in the season.
	Seed corn beetle (Iowa and Illinois only)			
	Reduction of wireworms (North Central States only)			
Application at Cultivation	Corn rootworms		Apply granules at time of cultivation in a band at base of plants just ahead of cultivator shovels so as to cover granules with soil. Apply in June as soon as rootworm activity is observed.	Do not make any later applications after cultivation treatment.
Application Over the Plant	European corn borer (1st brood) (North Central States only)	6.5 lbs. per acre (40 inch row spacing) or 8 ozs. per 1000 ft. of row for any row spacing (minimum 30-inch row spacing)	Apply granules into the whorl of the plant prior to tassel emergence.	Do not graze or cut for forage within 30 days of treatment. Do not make more than one application over the plant. Consult your State Experiment Station or State Extension Service for proper timing of application.
	Corn leaf aphids Mites		Apply granules into the whorl of the plant.	



POISON



IN CASE OF AN EMERGENCY ENDANGERING LIFE OR PROPERTY INVOLVING THIS PRODUCT, CALL COLLECT, DAY OR NIGHT, AREA CODE 201-835-3100.

ANTIDOTE: Atropine is an antidote. Consult your physician about obtaining a supply of 1/100 grain (0.6 milligram) tablets for emergency use. If symptoms include blurred vision, stomach cramps or tightness in chest, don't wait for a physician but take two tablets at once. Do not take atropine unless symptoms of poisoning have occurred. Anyone who has been sick enough to have taken atropine must be seen by a physician as soon as possible.

FIRST AID

IF SWALLOWED, induce vomiting by sticking finger down throat or by giving soapy or strong salty water to drink. Repeat until vomit is clear. Never give anything by mouth to an unconscious person.

IF INHALED, remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. If breathing is difficult, give oxygen. IN CASE OF CONTACT, immediately flush eyes or skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. Wash clothing before reuse.

CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING

NOTE TO PHYSICIANS: Warning symptoms include weakness, headache, tightness in chest, blurred vision, nonreactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps. Give atropine intramuscularly or intravenously, depending on severity of poisoning, 2 to 4 milligrams (3 to 6 tablets, 1/100 grain each) every 15 minutes until fully atropinized as shown by dilated pupils, dry flushed skin and tachycardia. Twenty to thirty milligrams, or more, may be required during the first 24 hours. Never give opiates or phenothiazine tranquilizers. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously for at least 48 hours. Allow no further exposure to any cholinesterase inhibitor until cholinesterase regeneration has taken place as determined by blood tests. Pralidoxime chloride (2-PAM; PROTOPAM chloride) may be effective as an adjunct to atropine. Use according to label directions.

© Trademark
Lot No.

Sold By
AMERICAN CYANAMID COMPANY
AGRICULTURAL DIVISION
PRINCETON, N. J. 08540

Made and Printed in U.S.A.