

PM16

241-145

1 of 15

Bag Label  
Front Panel

<b>ACCEPTED</b>
<b>JAN 7 1985</b>
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. <b>241-145</b>

**THIMET® 15-G**

soil and systemic insecticide

**Active Ingredient:**

Phorate (0,0-diethyl S-[(ethylthio) methyl]phosphorodithioate) . . . 15.0%

Inert Ingredients . . . . . 80.0%

EPA Reg. No. 241-145-AA

EPA Est. No.

(12 pt. type)

**KEEP OUT OF REACH OF CHILDREN**

**POISON**



**DANGER**

**PELIGRO!**

**PRECAUCION AL USUARIO:** Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

**FIRST AID**

**If swallowed, drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Avoid alcohol. Get medical attention.**

**If inhaled, remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. If breathing is difficult, give oxygen. Get medical attention.**

**If on skin, wash thoroughly with soap and water. Remove contaminated clothing and shoes. Wash clothing and decontaminate shoes before reuse.**

**If in eyes, immediately flush eyes with plenty of water. Get medical attention.**

**Antidote: Atropine is an antidote.**

PM16

241-145

2 of 15

-2-

Bag Label  
Front Panel (Cont'd.)

**NOTE TO PHYSICIANS:** Warning symptoms include weakness, headache, tightness in chest, blurred vision, nonreactive pinpoint pupils, salivation, nausea, vomiting, diarrhea, and abdominal cramps. Give atropine intramuscularly or intravenously, depending on severity of poisoning. 2 to 4 milligrams every 10 minutes until fully atropinized as shown by dilated pupils, dry flushed skin and tachycardia. Twenty to thirty milligrams, or more, may be required during the first 24 hours. Never give opiates or phenothiazine tranquilizers. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously for at least 48 hours. Allow no further exposure to any cholinesterase inhibitor until cholinesterase regeneration has taken place as determined by blood tests.

Pralidoxime chloride (2-PAM; PROTOPAM chloride) may be effective as an adjunct to atropine. Use according to label directions.

See Back Panel for Additional Precautionary Statements.

Net Weight: 50 lbs.  
22.68 kg.

24638-20

D46

● Registered trademark of American Cyanamid Company

**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS****DANGER!**

Fatal if swallowed, inhaled or absorbed through the skin. Do not breathe dust. Do not get in eyes, on skin, or on clothing.

Wear freshly laundered, long-sleeved work clothing daily. While transferring from package to equipment, wear a clean cap, gloves (rubber or cotton) and goggles. If cotton gloves are used, they must be laundered or discarded after each day's use. Rubber gloves should be washed with soap and water after each use. Do not wear the same gloves for other work. Destroy and replace gloves frequently.

In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and water. Launder clothing and decontaminate shoes before reuse. Wash thoroughly with soap and water before eating or smoking. Bathe at the end of the work day and change clothing.

**Do Not Breathe Dust**

Wear a face mask or other respiratory equipment while emptying bags of product into a hopper. While emptying bags into equipment, pour downwind and allow as little free fall as possible. Do not pour at face level and do not allow dust to reach the breathing zone.

**Do Not Contaminate Food or Feed Products**

Once a bag has been opened, use it completely. Make sure the hoppers are emptied while still in the field. Refer to STORAGE AND DISPOSAL statement for further instructions.

**Keep all unprotected persons out of operating areas.**

Do not apply this product in such a manner as to directly or through drift expose workers or other persons.

**Keep Out of Reach of Domestic Animals.****Not for Use or Store In or Around the Home.****RE-ENTRY STATEMENT**

Do not enter treated areas without protective clothing until treatments have been completed.

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: **DANGER.** Area treated with THIMET on (date of application). Do not enter without protective clothing.

Fields may be re-entered on day of treatment with soil applications of THIMET 15-G. Do not enter corn fields within 7 days after plants have been treated with foliar applications.

### ENVIRONMENTAL HAZARDS

Product is toxic to fish, shrimp, crab, birds and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and crab may be killed at application rate recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label.

This pesticide is toxic to bees exposed to direct application. Applications should be timed to coincide with periods of minimum bee activity, usually between late evening and early morning.

### STORAGE AND DISPOSAL

#### Storage

Store pesticide products in a secure locked area where children, unauthorized persons and animals cannot enter. Do not store in the same area with food or feed. Do not store opened bags.

#### Prohibitions

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

#### Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or Environmental Control Agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance.

#### Container Disposal

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

#### General

Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

PM16 . . .

241-145

5 of 15

-3-

**CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING.**

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This label must be in the possession of the user at the time of pesticide application.

**BEFORE USING, READ PRECAUTIONARY STATEMENTS.**

THIMET should be applied with a granular pesticide applicator properly calibrated to assure accurate placement and proper dosage. Cover granules that may be exposed on the ends of the treated rows and turns and loading areas by deep discing immediately after treating fields.

CROPS

Crop	Pests Controlled	Rate of THIMET 15-G	Application	
ALFALFA	Alfalfa weevil larvae	10-13 lbs. per acre	Broadcast the granules evenly over the field in the spring when first growth starts.	Do 35 c A to may infe
BARLEY  (Irrigated Barley only)	Aphids	6.5 lbs. per acre	Broadcast the granules evenly over the field when aphids first appear using air or ground equipment.	Irrig trea with Do graz
BEANS  At Planting	Mexican bean beetles	6-12.5 ozs. per 1,000 ft. of row for any row spacing (minimum 30-inch spacing).	Drill granules to the side of the seed.	Do gran con
	Leafhoppers Aphids Lygus bugs Thrips Mites Seedcorn maggots	6-9.3 ozs. per 1,000 ft. of row for any row spacing (minimum 30-inch spacing)		Do trea 60 c
	Seedcorn maggots		Apply granules in a band over the row.	

CROPS (Continued)

PM16

Crop	Pests Controlled	Rate of THIMET 15-G	Application	Remarks
<p>FIELD CORN AND SWEET CORN (excluding popcorn)</p> <p>At Planting</p>	<p>Corn rootworms Wireworms White grubs Seedcorn maggots Seedcorn beetles Flea beetles Mites</p>	<p>8 ozs. per 1,000 ft. of row for any row spacing (minimum 30-inch row spacing)</p>	<p>Place granules in a 7-inch band over the row directly behind the planter shoe in front of or behind the press wheel.</p>	<p>Do not place THIMET granules in direct contact with seed. One additional application can be applied at cultivation OR over corn later in the season.</p>
<p>FIELD CORN AND SWEET CORN (excluding popcorn)</p> <p>At Cultivation</p>	<p>Corn rootworms</p>		<p>Apply granules to base of plants or over the top of plants just ahead of cultivation shovels so as to cover granules with soil.</p>	<p>Do not make any applications of THIMET after cultivation treatment.</p>
<p>FIELD CORN (excluding sweet and popcorn)</p> <p>At Cultivation</p>	<p>Chinch bug nymphs (Colorado, Kansas, and Nebraska)</p>		<p>Apply granules to base of plants or over the top of plants just ahead of cultivation shovels so as to cover granules with soil. Apply as soon as chinch bug nymphs are observed at base of plants. Granules must be placed at the infested region of the plants and adequately covered with soil.</p>	<p>Later infestations of chinch bugs may not be controlled. Do not make any applications of THIMET after cultivation treatment.</p>

SHI-145

7815

CROPS (Continued)

PM 16

Crop	Pests Controlled	Rate of THIMET 15-G	Application	Remarks
<p>FIELD CORN (excluding sweet and popcorn)</p> <p>Over the Plant</p>	<p>European corn borer (1st brood)</p>	<p>6.7 lbs. per acre</p>	<p>Apply granules into the whorl of the plant prior to tassel emergence.</p>	<p>Do not graze or cut for forage within 30 days of treatment. Do not make more than one application over the plant. Do not apply under prolonged drought conditions. Consult your state experiment or state extension service for proper timing of application.</p>
	<p>Corn leaf aphids Mites</p>		<p>Broadcast the granules evenly over the top of plants by air or ground equipment.</p>	
<p>COTTON</p> <p>At Planting</p>	<p>Early season control of: Thrips Mites Leaf miners</p>	<p>4 ozs. per 1,000 ft. of row for any row spacing (minimum 36-inch spacing)</p>	<p>Distribute the granules evenly in the furrow at planting time. Soil temperature, moisture and tilth should be favorable for good germination and emergence.</p>	<p>Suggested for areas where a limited period of protection against early season pests is necessary for good cotton growth. Use 1.5 lbs. THIMET 20-G per acre for hill-dropped cotton.</p>
	<p>Mites Thrips Aphids Black cutworms Leafhoppers White flies Leaf miners</p>	<p>8-12 ozs. per 1,000 ft. of row for any row spacing (minimum 36-inch spacing)</p>	<p>Distribute the granules evenly in the furrow at planting time. Soil temperature, moisture and tilth should be favorable for good germination and emergence. Under adverse conditions plant stand may be affected.</p>	<p>Recommended for areas requiring an extended period of mite control as well as early season protection against other cotton pests. The higher rate is suggested for heavier soils.</p>

241-145

8915

CROPS (Continued)

PM16

Crop	Pests Controlled	Rate of THIMET 15-G	Application	Remarks
COTTON Side-Dressing (Irrigated Cotton Only)	Mites from spring to early summer	12-16 ozs. per 1,000 ft. of row for any row spacing (minimum 36-inch spacing)	Incorporate granules into the soil as a side-dress appli- cation deep enough to avoid disturbance by future cultivations.	Irrigate as soon as possible after treatment. Do not apply later than 60 days before harvest.
HOPS	Mites Aphids	13-20 lbs. per acre	Band over the row after crowns have begun to develop or shank in 2-4 inches deep.	Do not apply within 42 days of harvest. Do not feed crop refuse to livestock. Banded applications must be hilled over or watered in.
LETTUCE (Grown in the Northeastern United States)	Aster leafhoppers for reduction of lettuce yellows	4 ozs. per 1,000 ft. of row for any row spacing (minimum 18-inch spacing)	Apply the granules in the fertilizer band on one or both sides of the row at seeding time or when plants are transplanted in the field.	Do not place THIMET granule in direct contact with the seed. For best results seed should be treated with a registered fungicide in accordance with the recommendations on the label.
PEANUTS At Planting	Thrips Leafhoppers	7.3 ozs. per 1,000 ft. of row for any row spacing (minimum 24-inch spacing)	Distribute the granules evenly in the furrow at planting time. For best results, plant peanuts when soil temperatures are favorable for establishing good peanut stands.	Do not graze or feed treated hay or forage.

241-145

9815

CROPS (Continued)

Crop	Pests Controlled	Rate of THIMET 15-G	Application	Remarks
<p>PEANUTS At Pegging</p>	<p>Southern corn rootworms and Leafhoppers</p>	<p>14.6 ozs. per 1,000 ft. of row for any row spacing (minimum 24-inch spacing)</p>	<p>Distribute the granules as a band over the fruiting zone at pegging time. Work into the top few inches of soil immediately.</p>	<p>Do not graze or feed treated hay or forage.</p>
<p>POTATOES At Planting</p>	<p>Aphids Leafhoppers Leaf miners Psyllids Wireworms Flea beetle larvae and reduction of flea beetle adults</p>	<p>Light or Sandy Soils: 15 ozs. per 1,000 ft. of row for any row spacing (minimum 32-inch spacing)</p> <p>Heavy or Clay Soils: 23 ozs. per 1,000 ft. of row for any row spacing (minimum 32-inch spacing)</p>	<p>Distribute the granules evenly in the furrow or band on each side of the row.</p>	<p>Wait 90 days after treatment before harvesting potatoes.</p> <p>Do not use for Colorado potato beetle control in the Northeast United States.</p>
	<p>Colorado potato beetle (early season control)</p>	<p>Heavy or Clay Soils: 23 ozs. per 1,000 ft. of row for any row spacing (minimum 32-inch spacing)</p>		

PM 16.

241-145

100215

CROPS

PM16

Crop	Pests Controlled	Rate of THIMET 15-G	Application	Remarks
SORGHUM At Planting	Greenbugs	8 ozs. per 1,000 ft. of row for any row spacing (minimum 30-inch spacing)	Place granules in a 7-inch band over the row OR drill 1 inch below and 2 inches to the side of the seed.	Do not place THIMET granules in direct contact with seed. A second application of granules may be applied topically later in the season.
SORGHUM At Cultivation	Chinch bug nymphs (Colorado, Kansas, Nebraska)		Apply granules to base of plants or over the top of plants just ahead of cultivation shovels so as to cover granules with soil. Apply as soon as chinch bug nymphs are observed at base of plants. Granules must be placed at the infested region of the plants and adequately covered with soil.	Later infestations of chinch bugs may not be controlled.  Do not make any application of THIMET after cultivation treatment.  Do not feed foliage before grain harvest.
SORGHUM Over the Plant	Greenbugs Banks grass mites	6.5 lbs. per acre	Broadcast granules into the whorls of the plants by air or ground equipment.	Make only one application after plant emergence. Do not apply within 28 days of harvest.

841-145

11815

CROPS

Crop	Pests Controlled	Rate of THIMET 15-G	Application	Remarks
SOYBEANS At Planting	Mexican bean beetles Leafhoppers Lygus bugs Thrips Aphids Mites Seedcorn maggots	12 ozs. per 1,000 ft. of row for any row spacing (minimum 30-inch row spacing)	Drill granules to the side of the seed.	Do not place THIMET granules in direct contact with seed. Do not feed the foliage of treated soybeans. Crop injury may result when preplant incorporated or preemergence applications of metribuzine herbicides are used in conjunction with this product or other soil-applied organophosphate pesticides.
	Seedcorn maggots		Place granules in a 7-inch band over the row directly behind the planter shoe in front of the press wheel.	
SUGAR BEETS At Planting	Aphids Beet root maggots Leafhoppers Mites Leaf miners	6 ozs. per 1,000 ft. of row for any row spacing (minimum 20-inch spacing)	Drill granules to the side of the seed OR place granules in a band over the row.	Do not place THIMET granules in direct contact with the seed. Do not apply within 30 days of harvest. Do not feed sugar beet tops or silage to dairy cattle. Consult your state experiment station or state extension service for proper timing of application in your area.
SUGAR BEETS Postemergence	Mites Aphids	6.5-10 lbs. per acre	Apply to foliage when plants are dry. Use the higher rate on dense foliage for good coverage. Foliar application by air may be used for aphid control when wet soil prohibits ground application.	

PM 16

241-145

12/15

Crop	Pests Controlled	Rate of THIMET 15-G	Application	Remarks
SUGAR CANE (Florida)	Wireworms	26 lbs. per acre	Apply in a 10-12 inch band directly on the seed piece and surrounding soil in the open furrow immediately before covering.	
TOMATOES (Grown in the Southeastern United States)	Leaf miners	15 ozs. per 1,000 ft. of row for any row spacing (minimum 38-inch row spacing)	Apply the granules in the fertilizer band on either side of row at seeding time or when plants are transplanted in the field.	Do not apply THIMET granules in direct contact with the seed.
WHEAT At Planting	Grasshoppers Aphids (Western and North Central States only) Hessian fly	1.6 ozs. per 1,000 ft. or row for any row spacing (minimum 8-inch spacing)	Apply the granules in the seed furrow.	Do not feed or graze foliage within 45 days of treatment. Do not make any later applications after planting time treatment.
WHEAT Over the Plant	Aphids (Western States only)	6.5 lbs. per acre	Broadcast the granules evenly over the field by air equipment when aphids first appear.	Do not apply over plants if application was made at planting time. Do not make more than one application over the plants. Do not apply within 70 days of harvest of grain. Do not feed or graze foliage within 28 days of treatment.

PM 16

241-145

12415

PM16

### TYPICAL APPLICATION RATES FOR VARIOUS ROW SPACINGS

Rate Oz./1,000 Ft. of Row

	1.6 oz.	2 oz.	4 oz.	6 oz.	8 oz.	10 oz.	12 oz.	14 oz.	16 oz.	18 oz.	20 oz.	22 oz.	24 oz.
40"				4.9	6.5	8.2	9.8	11.4	13.1	14.7	16.3	18.0	19.6
38"				5.2	6.9	8.6	10.3	12.0	13.7	15.5	17.2	18.9	20.6
36"				5.4	7.3	9.1	10.9	12.7	14.5	16.3	18.2	20.0	21.8
34"				5.6	7.7	9.7	11.6	13.5	15.4	17.4	19.3	21.3	23.2
32"				6.1	8.2	10.2	12.3	14.3	16.3	18.3	20.5	22.4	24.6
30"				6.5	8.7	10.9	13.1	15.2	17.4	19.6	21.8	24.0	26.1
26"				7.1	9.5	11.8	14.2	16.6	18.9	21.3	23.7	26.0	28.4
24"			5.4	8.2	10.9	13.6	16.3	19.1	21.8	24.5	27.2	30.0	32.7
22"			5.9	8.9	11.9	14.9	17.9	21.0	24.0	26.9	29.9	33.0	35.9
20"			6.5	9.8	13.1	16.3	19.6	22.8	26.1	29.3	32.6	35.9	39.1
16"			8.2	12.3	16.3	20.5	24.6	28.7	32.7	36.8	40.9		
12"		5.4	10.9	16.3	21.8	27.2	32.7	38.1	43.6				
8"	6.5	8.2	16.3	24.6	32.7	40.9							
6"	8.7	10.9	21.8	32.7	43.6								

51-145

EQUIVALENT POUNDS PER ACRE

14815

POISON

6

DOT-E6296

ORGANIC PHOSPHATE COMPOUND  
MIXTURE, DRY

THIMET<sup>®</sup>  
soil and systemic insecticide