


DIRECTIONS FOR USE IN PEANUTS

**BEFORE USING, READ PRECAUTIONARY
STATEMENTS ON BAG LABEL**



It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. THIMET 15-G should be applied with a granular pesticide applicator properly calibrated to assure placement and proper dosage. See label for specific instructions.

Cover granules that may be exposed on the ends of the treated rows, turns and loading areas by deep disking immediately after treating fields.

CROP	PESTS CONTROLLED	RATE OF THIMET 15-G	APPLICATION	REMARKS
PEANUTS At Planting	Thrips Leafhoppers	7.3 ozs. per 1000 ft. of row for any row spacing. (minimum 24-inch spacing)	Distribute the granules evenly in the furrow at planting time. For best results, plant peanuts when soil temperatures are favorable for establishing good peanut stands.	Do not graze or feed treated hay or forage.
PEANUTS At Pegging	Southern corn rootworms Leafhoppers Suppression of: white mold <i>Sclerotium rolfsii</i>	14.6 ozs. per 1000 ft. of row for any row spacing. (minimum 24-inch spacing)	Distribute the granules as a band over the fruiting zone at pegging time. Work into the top few inches of soil immediately.	Do not graze or feed hay or treated forage.

**THIS LABEL SHOULD BE IN THE POSSESSION OF THE
USER AT THE TIME OF PESTICIDE APPLICATION**



Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470 © 1984

-3-
A stamped copy of the supplemental labeling is enclosed for your records.

Sincerely yours,

William E. Miller
Product Manager (16)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosure

SUPPLEMENTAL LABELING

**ACCEPTED
with COMMENTS
in EPA Letter Dated:**

NOV 5 1987

THIMET^(R) 15-G soil and systemic insecticide

EPA Reg No. 241-145-AA

Under the Fungicide, Insecticide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

241-145

BEFORE USING, READ THE PRECAUTIONARY STATEMENTS ON THE BAG

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. THIMET 15-G should be applied with a granular pesticide applicator properly calibrated to assure placement and proper dosage. See label for specific instructions.

Cover granules that may be exposed on the ends of the treated rows, turns and loading areas by deep discing immediately after treating fields.

Crop	Pests Controlled	Rate of THIMET 15-G	Application
FIELD CORN AND SWEET CORN (excluding popcorn) At Planting	Corn Rootworms Wireworms White grubs Seedcorn maggots Seedcorn beetles Flea beetles Mites	8 ozs per 1,000 ft of row for any row spacing (minimum 30-inch row spacing)	Place granules in a 7- inch band over the row directly behind the planter shoe in front of or behind the press wheel and lightly incorporate.

Remarks: Do Not use in-furrow application. One additional application can be applied at cultivation or over corn later in the season.

^(R)Registered Trademark of American Cyanamid Company



American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470

